



HAL
open science

Impaired interleukin-1 β and c-Fos expression in the hippocampus is associated with a spatial memory deficit in P2X(7) receptor-deficient mice

Virginie Labrousse, Laurence Costes, Agnès Aubert, Muriel Darnaudéry, Guillaume Ferreira, Thierry Amédée, Sophie Layé

► To cite this version:

Virginie Labrousse, Laurence Costes, Agnès Aubert, Muriel Darnaudéry, Guillaume Ferreira, et al.. Impaired interleukin-1 β and c-Fos expression in the hippocampus is associated with a spatial memory deficit in P2X(7) receptor-deficient mice. PLoS ONE, 2009, 4 (6), 9 p. (e6006). 10.1371/journal.pone.0006006 . hal-02665528

HAL Id: hal-02665528

<https://hal.inrae.fr/hal-02665528>

Submitted on 31 May 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Impaired Interleukin-1 β and c-Fos Expression in the Hippocampus Is Associated with a Spatial Memory Deficit in P2X₇ Receptor-Deficient Mice

Virginie F. Labrousse¹, Laurence Costes¹, Agnès Aubert¹, Muriel Darnaudéry^{1,2}, Guillaume Ferreira^{1,3}, Thierry Amédée¹, Sophie Layé^{1*}

1 Psychoneuroimmunologie, Nutrition et Génétique (PsyNuGen), INRA UMR 1286, CNRS UMR 5226, Université de Bordeaux, Bordeaux, France, **2** NEUROSTRESS EA 4347, "Université Lille Nord de France", Villeneuve d'Ascq, France, **3** Laboratoire Comportement, Neurobiologie et Adaptation, INRA UMR 85, CNRS UMR 6175, Université de Tours, Nouzilly, France

Abstract

Recent evidence suggests that interleukin-1 β (IL-1 β), which was originally identified as a proinflammatory cytokine, is also required in the brain for memory processes. We have previously shown that IL-1 β synthesis in the hippocampus is dependent on P2X₇ receptor (P2X₇R), which is an ionotropic receptor of ATP. To substantiate the role of P2X₇R in both brain IL-1 β expression and memory processes, we examined the induction of IL-1 β mRNA expression in the hippocampus of wild-type (WT) and homozygous P2X₇ receptor knockout mice (P2X₇R^{-/-}) following a spatial memory task. The spatial recognition task induced both IL-1 β mRNA expression and c-Fos protein activation in the hippocampus of WT but not of P2X₇R^{-/-} mice. Remarkably, P2X₇R^{-/-} mice displayed spatial memory impairment in a hippocampal-dependent task, while their performances in an object recognition task were unaltered. Taken together, our results show that P2X₇R plays a critical role in spatial memory processes and the associated hippocampal IL-1 β mRNA synthesis and c-Fos activation.

Citation: Labrousse VF, Costes L, Aubert A, Darnaudéry M, Ferreira G, et al. (2009) Impaired Interleukin-1 β and c-Fos Expression in the Hippocampus Is Associated with a Spatial Memory Deficit in P2X₇ Receptor-Deficient Mice. PLoS ONE 4(6): e6006. doi:10.1371/journal.pone.0006006

Editor: Bernhard T. Baune, James Cook University, Australia

Received: January 6, 2009; **Accepted:** May 27, 2009; **Published:** June 23, 2009

Copyright: © 2009 Labrousse et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Funding: This research was supported by Région Aquitaine grants n°20040305001A21 and n°20030305001 to S.L. and the INRA. V.F.L. is a PhD researcher working with the Région Aquitaine and the INRA. Quantitative PCR analyses were performed at the Bordeaux Genotyping and Sequencing facility (grants from the Conseil Régional d'Aquitaine n°20030304002FA and n°20040305003FA, and European Union FEDER grant n°2003227). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing Interests: The authors have declared that no competing interests exist.

* E-mail: sophie.laye@bordeaux.inra.fr

Introduction

The interleukin-1 (IL-1) family is composed of three closely-related proteins that are products of different genes: two agonists, IL-1 α and IL-1 β , and a naturally-occurring IL-1 receptor antagonist (IL-1ra) [1]. All the members of this family are synthesized and/or released in the brain by both glial cells and neurons [2]. Although most of the evidence gathered so far indicates that pathophysiological levels of IL-1 detrimentally affects neural plasticity and memory processes [3], recent evidence suggests that this cytokine is required for the physiological regulation of memory processes [4–7]. Blocking IL-1 receptors with IL-1ra impairs the maintenance of long-term potentiation (LTP) and hippocampal-dependent memory performance as measured in water maze and passive avoidance paradigms [7]. IL-1 receptor knockout mice exhibit a marked inability to express LTP in the hippocampus [4]. IL-1 β , rather than IL-1 α , could be involved in memory processes, since IL-1 β administration improves avoidance memory and contextual fear conditioning [7]. This is further reinforced by the observation that hippocampal IL-1 β mRNA expression is increased substantially after contextual fear conditioning and after LTP both *in vitro* and *in vivo* [5,6].

Extracellular ATP has been reported to trigger the release of IL-1 β by binding to type 2 purinergic receptors (P2 receptors) [8]. These receptors are subdivided into two subgroups: P2Y

metabotropic receptors (P2YR), which are G protein-coupled receptors, and P2X ionotropic receptors (P2XR), which are ligand-gated ion channels [9]. Interestingly, microglial cells, which are the main cell source of IL-1 β in the brain [2], express P2X₇R [10,11]. ATP binds to P2X₇R to induce the activation of caspase-1, the IL-1 β processing enzyme, also releasing mature IL-1 β in the process [8]. We and others have demonstrated *in vitro* that ATP triggers the maturation and release of IL-1 β in response to the bacterial endotoxin lipopolysaccharide (LPS) in microglia [12,13]. *In vivo*, LPS-induced IL-1 β mRNA expression is significantly lower in the brain of P2X₇R^{-/-} mice compared with WT mice [12].

Although the last decade has seen a wealth of knowledge on the effects of IL-1 β in the hippocampus, much less is known about the mechanism of IL-1 β release under physiological conditions, especially during hippocampal-dependent memory processes. We have recently identified P2X₇R as a key component in LPS-mediated mature IL-1 β release by microglia [12], and it has been shown that IL-1 β synthesis is induced in the hippocampus by memory processes [5]. Hence, the aim of the present study was to investigate the role of P2X₇R in memory processes and hippocampal IL-1 β production. For this purpose, we examined the memory performances of P2X₇R^{-/-} mice in two spontaneous recognition paradigms (Y maze and object recognition). Our results show that after 30 minutes of retention, spatial memory was impaired in P2X₇R^{-/-} mice, while their performances were

unaltered in an object recognition memory task. Moreover, the induction of IL-1 β synthesis and c-Fos expression in the hippocampus in response to the spatial task was abolished in P2X₇R^{-/-} mice, suggesting that hippocampal IL-1 β production may be involved in spatial memory performance.

Materials and Methods

Animals

Animal experiments were carried out according to the INRA Quality Reference System (1), and to relevant French (Directive 87–148, Ministère de l'Agriculture et de la Pêche) and international (Directive 86–609, November 24th 1986, European Community) legislation. Every effort was made to minimize suffering and the number of animals used. Experiments were performed on 3-month-old C57BL/6 (WT, n = 33) and homozygous P2X₇R^{-/-} (n = 33) male mice weighing 25–30 g. The P2X₇R^{-/-} mice were raised on a C57Bl/6 background and were kindly provided by Dr. Gabel (Pfizer, Groton, USA) [14]. Lack of the P2X₇ receptor was confirmed in the P2X₇R^{-/-} mice using PCR (not shown), as previously described [15]. P2X₇R^{-/-} and C57BL/6 mice were raised in the laboratory [12] and kept in groups of three in transparent polycarbonate cages (26.6×21.4×14.3 cm). They were maintained under standard colony conditions on corn cob litter in a temperature- (23±1°C) and humidity- (40%) controlled animal room under a 12 h light/dark cycle (7:00–19:00), with *ad libitum* access to food and water.

Behavioral measurements

Behavioral testing took place in the morning (between 8:00 AM and 11:00 AM). Mice were first handled for 5 min every day two weeks before launching the experiments. All the experiments were performed in a room adjacent to the *vivarium* under 78 lux of illumination.

Spatial recognition. The spontaneous spatial recognition in the Y-maze was used as a hippocampal-dependent test [16] as part of a two-trial procedure, as previously described [17–20]. The apparatus was a Y-shaped maze made of gray plastic. Each arm was 34 cm long, 8 cm wide and 14 cm high. Extramaze visual cues were placed in the testing room and kept constant during the test. Because the three arms of the maze were identical, discrimination of novelty versus familiarity was only based on the different aspects of the environment that the mouse can perceive from each of them. The floor of the maze was covered with dirty sawdust from the home cages of several animals, and was mixed between sessions in order to eliminate olfactory cues. In the first trial, one arm of the Y-maze was closed with a guillotine door and mice (n = 9/experimental group) were allowed to visit two arms of the Y-maze for 5 min. After a 30 min inter-trial interval (ITI), the mice were again placed in the start arm for the trial 2 and allowed free access to all three arms for 5 min. Start and closed arms were randomly assigned to each mouse. Arm entries were defined as all four paws entering the arm. Preference for novelty was also measured using a 2-min ITI between acquisition and retrieval. This 2 min ITI was employed to control for potential motivational disturbances and verify that all groups made more visits to the novel arm when mnemonic demand was minimal [17]. The different ITI tests were separated by a 7-day interval. Analyses were based on the time spent exploring the novel and the familiar arms during the first 3 minutes of the trial 2 [21].

Object recognition. The visual object recognition test was used as a hippocampal-independent task [22–25]. Mice (n = 9/experimental group) were acclimatized in a 40×40 cm square

cage made of white coated plywood with 16 cm-high walls for 15 min per day during the week before training. The floor of the maze was covered with corn cob litter which was mixed between each trial in order to remove olfactory cues. During a 10-min training period, two identical plastic blocks with particular shapes and colors were presented. After a 30 min ITI, one of the familiar blocks was replaced by a novel object, with a different shape and color, to test for memory retention. During the 5 min of testing, exploration of an object was defined as pointing the nose to the object and/or touching the object with the nose. Longer ITI (1-hour and 24-hour) were also employed to determine whether all groups recognized the novel object when retention was longer (data not shown) [22]. Analyses were based on the time spent exploring the novel and the familiar objects and the object recognition score was calculated as the difference in time spent exploring the novel and the familiar objects.

IL-1 β real-time PCR

WT and P2X₇R^{-/-} mice were sacrificed with a lethal dose of pentobarbital immediately after completion of behavioral testing [2-min ITI or 30-min ITI Y-maze (Y-maze group) or free exploratory behavior (control group)]. Control-group mice were allowed to freely explore a new cage for the same amount of time as their spatially-trained counterparts (i.e. twice during 5 min each with either a 2-min ITI or a 30-min ITI) in the experimental Y-maze room. This control condition allowed the assessment of activation that results from exploratory behavior, sensorimotor stimulation, stress, and other factors not related to the spatial task. Hence, difference between the two profiles, *i.e.* the activation induced in the Y-maze group and in the control group, may reflect specific learning-induced changes.

Mice (n = 6/experimental group) were transcardially perfused with Phosphate Buffer Saline (PBS) and the hippocampi and hypothalamus were quickly removed and frozen in dry ice. Two μ g of total RNA obtained from each brain area were reverse-transcribed with Moloney Murine Leukemia Virus reverse transcriptase (Invitrogen, Cergy-Pontoise, France). Quantitative PCR was then performed using the Applied Biosystems Assay-on-Demand Gene Expression Products protocol, as previously described [12]. Briefly, cDNAs for IL-1 β and housekeeping gene (β 2-microglobulin) were amplified by PCR by using an oligonucleotide probe with a 5' fluorescent reporter dye (6-FAM) and a 3' quencher dye (NFQ). Fluorescence was determined using an AB 7500 Real-Time PCR system (Applied Biosystems, Foster city, CA) and final quantification was performed using the comparative threshold (Ct) method as previously described [12]. For each experimental sample, the difference between target and housekeeping gene Ct values (Δ Ct) was used to normalize for differences in the amount of total nucleic acid added to each reaction and in the efficiency of the RT step. Values were then expressed as relative fold change to the mean Δ Ct value obtained for the group of control mice (free exploratory behavior, see below) (calibrator Δ Ct) by subtracting Δ Ct for each experimental sample from the calibrator Δ Ct (= $\Delta\Delta$ Ct). The amount of target gene (linear value) normalized to the housekeeping gene and relative to the calibrator was determined by $2^{-\Delta\Delta$ Ct} (relative fold change: RFC).

In order to assess regional specificity of IL-1 β induction, we analyzed IL-1 β mRNA in the hypothalamus immediately after the completion of the 30 min ITI spatial recognition task using the same protocol as described above.

c-Fos immunohistochemistry and immunocounting

WT and P2X₇R^{-/-} mice (n = 6/experimental group) were sacrificed with a lethal dose of pentobarbital 90 min after the

acquisition session of the spatial recognition task (Y-maze group) or free exploratory behavior (control group). This delay was chosen based on previous data showing that 90 min corresponds to the peak of the expression of c-Fos protein in the hippocampus in mice submitted to spatial memory tests [26]. WT and P2X₇R^{-/-} mice of the Y-maze groups were subjected to a 5-min test in the Y-maze with 30-min ITI. Control-group mice were allowed to freely explore a new cage for the same amount of time as their spatially trained counterparts were given in the experimental Y-maze room (i.e. two 5-min periods with a 30-min ITI).

c-Fos expression was measured by immunohistochemistry as previously described [27]. After intracardiac perfusion with PBS (pH 7.4) followed by 4% paraformaldehyde, the brains were removed, post-fixed in the same fixative for 4 h at 4°C, cryoprotected in 30% sucrose for 24 h, and then quickly frozen in liquid nitrogen and stored at -80°C before sectioning. Coronal cryostat sections of 30 μm width through the hippocampus (bregma: from -0.9 mm to -3.1 mm) were collected free-floating. Briefly, rabbit polyclonal antiserum raised against c-Fos (Santa Cruz Biotechnology, Santa Cruz, CA) was diluted 1:1000 in Tris Buffer Saline (TBS) containing 0.3% Triton X-100, 2% donkey serum and 1% bovine serum albumin (BSA), and sections were incubated overnight at room temperature (RT) before being incubated for 2 h with biotinylated donkey anti-rabbit antibody (1:1000; Amersham Pharmacia Biotech Europe, Freiburg, Germany), for 2 h with avidin-biotin peroxidase complex (1:1000; Vectastain ABC kit, Vector laboratories, Burlingame, CA), and finally revealed with diaminobenzidine via the nickel-enhanced glucose oxidase method [28]. Sections were then slide-mounted onto gelatin-coated slides. The procedure also included negative controls with omission of the primary antibody, which did not show any immunoreaction.

Brain sections were examined under a light microscope with a 10× lens (Nikon Eclipse E 400) and the images were captured using a high-resolution digital Nikon DXM 1200 camera (Nikon Corporation, Champigny-sur-Marne, France). Camera aperture, magnification, light power and exposure time were fixed for all images. ACT-1-generated images (Nikon Corporation, Champigny-sur-Marne, France) were stored. Image-editing software (Adobe Photoshop, Adobe Systems, San Jose, CA) was used to adjust image size, brightness, and contrast. Quantification of c-Fos immunoreactive cells was performed with the aid of Scion Image public domain NIH-imaging software (Frederick, MD). The number of c-Fos-positive cells was quantified in the dentate gyrus (DG) and CA1 of the hippocampus from consecutive sections (n = 5–6 sections per mice; n = 6 mice per group) across both hemispheres, and these counts were averaged to produce a mean. Dorsal and ventral regions of the hippocampus were analyzed, but only the data of the dorsal region were presented as there were no significant differences in the ventral part. The number of c-Fos-positive cells was also quantified in the arcuate nucleus of the hypothalamus to control for specific hippocampal activation.

Statistical analysis

All data are expressed as means ± SEM. A two-way analysis of variance (ANOVA) with genotype (WT vs. P2X₇R^{-/-}) as between factor and arm or object (novel vs. familiar) as within factor were performed for spatial and object recognition analysis. Specific comparisons between novel and familiar (arms or objects) or between WT and P2X₇R^{-/-} were assessed by paired and unpaired Student t-tests respectively. IL-1β mRNA and c-Fos quantification were analyzed by two-way ANOVA with genotype (WT vs. P2X₇R^{-/-}) and testing procedure (Control vs. Y-maze training) as between factors. When a significant interaction was reported, ANOVAs were

followed by Fisher LSD post-hoc comparisons. Relations between recognition score (calculated as the difference between the investigation time for the novel and the familiar stimulus) and c-Fos levels were evaluated using Bravais-Pearson's correlation tests. The level of statistical significance was set at p < 0.05.

Results

1- Spatial memory performance is impaired in P2X₇R^{-/-} mice

We first measured the spatial memory performances of WT and P2X₇R^{-/-} mice using a spatial recognition test designed as a hippocampus-dependent task [16]. After 30 minutes of retention, the spatial memory performance was different across the genotype (Fig. 1A). A two-way ANOVA revealed no effect of genotype ($F_{(1,16)} = 0.46$, n.s.) or arm ($F_{(1,16)} = 0.21$, n.s.), but there was a significant interaction between genotype and arm ($F_{(1,16)} = 4.5$, $p < 0.05$). Post-hoc analysis revealed that WT group spent less time in the familiar arm than P2X₇R^{-/-} group (Fisher LSD, $p < 0.05$). Further analysis indicated that WT mice significantly distinguished between the novel and the familiar arm (paired-t test, $t(8) = 5.7$, $p < 0.001$), unlike the P2X₇R^{-/-} group which exhibit a random exploration of the two arms ($p = 0.42$). No differences for the time spent in the starting arm between WT and P2X₇R^{-/-} mice were measured ($p = 0.176$, not shown). In order to evaluate whether the impairment revealed in P2X₇R^{-/-} mice was a memory deficit or a performance deficit related to motivational alteration, animals were tested in the spatial recognition test with a short ITI (2-min). When the mnemonic demand was minimal, all mice exhibited a clear preferential exploration of the novelty (arm effect, $F_{(1,16)} = 17.2$, $p < 0.001$, no effect of genotype, $F_{(1,16)} = 0.15$, n.s., or arm x genotype, $F_{(1,16)} = 1.64$, n.s.; Table 1).

We then measured performances of WT and P2X₇R^{-/-} mice in an object recognition task designed as a hippocampus-independent task [22]. As shown on Fig. 1B, after 30 minutes of retention, all animals showed greater exploration of the new object (object effect, $F_{(1,16)} = 41.6$, $p < 0.0001$, no effect of genotype, $F_{(1,16)} = 0.01$, n.s., or object x genotype, $F_{(1,16)} = 0.09$, n.s.), indicating that they readily recognized the familiar object (paired-t test, WT, $t(8) = 4.52$, $p < 0.01$; P2X₇R^{-/-}, $t(8) = 4.52$, $p < 0.01$). To evaluate whether longer ITI could reveal an impairment of object recognition in P2X₇R^{-/-} mice, animals were tested with a 1-h or a 24-h ITI. As was the case with the 30-min ITI, all animals preferred the novel object over the familiar object with a 1-h ITI (object effect, $F_{(1,16)} = 8.2$, $p = 0.01$), whereas there was no discrimination of the novel object with a 24-h ITI ($F_{(1,16)} < 1$ for each effect; data not shown).

2- IL-1β mRNA expression after the spatial memory task is blocked in the hippocampus of P2X₇R^{-/-} mice

The impaired behavioral performances of P2X₇R^{-/-} mice in the spatial recognition task prompted us to investigate whether hippocampal IL-1β synthesis was altered in these mice. For this purpose, we measured IL-1β mRNA expression by real-time PCR in the hippocampus of WT and P2X₇R^{-/-} mice immediately after the completion of two testing procedures: free exploratory behavior (control) or spatial recognition task (Y-maze with 2-min ITI or 30-min ITI).

Whatever the ITI, the Y maze experience increased IL-1β expression in the hippocampus (testing procedure effect, ITI 2-min, $F_{(1,20)} = 17.7$; $p < 0.001$; ITI 30-min, $F_{(1,20)} = 5.7$; $p < 0.05$). However, as shown on figure 2A, the IL-1β mRNA expression in the hippocampus was differentially affected by the Y maze exposure, depending of the group (genotype x testing procedure,




Figure 1. Recognition memory performance after 30 minutes of retention in P2X₇R^{-/-} mice. (A) Time spent (sec) in the novel or the familiar arm after a 30-min ITI. (B) Time spent (sec) exploring the novel or the familiar object after a 30-min ITI. Spatial recognition memory, but not object recognition memory, was impaired in P2X₇R^{-/-} mice after a 30-min ITI. ****p*<0.001, ***p*<0.01, **p*<0.05. doi:10.1371/journal.pone.0006006.g001

Table 1. Preference for novelty was measured using a 2-min ITI in WT and P2X₇R^{-/-} mice.

	Preference for Novelty (ITI = 2 min)	
	Novel arm (sec.)	Familiar arm (sec.)***
WT	84.28 ± 8.84	36.82 ± 6.60
P2X ₇ R ^{-/-}	74.81 ± 5.34	49.64 ± 7.90

Both strain exhibited a clear preferential exploration of the novelty (Arm effect, ****p*<0.001).

doi:10.1371/journal.pone.0006006.t001

2-min ITI, $F_{(1,20)} = 8.6$; $p < 0.01$; 30-min ITI, $F_{(1,20)} = 4.4$; $p < 0.05$). *Post-hoc* analysis revealed that the effect of the exposure to the Y maze was exclusively observed in WT mice (control vs. Y maze, 2-min ITI, $p < 0.0001$; 30-min ITI, $p < 0.01$), but not in P2X₇R^{-/-} mice (control vs. Y maze, 2-min ITI, $p = 0.37$; 30-min ITI, $p = 0.84$). Moreover, although there was no difference between WT and P2X₇R^{-/-} in the control condition (2-min ITI, $p = 0.33$; 30-min ITI, $p = 0.90$), IL-1 β mRNA expression was higher in WT mice than in P2X₇R^{-/-} mice in the Y-maze condition (2-min ITI, $p < 0.01$; 30-min ITI, $p < 0.01$).

Next, we analyzed the impact of Y maze exposure on IL-1B mRNA expression in a brain area not involved in spatial memory


IL-1 β mRNA expression

Figure 2. IL-1 β mRNA expression after the exposure to the Y maze in P2X₇R^{-/-} mice. IL-1 β mRNA was measured by real-time PCR on WT and P2X₇R^{-/-} mice sacrificed either after free exploration (control group) or completion of behavioral testing (Y-maze group). All data are expressed as mean relative fold change \pm SEM (see Materials and Methods for explanation). (A) IL-1 β mRNA on hippocampus extracts from the 2-min ITI Y-maze group and their controls (left-hand panel) and the 30-min ITI Y-maze group and their controls (right-hand panel). (B) IL-1 β mRNA on hypothalamus extracts from the 30-min ITI Y-maze group and their controls. Y maze exposure significantly increased the IL-1 β mRNA expression in the hippocampus of WT mice but not of P2X₇R^{-/-} mice. Whatever the group considered (WT or P2X₇R^{-/-}), no induction of IL-1 β mRNA expression was detected in hypothalamus after the Y maze experience. *** p <0.001, ** p <0.01. doi:10.1371/journal.pone.0006006.g002

(Fig. 2B). In order to assess the regional specificity of IL-1 β induction, we analyzed IL-1 β mRNA in the hypothalamus of the mice immediately after the completion of spatial recognition (Y-maze 30 min ITI) or free exploratory behavior (control). As expected, the Y maze exposure did not induce IL-1 β mRNA expression in the hypothalamus either in WT or P2X₇R^{-/-} mice (testing procedure effect, $F_{(1,20)} = 0.98$; n.s.; genotype, $F_{(1,20)} = 0.83$; p <0.01; genotype \times testing procedure, $F_{(1,20)} = 2.18$; n.s.).

3- Spatial memory deficits in P2X₇R^{-/-} mice are associated with blocked c-Fos activation in the hippocampus

It has previously been shown that c-Fos immunoreactivity is a cellular activity marker that provides information on brain regions involved in learning and memory processes [29]. Because spatial memory performance and task-induced IL-1 β mRNA expression in the hippocampus were impaired in our P2X₇R^{-/-} mice, we sought to investigate whether cellular activity in the hippocampus had been affected. We therefore measured by immunohistochemistry, c-Fos expression in the hippocampus (DG, CA1) of WT and P2X₇R^{-/-} submitted (Y-maze 30 min ITI) or not (control, i.e. free exploratory behavior) to a spatial recognition task (Figure 3A,

B, left-hand panel). Two-way ANOVA revealed a significant effect of genotype (DG, $F_{(1,20)} = 5.4$; p <0.05; CA1, $F_{(1,20)} = 4.5$; p <0.05) and testing procedure (DG, $F_{(1,20)} = 8.5$; p <0.01; CA1, $F_{(1,20)} = 14.9$; p <0.001). However, the impact of Y maze exposure differed across the two groups of animals, as suggested by the trend towards genotype \times testing procedure interaction in DG ($F_{(1,20)} = 3.6$; $p = 0.07$) and the significant interaction in CA1 ($F_{(1,20)} = 13.4$; p <0.001). Subsequent post-hoc analysis revealed that the spatial recognition task significantly up-regulated c-Fos expression in the DG and CA1 of WT (DG, p <0.01; CA1, p <0.001) mice but not P2X₇R^{-/-} mice (DG, $p = 0.47$; CA1, p <0.89). Moreover, although there was no difference between WT and P2X₇R^{-/-} in the control condition, c-Fos expression was higher in WT mice than in P2X₇R^{-/-} mice in the Y-maze condition (DG, p <0.01; CA1, p <0.001). Assessment of c-Fos expression in the hypothalamus (Fig. 3B, right-hand panel) after the completion of spatial recognition did not show any effect of testing procedure either in WT or P2X₇R^{-/-} mice (testing procedure effect, $F_{(1,20)} = 0.96$, n.s.; genotype \times testing procedure effect $F_{(1,20)} = 0.008$, n.s.).

We then examined the relationship between the brain c-Fos expression after the Y maze and the spatial memory performance in this task. As demonstrated in the Figure 3C, c-Fos expression in the hippocampus was positively correlated to the spatial




Figure 3. c-Fos expression after the exposure to the Y maze in P2X₇R^{-/-} mice. c-Fos immunohistochemistry analysis was performed on the hippocampus and the hypothalamus of WT and P2X₇R^{-/-} mice sacrificed 90 min after either the spatial recognition acquisition session (Y-maze group) or free exploration (control group). (A) Representative images of c-Fos immunohistochemistry in the DG (left-hand panel) and CA1 (middle panel) regions of the hippocampus and in the hypothalamus (right-hand panel). Scale bar = 100 μ m. (B) Quantification of c-Fos-positive cells was performed in the DG (left-hand panel) and CA1 (middle panel) regions of the hippocampus and in the hypothalamus (right-hand panel). The spatial recognition in the Y maze task significantly up-regulated c-Fos expression in the DG and CA1 of WT mice but not of P2X₇R^{-/-} mice. No increase of c-Fos expression was detected in hypothalamus after the Y maze experience. *** $p < 0.001$; ** $p < 0.01$. (C) Correlation between the spatial recognition score (sec) and the number of c-Fos-positive cells induced by the Y maze task in the DG (left-hand), in the CA1 (middle panel) and in the hypothalamus (right-hand panel). Pearson's correlation coefficients (r) and corresponding significance (p) are displayed within each correlation windows. The spatial recognition performance in the Y maze after a 30-min ITI was positively correlated to the number of c-Fos positive cells in the DG and the CA1 of the hippocampus, but not with the number of c-Fos positive cells in the hypothalamus. doi:10.1371/journal.pone.0006006.g003

recognition score (DG, $r = 0.64$, $p < 0.05$; CA1, $r = 0.81$, $p < 0.01$). In contrast, no significant correlation was reported between the c-Fos expression in the hypothalamus and the memory performance (Figure 3C, right-hand panel, $r = 0.02$, $p = 0.93$).

Discussion

We show here that compared to WT mice, P2X₇R^{-/-} mice exhibit spatial memory deficits in the Y maze, a hippocampal dependent task, whereas their memory performance was intact in an object recognition paradigm. We further report that, contrary to the WT animals, P2X₇R^{-/-} mice had no increase of hippocampal IL-1 β mRNA and c-Fos expression after the Y maze exposure. Taken together, our results clearly support a crucial role for P2X₇R-mediated IL-1 β expression in memory processes involving the hippocampus.

Our study employed spontaneous recognition tasks (spatial and object), which exploit the natural tendency of rodents to explore novel place or stimuli more than familiar ones and have been widely validated as measure of recognition memory (reviewed in [16,25]). These paradigms do not require the learning of a rule, and factors that may particularly influence performance in mice such as emotional states generated by commonly used procedures (food deprivation, water avoidance or electric foot-shock) are minimized, making it useful for studying memory in rodents [30]. Spontaneous recognition is based on novelty preference, however, when a delay of several minutes (e.g. 30 min), the decrease of the exploration of the familiar arm (or object) during the trial 2 reflects the memory of the previous exploration during the first trial [17,19,20,30–33]. Following a 30-min ITI, WT mice showed significant spatial recognition, while P2X₇R^{-/-} mice did not. Furthermore, P2X₇R^{-/-} mice display intact performance in an object recognition task with the same 30-min ITI, suggesting that memory is not affected in a non-spatial task. Even if we cannot rule out that with longer ITI (such as 1h30 or 2h), the ability of mice to recognize object would decrease faster in P2X₇R^{-/-} mice than in WT mice, the differential impairment between spatial and object recognition tasks for a similar delay is particularly interesting regarding the considerable literature showing that spatial memory and object memory do not involve similar brain areas [22–25,34]. Hippocampus is clearly involved for spatial memory; in contrast, visual/object recognition requires intact perirhinal area, but hippocampus seems less crucial [35]. Taken together, our results suggest that the lack of P2X₇R specifically affects hippocampal functioning as well as the downstream events stemming from it that are specifically involved in spatial recognition. This possibility is well supported by the high expression of P2X₇R both by presynaptic terminals of mossy fibers targeting CA3 pyramidal neurons [36] and by excitatory synaptic terminals in the CA1 and dentate gyrus regions of the hippocampus [37].

An important new finding of our study is the demonstration of the reduced expression of learning-induced IL-1 β mRNA in the hippocampus of P2X₇R^{-/-} mice. Interestingly, IL-1 β was induced in the hippocampus but not in the hypothalamus, suggesting that IL-1 β induction is specifically linked to spatial memory performance. This observation is consistent with previous reports showing that IL-1 β gene expression is induced in the hippocampus by LTP in freely-moving rats [6,38] and by two other hippocampal-dependent memory tasks, i.e. passive avoidance and contextual fear conditioning [5,39]. Since, spontaneous recognition memory tasks do not allow a subtle assessment of memory processes (learning, flexibility, reference memory etc.), it would be interesting to evaluate whether the impaired spatial

recognition memory in P2X₇R^{-/-} mice can be generalized to other hippocampal-dependent tasks. In this context, it has previously been shown that IL-1 receptor activation is necessary for memory tasks such as passive avoidance, contextual fear conditioning or the Morris water maze [4–7].

IL-1 β mRNA expression was induced in the hippocampus of WT but not P2X₇R^{-/-} mice submitted to the spatial recognition task, independently of the ITI. These results suggest that P2X₇R^{-/-} is a prerequisite for IL-1 β mRNA expression, as previously demonstrated after LPS treatment [12]. In the hippocampus, P2X₇R stimulation may either directly or indirectly activate neurons and elicit changes in their excitability [36,37,40]. Since an intact performance was reported in P2X₇R^{-/-} mice after the 2-min ITI, despite a lack of IL-1 β after the Y maze, we hypothesized that the induction of IL-1 β in the hippocampus was an important factor underlying the spatial memory processes (e.g. retention of the spatial information), but not the spatial novelty preference. We did not identify the cell type that synthesizes and releases IL-1 β in the hippocampus after the spatial task. Both neurons and microglia have been reported to express IL-1 β in the non-pathological brain. There is a large body of evidence demonstrating the presence of P2X₇R on microglia [8,13,41,42]. The activation of these receptors, via downstream inflammatory stimuli such as LPS or direct activation by ATP, triggers microglial release of IL-1 β , as we and others have previously demonstrated *in vitro* [8,12,13,15,41].

Similarly to IL-1 β mRNA expression, c-Fos expression was induced in the hippocampus of WT but not P2X₇R^{-/-} mice submitted to the spatial recognition task. c-Fos is an immediate early gene rapidly induced by a variety of stimuli, including stimuli involved in regulating synaptic plasticity [43]. This suggests that the increased neuronal activity in hippocampus observed after the exposure to the spatial task is linked to the hippocampal activation of P2X₇R. Moreover, in our study, we found that c-Fos activation in the DG and CA1 of the hippocampus is positively correlated to spatial recognition performance. Indeed, mice exhibiting the best recognition scores also show the highest hippocampal c-Fos expression after the spatial task. In contrast, c-Fos levels in the hypothalamus were not linked to the spatial recognition score. The hypothalamus showed neither c-Fos expression nor IL-1 β mRNA induction after the spatial task. This last observation further reinforces the link between P2X₇R, IL-1 β and c-Fos activation in the hippocampus in response to spatial memory performance.

IL-1 β is synthesized in the cell cytoplasm as a leaderless procytokine (pro-IL-1 β) which tends to accumulate in the absence of a maturation-promoting second stimulus. IL-1 β -converting enzyme (ICE) is the key protease responsible for the intracellular processing of pro-IL-1 β into mature IL-1 β . Numerous *in vitro* and *in vivo* experiments using P2X₇R^{-/-} mice have conclusively linked the stimulation of P2X₇R by extracellular ATP to the activation of ICE and the downstream production of mature IL-1 β [10,14,44–46]. Therefore, there is little doubt that a lack of P2X₇R necessarily impacts IL-1 β production, resulting in severe reduction if not a complete loss. Very importantly, IL-1 β induced its own synthesis in the brain, and this process is dependent on P2X₇R [12]. It is therefore possible that an initial small increase in IL-1 β secretion through the recruitment of P2X₇R could up-regulate the IL-1 β mRNA induction in the hippocampus linked to a spatial recognition memory task. This idea is supported by the fact that contextual fear conditioning induces IL-1 β mRNA expression in the hippocampus of WT mice but not of IL-1 receptor knockout mice [5]. The impaired memory processes in P2X₇R^{-/-} mice could thus be linked to hippocampus-deficient IL-1 β synthesis. Therefore, our results are in accordance with recent studies

demonstrating that physiological levels of hippocampal IL-1 β are required for short-term spatial memory performance, whereas deficient hippocampal IL-1 β expression (as here) results in impaired memory.

The mechanisms underlying the role of P2X₇R in the synthesis of IL-1 β mRNA in relation to memory processes remain an enigma. P2X₇R has been associated to IL-1 β precursor processing, through the rapid activation of a specific caspase-1-dependent multiprotein complex [47]. The ATP-induced activation of P2X₇R decreases cytoplasmic potassium which in turn induces the maturation and the release of mature IL-1 β by monocytes [48,49], microglia [12] and Schwann cells [48,49]. In addition ATP by acting on presynaptic P2X₇R, modulates hippocampal synaptic transmission [36,37]. In the present work, it could be hypothesized that the Y maze exposure stimulates the ATP release by neighboring glial cells *via* an increase of the glutamate release [50]. This could in turn affect presynaptic P2X₇R. Alternatively, or additionally, the memory task may modulate the neuronal release of ATP, which could then act on nearby P2X₇R since ATP is co-stored and co-released with classical neurotransmitters from brain neurons [37,51] in an exocytotic manner [52] and acts as a fast neurotransmitter within the brain [53–55]. However, up to

now, the sole direct demonstration of P2X₇R modulation of hippocampal synaptic transmission reported a depression [36].

In conclusion, our study demonstrated that P2X₇R is involved in spatial memory processes and in the associated neuronal activity and IL-1 β mRNA synthesis in the hippocampus. These results therefore corroborate clinical observations showing that the impairment of P2X₇R function, due to mutation or polymorphism, leads to both immunological [56,57] and brain dysfunction [58–60], putatively due to the lack of IL-1 β synthesis and release.

Acknowledgments

Quantitative PCR analyses were performed at the Bordeaux Genotyping and Sequencing facility. We thank Françoise Moos for proofreading the Manuscript.

Author Contributions

Conceived and designed the experiments: VFL SL. Performed the experiments: VFL LC AA. Analyzed the data: VFL MD GF SL. Contributed reagents/materials/analysis tools: TA. Wrote the paper: VFL MD GF TA SL.

References

- Dinarello CA (2002) The IL-1 family and inflammatory diseases. *Clin Exp Rheumatol* 20: S1–13.
- Rothwell NJ (1991) Functions and mechanisms of interleukin 1 in the brain. *Trends Pharmacol Sci* 12: 430–436.
- Coogan AN, O'Connor JJ (1999) Interleukin-1beta inhibits a tetraethylammonium-induced synaptic potentiation in the rat dentate gyrus *in vitro*. *Eur J Pharmacol* 374: 197–206.
- Avital A, Goshen I, Kamsler A, Segal M, Iverfeldt K, et al. (2003) Impaired interleukin-1 signaling is associated with deficits in hippocampal memory processes and neural plasticity. *Hippocampus* 13: 826–834.
- Goshen I, Kreisel T, Ounallah-Saad H, Renbaum P, Zalsstein Y, et al. (2007) A dual role for interleukin-1 in hippocampal-dependent memory processes. *Psychoneuroendocrinology* 32: 1106–1115.
- Schneider H, Pitossi F, Balschun D, Wagner A, del Rey A, et al. (1998) A neuromodulatory role of interleukin-1beta in the hippocampus. *Proc Natl Acad Sci U S A* 95: 7778–7783.
- Yirmiya R, Winocur G, Goshen I (2002) Brain interleukin-1 is involved in spatial memory and passive avoidance conditioning. *Neurobiol Learn Mem* 78: 379–389.
- Ferrari D, Pizzirani C, Adinolfi E, Lemoli RM, Curti A, et al. (2006) The P2X₇ receptor: a key player in IL-1 processing and release. *J Immunol* 176: 3877–3883.
- Di Virgilio F, Borea PA, Illes P (2001) P2 receptors meet the immune system. *Trends Pharmacol Sci* 22: 5–7.
- Ferrari D, Villalba M, Chiozzi P, Falzoni S, Ricciardi-Castagnoli P, et al. (1996) Mouse microglial cells express a plasma membrane pore gated by extracellular ATP. *J Immunol* 156: 1531–1539.
- Yu Y, Ugawa S, Ueda T, Ishida Y, Inoue K, et al. (2008) Cellular localization of P2X₇ receptor mRNA in the rat brain. *Brain Res* 1194: 45–55.
- Mingam R, De Smedt V, Amedee T, Bluthé RM, Kelley KW, et al. (2008) *In vitro* and *in vivo* evidence for a role of the P2X₇ receptor in the release of IL-1 beta in the murine brain. *Brain Behav Immun* 22: 234–244.
- Sanz JM, Di Virgilio F (2000) Kinetics and mechanism of ATP-dependent IL-1 beta release from microglial cells. *J Immunol* 164: 4893–4898.
- Solle M, Labasi J, Perregaux DG, Stam E, Petrushova N, et al. (2001) Altered cytokine production in mice lacking P2X₇ receptors. *J Biol Chem* 276: 125–132.
- Le Feuvre RA, Brough D, Iwakura Y, Takeda K, Rothwell NJ (2002) Priming of macrophages with lipopolysaccharide potentiates P2X₇-mediated cell death via a caspase-1-dependent mechanism, independently of cytokine production. *J Biol Chem* 277: 3210–3218.
- Conrad CD, Galea LA, Kuroda Y, McEwen BS (1996) Chronic stress impairs rat spatial memory on the Y maze, and this effect is blocked by tianeptine pretreatment. *Behav Neurosci* 110: 1321–1334.
- Dellu F, Contarino A, Simon H, Koob GF, Gold LH (2000) Genetic differences in response to novelty and spatial memory using a two-trial recognition task in mice. *Neurobiol Learn Mem* 73: 31–48.
- Vallee M, MacCari S, Dellu F, Simon H, Le Moal M, et al. (1999) Long-term effects of prenatal stress and postnatal handling on age-related glucocorticoid secretion and cognitive performance: a longitudinal study in the rat. *Eur J Neurosci* 11: 2906–2916.
- Contarino A, Dellu F, Koob GF, Smith GW, Lee KF, et al. (1999) Reduced anxiety-like and cognitive performance in mice lacking the corticotropin-releasing factor receptor 1. *Brain Res* 835: 1–9.
- Palin K, Bluthé RM, Verrier D, Tridon V, Dantzer R, et al. (2004) Interleukin-1beta mediates the memory impairment associated with a delayed type hypersensitivity response to bacillus Calmette-Guérin in the rat hippocampus. *Brain Behav Immun* 18: 223–230.
- Mayo W, Dellu F, Robel P, Cherkaoui J, Le Moal M, et al. (1993) Infusion of neurosteroids into the nucleus basalis magnocellularis affects cognitive processes in the rat. *Brain Res* 607: 324–328.
- Dere E, Huston JP, De Souza Silva MA (2007) The pharmacology, neuroanatomy and neurogenetics of one-trial object recognition in rodents. *Neurosci Biobehav Rev* 31: 673–704.
- Broadbent NJ, Squire LR, Clark RE (2004) Spatial memory, recognition memory, and the hippocampus. *Proc Natl Acad Sci U S A* 101: 14515–14520.
- Bevins RA, Besheer J (2006) Object recognition in rats and mice: a one-trial non-matching-to-sample learning task to study 'recognition memory'. *Nat Protoc* 1: 1306–1311.
- Winters BD, Saksida LM, Bussey TJ (2008) Object recognition memory: neurobiological mechanisms of encoding, consolidation and retrieval. *Neurosci Biobehav Rev* 32: 1055–1070.
- Farivar R, Zangenehpour S, Chaudhuri A (2004) Cellular-resolution activity mapping of the brain using immediate-early gene expression. *Front Biosci* 9: 104–109.
- Nadjar A, Bluthé RM, May MJ, Dantzer R, Parnet P (2005) Inactivation of the cerebral NFkappaB pathway inhibits interleukin-1beta-induced sickness behavior and c-Fos expression in various brain nuclei. *Neuropsychopharmacology* 30: 1492–1499.
- Shu SY, Ju G, Fan LZ (1988) The glucose oxidase-DAB-nickel method in peroxidase histochemistry of the nervous system. *Neurosci Lett* 85: 169–171.
- Herdegen T, Leah JD (1998) Inducible and constitutive transcription factors in the mammalian nervous system: control of gene expression by Jun, Fos and Krox, and CREB/ATF proteins. *Brain Res Brain Res Rev* 28: 370–490.
- Dellu F, Mayo W, Cherkaoui J, Le Moal M, Simon H (1992) A two-trial memory task with automated recording: study in young and aged rats. *Brain Res* 588: 132–139.
- Dillon GM, Qu X, Marcus JN, Dodart JC (2008) Excitotoxic lesions restricted to the dorsal CA1 field of the hippocampus impair spatial memory and extinction learning in C57BL/6 mice. *Neurobiol Learn Mem* 90: 426–433.
- Niu L, Cao B, Zhu H, Mei B, Wang M, et al. (2008) Impaired *in vivo* synaptic plasticity in dentate gyrus and spatial memory in juvenile rats induced by prenatal morphine exposure. *Hippocampus*.
- Sanderson DJ, Gray A, Simon A, Taylor AM, Deacon RM, et al. (2007) Deletion of glutamate receptor-A (GluR-A) AMPA receptor subunits impairs one-trial spatial memory. *Behav Neurosci* 121: 559–569.
- Brown MW, Aggleton JP (2001) Recognition memory: what are the roles of the perirhinal cortex and hippocampus? *Nat Rev Neurosci* 2: 51–61.
- Eichenbaum H, Yonelinas AP, Ranganath C (2007) The medial temporal lobe and recognition memory. *Annu Rev Neurosci* 30: 123–152.
- Armstrong JN, Brust TB, Lewis RG, MacVicar BA (2002) Activation of presynaptic P2X₇-like receptors depresses mossy fiber-CA3 synaptic transmission through p38 mitogen-activated protein kinase. *J Neurosci* 22: 5938–5945.

37. Sperlagh B, Kofalvi A, Deuchars J, Atkinson L, Milligan CJ, et al. (2002) Involvement of P2X₇ receptors in the regulation of neurotransmitter release in the rat hippocampus. *J Neurochem* 81: 1196–1211.
38. Balschun D, Randolph A, Pitossi F, Schneider H, Del Rey A, et al. (2003) Hippocampal interleukin-1 beta gene expression during long-term potentiation decays with age. *Ann N Y Acad Sci* 992: 1–8.
39. Depino AM, Alonso M, Ferrari C, del Rey A, Anthony D, et al. (2004) Learning modulation by endogenous hippocampal IL-1: blockade of endogenous IL-1 facilitates memory formation. *Hippocampus* 14: 526–535.
40. Ireland MF, Noakes PG, Bellingham MC (2004) P2X₇-like receptor subunits enhance excitatory synaptic transmission at central synapses by presynaptic mechanisms. *Neuroscience* 128: 269–280.
41. Brough D, Le Feuvre RA, Iwakura Y, Rothwell NJ (2002) Purinergic (P2X₇) receptor activation of microglia induces cell death via an interleukin-1-independent mechanism. *Mol Cell Neurosci* 19: 272–280.
42. Boucsein C, Zacharias R, Farber K, Pavlovic S, Hanisch UK, et al. (2003) Purinergic receptors on microglial cells: functional expression in acute brain slices and modulation of microglial activation in vitro. *Eur J Neurosci* 17: 2267–2276.
43. Guzowski JF (2002) Insights into immediate-early gene function in hippocampal memory consolidation using antisense oligonucleotide and fluorescent imaging approaches. *Hippocampus* 12: 86–104.
44. Ferrari D, Chiozzi P, Falzoni S, Dal Susino M, Melchiorri L, et al. (1997) Extracellular ATP triggers IL-1 beta release by activating the purinergic P2Z receptor of human macrophages. *J Immunol* 159: 1451–1458.
45. Labasi JM, Petrushova N, Donovan C, McCurdy S, Lira P, et al. (2002) Absence of the P2X₇ receptor alters leukocyte function and attenuates an inflammatory response. *J Immunol* 168: 6436–6445.
46. Chessell IP, Hatcher JP, Bountra C, Michel AD, Hughes JP, et al. (2005) Disruption of the P2X₇ purinoceptor gene abolishes chronic inflammatory and neuropathic pain. *Pain* 114: 386–396.
47. Mariathasan S, Weiss DS, Newton K, McBride J, O'Rourke K, et al. (2006) Cryopyrin activates the inflammasome in response to toxins and ATP. *Nature* 440: 228–232.
48. Perregaux D, Gabel CA (1994) Interleukin-1 beta maturation and release in response to ATP and nigericin. Evidence that potassium depletion mediated by these agents is a necessary and common feature of their activity. *J Biol Chem* 269: 15195–15203.
49. Colomar A, Marty V, Combe C, Medina C, Parnet P, et al. (2003) [The immune status of Schwann cells: what is the role of the P2X₇ receptor?]. *J Soc Biol* 197: 113–122.
50. Werry EL, Liu GJ, Bennett MR (2006) Glutamate-stimulated ATP release from spinal cord astrocytes is potentiated by substance P. *J Neurochem* 99: 924–936.
51. Sperlagh B, Sershen H, Lajtha A, Vizi ES (1998) Co-release of endogenous ATP and [3H]noradrenaline from rat hypothalamic slices: origin and modulation by alpha2-adrenoceptors. *Neuroscience* 82: 511–520.
52. North RA, Verkhratsky A (2006) Purinergic transmission in the central nervous system. *Pflügers Arch* 452: 479–485.
53. Edwards FA, Gibb AJ, Colquhoun D (1992) ATP receptor-mediated synaptic currents in the central nervous system. *Nature* 359: 144–147.
54. Norenberg W, Illes P (2000) Neuronal P2X receptors: localisation and functional properties. *Naunyn Schmiedebergs Arch Pharmacol* 362: 324–339.
55. Khakh BS (2001) Molecular physiology of P2X receptors and ATP signalling at synapses. *Nat Rev Neurosci* 2: 165–174.
56. Chen L, Brosnan CF (2006) Regulation of immune response by P2X₇ receptor. *Crit Rev Immunol* 26: 499–513.
57. Di Virgilio F (2007) Purinergic signalling in the immune system. A brief update. *Purinergic Signal* 3: 1–3.
58. Barden N, Harvey M, Gagne B, Shink E, Tremblay M, et al. (2006) Analysis of single nucleotide polymorphisms in genes in the chromosome 12Q24.31 region points to P2RX7 as a susceptibility gene to bipolar affective disorder. *Am J Med Genet B Neuropsychiatr Genet* 141B: 374–382.
59. Burnstock G (2008) Purinergic signalling and disorders of the central nervous system. *Nat Rev Drug Discov* 7: 575–590.
60. Romagnoli R, Baraldi PG, Cruz-Lopez O, Lopez-Cara C, Preti D, et al. (2008) The P2X₇ receptor as a therapeutic target. *Expert Opin Ther Targets* 12: 647–661.