

Evidence for horizontal transfer of a secondary metabolite gene cluster between fungi

Nora Khaldi, Jérôme Collemare, Marc-Henri M.-H. Lebrun, Kenneth Wolfe

▶ To cite this version:

Nora Khaldi, Jérôme Collemare, Marc-Henri M.-H. Lebrun, Kenneth Wolfe. Evidence for horizontal transfer of a secondary metabolite gene cluster between fungi. Genome Biology, 2008, 9, pp.R18. 10.1186/gb-2008-9-1-r18 . hal-02666071

HAL Id: hal-02666071 https://hal.inrae.fr/hal-02666071

Submitted on 31 May 2020 $\,$

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

Research



Evidence for horizontal transfer of a secondary metabolite gene cluster between fungi Nora Khaldi^{**}, Jérôme Collemare^{**}, Marc-Henri Lebrun[†] and Kenneth H Wolfe^{*}

Addresses: *Smurfit Institute of Genetics, Trinity College Dublin, Dublin 2, Ireland. †2UMR5240 CNRS/UCB/INSA/BCS, Bayer Cropscience, 69263 Lyon cedex 09, France.

× These authors contributed equally to this work.

Correspondence: Kenneth H Wolfe. Email: khwolfe@tcd.ie

Published: 24 January 2008

Genome Biology 2008, 9:R18 (doi:10.1186/gb-2008-9-1-r18)

The electronic version of this article is the complete one and can be found online at http://genomebiology.com/2008/9/1/R18

© 2008 Khaldi et al.; licensee BioMed Central Ltd.

This is an open access article distributed under the terms of the Creative Commons Attribution License (http://creativecommons.org/licenses/by/2.0), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Received: 9 October 2007 Revised: 21 December 2007

Accepted: 24 January 2008

Abstract

Background: Filamentous fungi synthesize many secondary metabolites and are rich in genes encoding proteins involved in their biosynthesis. Genes from the same pathway are often clustered and co-expressed in particular conditions. Such secondary metabolism gene clusters evolve rapidly through multiple rearrangements, duplications and losses. It has long been suspected that clusters can be transferred horizontally between species, but few concrete examples have been described so far.

Results: In the rice blast fungus *Magnaporthe grisea*, the avirulence gene *ACE1* that codes for a hybrid polyketide synthase-nonribosomal peptide synthetase (PKS-NRPS) belongs to a cluster of 15 genes involved in secondary metabolism. Additional related clusters were detected in the ascomycetes *Chaetomium globosum*, *Stagonospora nodorum* and *Aspergillus clavatus*. Gene-by-gene phylogenetic analysis showed that in *C. globosum* and *M. grisea*, the evolution of these *ACE1*-like clusters is characterized by successive complex duplication events including tandem duplication within the *M. grisea* cluster. The phylogenetic trees also present evidence that at least five of the six genes in the homologous *ACE1* gene cluster in *A. clavatus* originated by horizontal transfer from a donor closely related to *M. grisea*.

Conclusion: The ACE1 cluster originally identified in *M. grisea* is shared by only few fungal species. Its sporadic distribution within euascomycetes is mainly explained by multiple events of duplication and losses. However, because *A. clavatus* contains an ACE1 cluster of only six genes, we propose that horizontal transfer from a relative of *M. grisea* into an ancestor of *A. clavatus* provides a much simpler explanation of the observed data than the alternative of multiple events of duplication and losses of parts of the cluster.

Background

In filamentous fungi, genes involved in the same secondary metabolite biosynthetic pathway are often located at the same locus in the genome and co-expressed, defining gene clusters [1]. Genomic clustering of genes with related cellular functions (but unrelated sequences) also occurs in other eukaryotes including mammals, nematodes and plants [2-4]. In mammals, it has been shown that clusters of co-expressed genes tend not to be rearranged among species, which indicates that natural selection can act to conserve gene order [5,6]. Similarly in fungi, natural selection seems to act to conserve gene clusters as exemplified in Aspergillus species by the cluster for the biosynthesis of aflatoxin and sterigmatocystin that has been maintained as a cluster, despite many internal rearrangements, for at least 120 million years [7,8]. The evolutionary mechanisms by which these clusters are created and maintained are unclear, but there is evidence that some instances of clustering result from strong natural selection. For example, the DAL cluster involved in nitrogen metabolism in Saccharomyces cerevisiae was formed relatively recently by a series of near-simultaneous relocations of genes that were previously scattered around the genome [9]. Other mechanisms involved in the formation and maintenance of clusters include selection for co-regulation by chromatin remodelling, epistatic selection for tight linkage between genetically interacting genes, and the "selfish operon" hypothesis of origin by horizontal gene transfer (HGT) [2,10-13]. Indeed, the clustering of the genes from a pathway at a single locus certainly facilitates HGT of genes involved in the same cellular function [10,14], increasing its likelihood.

Despite frequent speculation (reviewed in [15]), and even though some clear examples of HGT of single genes between fungal species [16] or from bacteria to fungi [17] are known, there are few reports that conclusively demonstrate HGT of a fungal secondary metabolite cluster. The strongest candidate reported so far is the epipolythiodioxopiperazine (ETP) synthase gene cluster, recently analyzed by Patron et al [18], but even in this instance alternative evolutionary scenarios can be contemplated (see Discussion). One of the best-known cases of possible HGT of a fungal secondary metabolite cluster concerns the fungal β-lactam (penicillin) antibiotic biosynthetic genes of *Penicillium* species. This proposal was originally made when bacterial and fungal isopenicillin-N-synthetases were found to have unexpectedly highly similar protein sequences [19-21]. However, subsequent phylogenetic analyses of these proteins failed to provide robust support for their HGT [22,23].

The rice blast fungus *Magnaporthe grisea* is one of the richest known fungi in terms of secondary metabolite gene clusters [24,25]. One of them contains the avirulence gene *ACE1* that encodes a hybrid polyketide synthase-nonribosomal peptide synthetase (PKS-NRPS) likely involved in the biosynthesis of an avirulence signal recognized by rice cultivars car-

rying the resistance gene Pi33 [26]. The ACE1 cluster contains 15 genes that are co-expressed specifically during the appressorium mediated penetration of the fungus into host tissues (Collemare et al, unpublished results). During annotation of the ACE1 cluster, a similar cluster was identified in the related animal pathogen Chaetomium globosum. We were then interested in identifying possible homologous clusters in other fungi in order to decipher its evolutionary history. In the present study, we combine phylogenetics and comparative genomics to identify orthologs of the M. grisea ACE1 cluster in other ascomycetes. We define a set of three genes that are shared across all instances of the cluster and hence are probably ancestral to it. This analysis revealed that the cluster in M. grisea expanded by internal duplication, and that after this duplication, part of the ACE1 cluster was likely horizontally transferred from an *M. grisea*-like ancestor into an ancestor of Aspergillus clavatus.

Results

Identification of homologous ACE1 clusters in other filamentous fungi

The ACE1 secondary metabolism gene cluster of M. grisea comprises 15 genes: ACE1 and SYN2 are PKS-NRPS hybrid genes; RAP1 and RAP2 code for enoyl reductases; CYP1-CYP4 for cytochrome P450 monoxygenases; ORFZ for an α/β hydrolase; OXR1 and OXR2 for oxidoreductases; MFS1 codes for a transporter in the MFS superfamily; BC2 codes for a binuclear zinc finger transcription factor; OME1 codes for an O-methyl transferase; and ORF3 has no homology to known proteins (Collemare *et al*, unpublished results). To find gene clusters homologous to the ACE1 cluster in other fungal species, we used an algorithm that searched 26 fungal genomes for loci where at least three likely orthologs of genes from the ACE1 cluster were linked (see Materials and methods). This search identified nine similar clusters in seven fungal species from the subphylum Pezizomycotina: three Sordariomycetes (Chaetomium globosum, Fusarium oxysporum and F. verticillioides), one Dothideomycete (Stagonospora nodorum) and three Eurotiomycetes (Aspergillus clavatus, Coccidioides immitis and Uncinocarpus reesii) (Figure 1).

Two types of clusters related to the *ACE1* cluster were identified: large clusters with eight or more genes are found in *M. grisea*, *C. globosum* and *S. nodorum*, whereas smaller clusters with three to six genes are found in the three Eurotiomycetes and in *Fusarium* species (Figure 1). *C. globosum* is unusual as its genome contains two large *ACE1*-like clusters, which we refer to as clusters 1 and 2. Similarly, the *A. clavatus* genome has two clusters as discussed below. Interestingly, a core set of three genes (homologs of *ACE1*, *RAP1* and *ORF3*; boxed in Figure 1) is present in all eight species. The presence of this core suggests that the physical linkage between these three genes is ancient and can be inferred to have existed in the common ancestor of all the genomes considered in Figure 1. As well as the genes in the eight clusters shown in Figure 1,



Figure I

ACE I and ACE I-like gene clusters in filamentous fungi. Colors indicate gene orthology in different species and paralogs in the same species. Horizontal lines indicate genes that are adjacent in the genome, with gene orientations as shown. Genomic regions are not drawn to scale. Parts A and B of the *M. grisea* cluster as identified in the text are marked. The core set of three genes inferred to have been present in the ancestral cluster are boxed. Vertical lines indicate the closest relatives of genes in the *M. grisea* cluster and one of the *A. clavatus* clusters, based on phylogenetic analyses (Figure 2 and Additional data file 1). The species phylogeny is based on the whole-genome supertree analysis of Fitzpatrick *et al* [27]; in that study the placement of Dothideomycetes relative to Sordariomycetes and Eurotiomycetes varied depending on the method of analysis, so we have shown it as a trichotomy. The analysis of Hane *et al* of the complete *S. nodorum* genome placed Dothideomycetes and Sordariomycetes in a clade with Eurotiomycetes outside [47]. Species-specific gene nomenclature is shown, except for *M. grisea* (Collemare *et al*, unpublished results). Red, green and blue coloring of species names corresponds to the labelling of individual genes from the clusters in Figure 2 and Additional data file 1.0.

we also identified a small number of single homologs of genes from the *M. grisea ACE1* cluster that are located at dispersed genomic locations in other species.

Phylogenetic analysis of the ACE1 cluster in filamentous fungi

Gene-by-gene phylogenetic analyses were carried out to decipher the evolutionary history of the loci using homologs (even at dispersed locations) of genes from *M. grisea ACE1* cluster (Figure 2 and Additional data file 1). The first trend evident from this phylogenetic analysis is that genes from clusters in Eurotiomycetes and *Fusarium* spp. are distant from those of the *M. grisea*, *C. globosum* and *S. nodorum* clusters. Indeed, genes in clusters from these last three species define clades supported by high bootstrap values (> 91%), to the exclusion of genes from Eurotiomycetes and *Fusarium* species (Figure 2a,b,e,f). Interestingly, genes from one of the *M. grisea*

ACE1 cluster than to those in ACE1-like clusters from other Eurotiomycetes (see below). In view of the gene contents of the clusters and their phylogenetic relationships, we refer to the large clusters in *M. grisea*, *C. globosum*, *S. nodorum* and the larger of the two clusters in *A. clavatus* as "ACE1 clusters", and to the smaller clusters in Eurotiomycetes and Fusarium spp. as "ACE1-like clusters". These two types of cluster have probably had a long history of independent evolution, although they certainly share a common ancestor.

We then focused on the origins of the duplicated genes in the *M. grisea* cluster. Phylogenetic trees show clearly that in *M. grisea* RAP2 is a paralog of RAP1, CYP3 is a paralog of CYP2, CYP4 is a paralog of CYP1, and SYN2 is a paralog of ACE1 (Figure 2a-d). Notably, in each of these pairs, one gene is located on the left-hand side of the *M. grisea* cluster and the other is on the right-hand side. Thus the *M. grisea* cluster appears to have undergone partial tandem duplication at



Figure 2

Maximum likelihood trees for ACE1 cluster genes and their homologs. (a) ACE1 and SYN2; (b) RAP1 and RAP2; (c) CYP1 and CYP4; (d) CYP2 and CYP3; (e) ORF3; (f) ORFZ. In each tree, genes that appear in Figure 1 are named in color or bold black. Yellow highlighting shows the five genes in the A. clavatus ACE1 cluster whose closest relatives are genes from part B of the M. grisea cluster. Bootstrap percentages are shown for all nodes. Trees were constructed from amino acid sequences as described in Methods using PHYML after alignment with ClustalW and Gblocks filtering. Trees for the other five genes in the ACE1 cluster are shown in Additional data file 1. The values of the shape parameter (α) for the gamma distribution were estimated from the data as 1.329, 1.441, 2.476, 2.615, 2.536 and 0.961 for panels a-f, respectively. The proportions of invariant sites are 0.028, 0.035, 0.030, 0.068, 0.000 and 0.000, respectively. The M. grisea SYN2 gene corresponds to parts of the automatically-annotated gene models MGG_12452.5 and MGG_12451.5.

some stage during its evolution, although the gene order is not conserved between the two parts. The presence of two *ACE1* clusters in *C. globosum* is suggestive of a second blockduplication event in this species. However, for most genes present in both *C. globosum ACE1* clusters, the copy from cluster 1 forms a clade with their *M. grisea* homologs. This close phylogenetic relationship is observed for *ACE1*, *RAP1*, *ORFZ*, *OXR1*, *CYP1*, and *OXR2*. The only exception to this pattern is *M. grisea ORF3*, which is marginally closer to the *C. globosum* cluster 2 gene, but with low bootstrap support (Figure 2 and Additional data file 1). This observation suggests that the duplication that gave rise to the current *C. globosum* clusters 1 and 2 occurred in a common ancestor of *C. globosum* and *M. grisea*, and that the corresponding cluster 2 in *M. grisea* was lost.

On the basis of this analysis, we divided the *M. grisea* cluster into two parts, A and B, so that each of the duplicated genes in *M. grisea* has one copy in part A and one in part B (Figure 1). Part A in *M. grisea* consists of nine genes, all of which have orthologs in one or both of the clusters in its closest relative *C. globosum*. The clusters in other species consist of homologs of genes from *M. grisea* part A, plus one gene from part B (*ORF3*; see Discussion). The order of the part A genes is not conserved among *M. grisea*, *C. globosum* and *S. nodorum*.

Surprisingly, this phylogenetic analysis shows that five of the six genes from part B of the M. grisea ACE1 cluster group with genes from the larger of the two clusters in A. clavatus, rather than with the genes in the more closely related (Sordariomycete) species C. globosum, or with their part A paralogs in M. grisea. Bootstrap values for grouping the M. grisea part B genes SYN2, RAP2, CYP4, CYP3 and ORF3 with their A. clavatus homologs are 98-100% (Figure 2a-e). The only gene from part B of the *M. grisea* cluster that does not group with A. clavatus is OME1 (panel e of Additional data file 1), but this is also the only gene whose detected homolog in A. clavatus (ACLA_002520) is not physically clustered with the others, which calls its orthology into question. The consistency of this phylogenetic result for part B genes, and its disagreement with the expected species relationships, are indicative of HGT between A. clavatus and part B of the M. grisea cluster. In contrast seven of the nine genes from part A of the M. grisea cluster, including ACE1 itself, lie at the expected phylogenetic position forming a clade with C. globosum (Figure 2 and Additional file 1; the two exceptions are CYP2, which is discordant but has a low bootstrap value of 66%, and MFS1, which cannot be analyzed because there is no homolog in the C. globosum clusters).

For the four panels in Figure 2 that include sequences from other Eurotiomycetes (*C. immitis* and *U. reesii*) as well as *A. clavatus*, we used the likelihood ratio test (LRT) to test whether the topologies shown (Figure 2a,b,e,f) have significantly higher likelihoods than alternative trees where the Eurotiomycetes were constrained to form a monophyletic group. In all four cases the topology shown in Figure 2 is significantly more likely than the tree expected if genes were inherited vertically (p < 0.001 for each).

Identifying the direction of gene transfer

To determine whether part B of the cluster was transferred from an *M. grisea*-like donor to an ancestor of *A. clavatus*, or vice versa, we examined phylogenetic trees constructed from those genes that have orthologs both in species that are close relatives of *M. grisea* and in species that are closer to *A. clavatus*. We would predict that if an ancestor of *A. clavatus* was the recipient of HGT, then the genes in its *ACE1* cluster would not show the expected close relationship to other Eurotiomycete species such as *C. immitis* and *U. reesii* (Figure 1), and would instead form a clade with the donor lineage (represented by *M. grisea*). Conversely, if the direction of transfer was from an *A. clavatus*-like donor into the *M. grisea* lineage, we would expect the *M. grisea* part B genes not to form a monophyletic clade with the other Sordariomycete species *C. globosum*, and instead to group with *A clavatus*.

In the phylogenetic tree of ORF3 sequences, the shared A. clavatus-M. grisea branch lies within a clade that contains homologs from the two clusters in C. globosum, as well as the Dothideomycete S. nodorum (Figure 2e). The ORF3 orthologs from C. immitis and U. reesii clearly lie outside this clade with 95% bootstrap support. Similarly, the phylogenetic tree of RAP1 and RAP2 orthologs (Figure 2b) shows that the shared branch containing the A. clavatus gene and the part B M. grisea gene (RAP2) lies within a larger clade that includes the C. globosum and M. grisea part A (RAP1) orthologs. The homologs from C. immitis and U. reesii lie outside (91% bootstrap support). Likewise, the phylogenetic tree of the ACE1-SYN2 pair (Figure 2a) places the A. clavatus sequence within a Sordariomycete/Dothideomycete clade, distant from the other Eurotiomycetes (C. immitis and U. reesii). These topologies all indicate that an ancestor of M. grisea was the donor of the transferred part B genes, and an ancestor of A. clavatus was the recipient.

ORFZ is the only gene in the *A. clavatus ACE1* cluster that does not have a homolog in part B of the *M. grisea* cluster. The origin of this gene in *A. clavatus* is not clear. Phylogenetic analysis (Figure 2f) indicates that *A. clavatus ORFZ* does not group with the *C. immitis* and *U. reesii* genes, and this conclusion is supported by the LRT. This result suggests a foreign origin for *A. clavatus ORFZ*, but the absence of a homolog in *M. grisea* part B makes it impossible to test whether this gene has a similar origin to its five neighboring genes in *A. clavatus*.

We conclude that there is phylogenetic support for the hypothesis that at least five of the six genes in the *ACE1* cluster of *A. clavatus* originated by HGT, and that the most probable single donor is a Sordariomycete ancestor related to *M. grisea*.

Discussion

The ACEI cluster is specific to few fungal species

A complete ACE1 cluster is present in only four of the 23 sequenced Pezizomycotina genomes (M. grisea, C. globosum, S. nodorum and A. clavatus). Such a sporadic distribution could be the result of either independent HGTs or frequent losses of the whole cluster in different lineages (Figure 3). We favor the latter explanation because - with the exception of A. clavatus - our phylogenetic trees of genes from the cluster have topologies that are in broad agreement with the expected species phylogeny [27]. We suggest that an ACE1like cluster consisting of at least three genes (homologous to ACE1, RAP1 and ORF3) existed in the common ancestor of Pezizomycotina, but this cluster has been lost in many lineages subsequently. The scheme in Figure 3 identifies four independent lineages (shown by dashed lines) in which all copies of the cluster have been lost. We cannot tell, with current data, whether genes such as OXR1 that are present in the ACE1 clusters of Sordariomycetes and Dothideomycetes but not in the ACE1-like clusters of Eurotiomycetes correspond to lineage-specific additions or losses.

Any tree showing apparent HGT of a gene can also be explained by an alternative scenario of gene duplications and losses. However, the situation reported here is rather different to typical cases of possible HGT of individual genes, because it involves multiple genes that are arranged as a large tandem duplication (in M. grisea). The fact that the A. clavatus ACE1 cluster forms a clade with the M. grisea part B genes (to the exclusion of the part A genes) means that the only alternative scenario to HGT is one where the part A/part B tandem duplication occurred right at the base of the tree in Figure 3. This scenario would then necessitate at least four events of precise loss of exactly one part of the tandemly duplicated set of genes: part B in C. globosum, part B in the ancestor of C. immitis and U. reesii, part B in S. nodorum, and part A in A. clavatus. Because of the precise nature of the deletion required (and choice of gene copy to delete), we do not regard this scenario as likely.

The discontinuous distribution of the ACE1 cluster among fungal species suggests that evolutionary constraints act to maintain this cluster only in few species. As M. grisea, S. nodorum and C. globosum are plant or animal pathogens, it is tempting to speculate that the ACE1 cluster is involved in the infection process of these three species. The metabolite produced by this biosynthetic pathway may be an important pathogenicity factor, but such a role remains to be determined. A. clavatus is different as it is not pathogenic. The presence of the ACE1 cluster in A. clavatus may arise from selection involving an unknown biological role of this metabolite in this fungus. Identifying the molecules made by these different clusters will be necessary to understand the role of the ACE1 cluster in fungal biology and could give clues about evolution of the ancestral biosynthetic pathway controlled by this cluster.

ACE1 cluster evolution in Sordariomycetes involved several duplication events

The ACE1 cluster has a complex history with multiple events of large-scale duplication and multiple losses. The scenario we infer is summarized in Figure 3. An ancient duplication produced the large ACE1 and smaller ACE1-like clusters. A second duplication event in an ancestral Sordariomycete gave rise to the two clusters (1 and 2) presently seen in C. globosum. This event occurred prior to the speciation between C. globosum and M. grisea, but M. grisea later lost its counterpart of cluster 2. Independently, cluster 1 underwent a tandem duplication event, generating parts A and B. This tandem duplication survived in M. grisea, but in C. globosum the addition (part B of cluster 1) was lost again. It might seem simpler to suggest that the part A/B tandem duplication was an event that occurred specifically in M. grisea after it diverged from C. globosum, but we know that this is incorrect because the part B genes from M. grisea form outgroups to a clade consisting of C. globosum and M. grisea part A genes. We can also be sure that the surviving duplications seen in M. grisea and C. globosum were separate events because of the topology of the phylogenetic trees: if the surviving genes were descended from the same duplication event we would expect that in the ACE1-SYN2 tree, for example, M. grisea ACE1 and SYN2 should each form a separate monophyletic group with one of the C. globosum genes, but that is not seen (Figure 2a). Instead we interpret the trees as indicative of two duplications of the whole cluster in a Sordariomycete ancestor of M. grisea and C. globosum, the first of which was non-tandem and the second of which was tandem. After this tandem duplication, the *M. grisea* lineage lost its ortholog of cluster 2 of *C*. globosum, and the C. globosum lineage lost its ortholog of part B of M. grisea (Figure 3). This pattern of frequent loss is consistent with the cluster's sporadic distribution in fungi.

ORF3 is unusual as it is inferred to have been present in the ancestor of all *ACE1* and *ACE1*-like clusters, but in *M. grisea* it is not duplicated and it shows phylogenetic affinity to *A. clavatus* rather than to *C. globosum* or *S. nodorum* (Figure 2e). These properties suggest that a homolog of *ORF3* was lost from part A of the *M. grisea* cluster, after the tandem duplication occurred. Furthermore, we speculate that the location of *ORF3* on the boundary between parts A and B may indicate that the tandem duplication event visible in *M. grisea* involved a recombination between two copies of this gene.

Gene order and orientation is quite poorly conserved among the *ACE1* clusters, as is typical of many secondary metabolism gene clusters [7,8,28]. This makes it all the more striking that the duplicated *M. grisea* genes each have one copy in the part A and one copy in part B. Because the tandem duplication that is evident in the *M. grisea* genome is not particularly recent (it predates the *M. grisea*/*C. globosum* speciation), we suggest that some form of selection has acted on gene order in the cluster, preventing intermixing of the two parts. In this context it is notable that recombination seems to be inhibited



Figure 3

Inferred history of ACE1 and ACE1-like clusters in filamentous fungi. The gray rectangle corresponds to the ancient core cluster of three genes (ACE1, RAP1, ORF3) that is common to all ACE1 clusters (pink) and ACE1-like clusters (orange). The black arrow denotes the inferred HGT of part B of the cluster from a donor related to *M. grisea* to the *A. clavatus* recipient. Dashed branches and smaller fonts indicate euascomycetes that were included in our analysis but lack the clusters entirely. Phylogenetic relationships are based on [27] and N Fedorova and N Khaldi, unpublished data, for the topology within the genus Aspergillus. The tree is not drawn to scale.

in the M. grisea ACE1 cluster, because it displays a low frequency of targeted gene replacement, even in a KU80 null mutant background where homologous recombination rates are increased ([29]; Collemare et al, unpublished results). The way that part A and part B genes of the ACE1 cluster are distributed among species may indicate that they are involved in the biosynthesis of different molecules. Alternatively, parts A and B of the ACE1 cluster may be each involved in the biosynthesis of independent polyketide precursors that are fused into a final complex molecule as observed for lovastatin [25,30,31]. The fact that all 15 genes in the M. grisea ACE1 cluster are co-expressed at a very specific stage of the infection process (Collemare et al, unpublished results) favors the hypothesis that both part A and part B genes are involved in same biosynthetic pathway. However, gene knockout experiments have shown that two part B genes (RAP2 and SYN2) are not essential for the avirulence function supported up to now only by the part A gene ACE1 (Collemare et al, unpublished results). These latter results suggest that part A and part B genes could be involved in the biosynthesis of two different molecules, with only one (ACE1, part A pathway) being recognized by resistant rice cultivars. However, these two hypotheses are both plausible, and await the biochemical characterization of the Ace1 metabolite.

HGT of a fungal secondary metabolism gene cluster

Although the genomics era has uncovered evidence for widespread horizontal gene transfer among prokaryotes [32,33], and from prokaryotes to eukaryotes [17,34-37] or vice versa [38,39], relatively few instances of horizontal gene transfer have been documented from one eukaryote to another [40-42]. Among fungi, the best documented is the transfer of a virulence gene from S. nodorum to Pyrenophora triticirepens, which occurred only about 70 years ago [16]. In that case, the transferred DNA fragment was about 11 kb in size but contained only one gene. In this study we showed that part B of the ACE1 cluster (30 kb in size, containing 5-6 genes) was likely horizontally transferred from a close ancestor of M. grisea (a Sordariomycete) into an ancestor of A. clavatus (a Eurotiomyete). The mechanism by which HGT might have occurred remains a matter of speculation, but could perhaps have involved hyphal fusion between species, or endocytosis. Our inference of HGT is valid only if the Sordariomycete and Eurotiomycete clades are monophyletic as shown in Figure 1, but their monophyly is supported by several molecular and systematic analyses [27,43-47].

To our knowledge, our study and the recent work of Patron *et al* [18] are the first reported instances of HGT of groups of linked genes involved in the same pathway between eukaryotic species. In both cases these secondary metabolite clusters show a punctate (sporadic) distribution among other species, with an ancestral cluster apparently having been lost by more species than the number that retain it. This pattern of frequent losses of genes and their occasional reacquisition by HGT resembles the pattern of evolution of "dispensable path-

way" genes in ascomycete yeasts [48]. Hall and Dietrich [48] noted that genes whose products function in dispensable pathways are one of the few categories of genes in S. cerevisiae that are physically organized into gene clusters. They found that the pathway for biotin synthesis was lost in a yeast ancestor and then regained in the S. cerevisiae lineage by a combination of HGT from bacteria and gene duplication with neofunctionalization. One possible explanation for this strange pattern of evolution could be that an intermediate in the pathway is toxic [48], although there is no direct experimental evidence of this. If a pathway can confer a selective advantage in some circumstances but also involves the production of a toxic intermediate, there can be strong selection in favor of the pathway in some conditions and strong selection against it in others. The consequences of such a situation could include the formation of physical gene clusters (to reduce the chances of coding for only part of the pathway, or for strong repression of transcription mediated by chromatin remodelling), and occasional selection for re-gain of function by HGT. Further exploration of this hypothesis will require the discovery of more examples of similar sets of genes, and detailed characterization of the biochemical pathways involved.

Materials and methods

We set up a local basic local alignment search tool (BLAST) database of the proteins encoded in 26 completely sequenced fungal genomes (A. niger, A. nidulans, A. terreus, A. flavus, A. oryzae, A. clavatus, N. fischeri, A. fumigatus Af293, A. fumigatus CEA10, C. immitis, C. posadasii, P. chrysogenum, U. reesii, S. sclerotiorum, F. graminearum, F. oxysporum, F. verticillioides, M. grisea, N. crassa, C. globosum, H. jecorina (T. reesei), N. haematococca (F. solani), P. chrysosporium, S. nodorum (P. nodorum), C.neoformans, U. maydis). To find candidate ACE1-like clusters in other fungi, we used a two-step process outlined below.

In the first step, each protein encoded by the *M. grisea ACE1* cluster was used as a query in protein-protein BLAST (BLASTP) searches against this database, and for each query the top 25 hits were retained provided that their E-values were less than 1e-4. Each set of proteins was aligned using ClustalW [49] and poorly aligned regions were removed using Gblocks [50]. Sequence alignments are available as Additional data file 2. Maximum likelihood trees were constructed using PHYML [51] with the JTT amino acid substitution matrix and four categories of substitution rates. Bootstrapping was done using the default options in PHYML with 100 replicates per run. To avoid long branch attraction problems we withdrew highly divergent sequences and repeated the alignment and tree reconstruction steps on the new sets. We also verified at each step that the alignment obtained after running Gblocks represented at least 30% of the initial protein sequence. Genes were considered as orthologs of an M. *grisea ACE1* cluster gene if they grouped in a monophyletic clade with a bootstrap support of \geq 70%.

Many of the genes identified in this first step were located in gene clusters. For each cluster so identified (defined as the presence of at least two homologs of *M. grisea ACE1* cluster genes adjacent to one another) we then made a second step of analysis, examining any other genes that are physically located within these clusters but which were not picked up at the first step (either because their BLASTP E-values were too weak, or because they were not in the top 25 hits when the database was searched). This process added genes CHG05286.1, CHG05287.1, SNU00307.1 and FVEG_12610 to the analyses.

Abbreviations

BLAST, basic local alignment search tool; HGT, horizontal gene transfer; LRT, likelihood ratio test.

Authors' contributions

JC and MHL isolated the *M. grisea ACE1* cluster and identified initial evidence of HGT. NK and JC conducted genome searches and phylogenetic analyses. KHW drew the figures. All authors contributed to writing the manuscript. All authors read and approved the final manuscript.

Additional data files

The following additional data are available with the online version of this article: a figure (Additional data file 1) showing maximum likelihood trees for the *ACE1* cluster genes that are not included in Figure 2 (*OXR1*, *BC2*, *OXR2*, *MFS1* and *OME1*), and a data file (Additional data file 2) containing the sequence alignments used to produce Figure 2 and Additional data file 1.

Acknowledgements

NK and KHW are supported by Science Foundation Ireland. MHL and JC are supported by CNRS, and Bayer Cropscience, France.

References

- Keller NP, Hohn TM: Metabolic pathway gene clusters in filamentous fungi. Fungal Genet Biol 1997, 21:17-29.
- Hurst LD, Pal C, Lercher MJ: The evolutionary dynamics of eukaryotic gene order. Nat Rev Genet 2004, 5:299-310.
- Qi X, Bakht S, Leggett M, Maxwell C, Melton R, Osbourn A: A gene cluster for secondary metabolism in oat: implications for the evolution of metabolic diversity in plants. Proc Natl Acad Sci USA 2004, 101:8233-8238.
- Shimura K, Okada A, Okada K, Jikumaru Y, Ko KW, Toyomasu T, Sassa T, Hasegawa M, Kodama O, Shibuya N, Koga J, Nojiri H, Yamane H: Identification of a biosynthetic gene cluster in rice for momilactones. J Biol Chem 2007, 282:34013-34018.
- Singer GAC, Lloyd AT, Huminiecki LB, Wolfe KH: Clusters of coexpressed genes in mammalian genomes are conserved by natural selection. *Mol Biol Evol* 2005, 22:767-775.
- 6. Semon M, Duret L: Evolutionary origin and maintenance of

coexpressed gene clusters in mammals. Mol Biol Evol 2006, 23:1715-1723.

- Cary JW, Chang P-K, Bhatnagar D: Clustered metabolic pathway genes in filamentous fungi. In Applied Mycology and Biotechnology, Agriculture and Food Production Volume 1. Edited by: Khachatourians GG, Arora DK. Amsterdam: Elsevier; 2001:165-198.
- Cary JW, Ehrlich KC: Aflatoxigenicity in Aspergillus: molecular genetics, phylogenetic relationships and evolutionary implications. Mycopathologia 2006, 162:167-177.
- 9. Wong S, Wolfe KH: Birth of a metabolic gene cluster in yeast by adaptive gene relocation. *Nat Genet* 2005, **37**:777-782.
- Lawrence J: Selfish operons: the evolutionary impact of gene clustering in prokaryotes and eukaryotes. Curr Opin Genet Devel 1999, 9:642-648.
- Keller NP, Turner G, Bennett JW: Fungal secondary metabolism - from biochemistry to genomics. Nat Rev Microbiol 2005, 3:937-947.
- 12. Batada NN, Urrutia AO, Hurst LD: Chromatin remodelling is a major source of coexpression of linked genes in yeast. *Trends Genet* 2007, 23:480-484.
- Batada NN, Hurst LD: Evolution of chromosome organization driven by selection for reduced gene expression noise. Nat Genet 2007, 39:945-949.
- 14. Walton JD: Horizontal gene transfer and the evolution of secondary metabolite gene clusters in fungi: an hypothesis. *Fun*gal Genet Biol 2000, **30**:167-171.
- 15. Rosewich UL, Kistler HC: Role of horizontal gene transfer in the evolution of fungi. Annu Rev Phytopathol 2000, 38:325-363.
- Friesen TL, Stukenbrock EH, Liu Z, Meinhardt S, Ling H, Faris JD, Rasmussen JB, Solomon PS, McDonald BA, Oliver RP: Emergence of a new disease as a result of interspecific virulence gene transfer. Nat Genet 2006, 38:953-956.
- Wenzl P, Wong L, Kwang-Won K, Jefferson RA: A functional screen identifies lateral transfer of beta-glucuronidase (gus) from bacteria to fungi. Mol Biol Evol 2005, 22:308-316.
- Patron NJ, Waller RF, Cozijnsen AJ, Straney DC, Gardiner DM, Nierman WC, Howlett BJ: Origin and distribution of epipolythiodioxopiperazine (ETP) gene clusters in filamentous ascomycetes. BMC Evol Biol 2007, 7:174.
- Weigel BJ, Burgett SG, Chen VJ, Skatrud PL, Frolik CA, Queener SW, Ingolia TD: Cloning and expression in Escherichia coli of isopenicillin-N-synthetase genes from Streptomyces lipmanii and Aspergillus nidulans. J Bacteriol 1988, 170:3817-3826.
- 20. Aharonowitz Y, Cohen G, Martin JF: Penicillin and cephalosporin biosynthetic genes: structure, organization, regulation, and evolution. Ann Rev Microbiol 1992, 46:461-495.
- 21. Liras P, Martin JF: Gene clusters for beta-lactam antibiotics and control of their expression: why have clusters evolved, and from where did they originate? *Int Microbiol* 2006, **9**:9-19.
- Smith MW, Feng DF, Doolittle RF: Evolution by acquisition: the case for horizontal gene transfers. Trends Biochem Sci 1992, 17:489-493.
- Buades C, Moya A: Phylogenetic analysis of the isopenicillin-N -synthetase horizontal gene transfer. J Mol Evol 1996, 42:537-542.
- Dean RA, Talbot NJ, Ebbole DJ, Farman ML, Mitchell TK, Orbach MJ, Thon M, Kulkarni R, Xu JR, Pan H, Read ND, Lee Y-H, Carbone I, Brown D, Yee Oh Y, Donofrio N, Jeong JS, Soanes DM, Djonovic S, Kolomiets E, Rehmeyer C, Li W, Harding M, Kim S, Lebrun M-H, Bohnert H, Coughlan S, Butler J, Calvo S, Ma L-J, Nicol R, Purcell S, Nusbaum C, Galagan JE, Birren BW: The genome sequence of the rice blast fungus Magnaporthe grisea. Nature 2005, 434:980-986.
- Collemare J, Billard A, Böhnert H, Lebrun M-H: Secondary metabolism of the rice blast fungus Magnaporthe grisea: the role of hybrid PKS-NRPS in pathogenicity. Mycol Res 2007. doi: 10.1016/j.mycres.200708.003
- Bohnert HÚ, Fudal I, Dioh W, Tharreau D, Notteghem JL, Lebrun MH: A putative polyketide synthase/peptide synthetase from Magnaporthe grisea signals pathogen attack to resistant rice. Plant Cell 2004, 16:2499-2513.
- 27. Fitzpatrick DA, Logue ME, Stajich JE, Butler G: A fungal phylogeny based on 42 complete genomes derived from supertree and combined gene analysis. *BMC Evol Biol* 2006, 6:99.
- Gardiner DM, Cozijnsen AJ, Wilson LM, Pedras MS, Howlett BJ: The sirodesmin biosynthetic gene cluster of the plant pathogenic fungus Leptosphaeria maculans. Mol Microbiol 2004, 53:1307-1318.
- 29. Villalba F, Collemare J, Landraud P, Lambou K, Brozek V, Cirer B,

Morin D, Bruel C, Beffa R, Lebrun MH: Improved gene targeting in *Magnaporthe grisea* by inactivation of *MgKU80* required for non-homologous end joining. *Fungal Genet Biol* 2007. 2000, **17:**540-552.

- Guindon S, Gascuel O: A simple, fast, and accurate algorithm to estimate large phylogenies by maximum likelihood. Syst Biol 2003, 52:696-704.
- Kennedy J, Auclair K, Kendrew SG, Park C, Vederas JC, Hutchinson CR: Modulation of polyketide synthase activity by accessory proteins during lovastatin biosynthesis. Science 1999, 284:1368-1372.
- Hutchinson CR, Kennedy J, Park C, Kendrew S, Auclair K, Vederas J: Aspects of the biosynthesis of non-aromatic fungal polyketides by iterative polyketide synthases. Antonie Van Leeuwenhoek 2000, 78:287-295.
- 32. Doolittle WF: **Phylogenetic classification and the universal** tree. *Science* 1999, **284**:2124-2129.
- 33. Beiko RG, Harlow TJ, Ragan MA: Highways of gene sharing in prokaryotes. Proc Natl Acad Sci USA 2005, 102:14332-14337.
- Kondo N, Nikoh N, Ijichi N, Shimada M, Fukatsu T: Genome fragment of Wolbachia endosymbiont transferred to X chromosome of host insect. Proc Natl Acad Sci USA 2002, 99:14280-14285.
- Dunning Hotopp JC, Clark ME, Oliveira DC, Foster JM, Fischer P, Torres MC, Giebel JD, Kumar N, Ishmael N, Wang S, Ingram J, Nene RV, Shepard J, Tomkins J, Richards S, Spiro DJ, Ghedin E, Slatko BE, Tettelin H, Werren JH: Widespread lateral gene transfer from intracellular bacteria to multicellular eukaryotes. Science 2007, 317:1753-1756.
- Hall C, Brachat S, Dietrich FS: Contribution of horizontal gene transfer to the evolution of Saccharomyces cerevisiae. Eukaryot Cell 2005, 4:1102-1115.
- Woolfit M, Rozpedowska E, Piskur J, Wolfe KH: Genome survey sequencing of the wine spoilage yeast Dekkera (Brettanomyces) bruxellensis. Eukaryot Cell 2007, 6:721-733.
- Guljamow A, Jenke-Kodama H, Saumweber H, Quillardet P, Frangeul L, Castets AM, Bouchier C, Tandeau de Marsac N, Dittmann E: Horizontal gene transfer of two cytoskeletal elements from a eukarvote to a cvanobacterium. *Curr Biol* 2007. 17:R757-759
- eukaryote to a cyanobacterium. Curr Biol 2007, 17:R757-759.
 39. Rogers MB, Patron NJ, Keeling PJ: Horizontal transfer of a eukaryotic plastid-targeted protein gene to cyanobacteria. BMC Biol 2007, 5:26.
- Andersson JO: Lateral gene transfer in eukaryotes. Cell Mol Life Sci 2005, 62:1182-1197.
- Richards TA, Dacks JB, Jenkinson JM, Thornton CR, Talbot NJ: Evolution of filamentous plant pathogens: gene exchange across eukaryotic kingdoms. Curr Biol 2006, 16:1857-1864.
- Andersson JO, Sjogren AM, Horner DS, Murphy CA, Dyal PL, Svard SG, Logsdon JM Jr, Ragan MA, Hirt RP, Roger AJ: A genomic survey of the fish parasite Spironucleus salmonicida indicates genomic plasticity among diplomonads and significant lateral gene transfer in eukaryote genome evolution. BMC Genomics 2007, 8:51.
- James TY, Kauff F, Schoch CL, Matheny PB, Hofstetter V, Cox CJ, Celio G, Gueidan C, Fraker E, Miadlikowska J, Lumbsch HT, Rauhut A, Reeb V, Arnold AE, Amtoft A, Stajich JE, Hosaka K, Sung G-H, Johnson D, O'Rourke B, Crockett M, Binder M, Curtis JM, Slot JC, Wang Z, Wilson AW, Schüler A, Longcore JE, O'Donnell K, et al.: Reconstructing the early evolution of fungi using a six-gene phylogeny. Nature 2006, 443:818-822.
- Blackwell M, Hibbett DS, Taylor JW, Spatafora JW: Research coordination networks: a phylogeny for kingdom Fungi (deep hypha). Mycologia 2006, 98:829-837.
- Kuramae EE, Robert V, Snel B, Weiss M, Boekhout T: Phylogenomics reveal a robust fungal tree of life. FEMS Yeast Res 2006, 6:1213-1220.
- Robbertse B, Reeves JB, Schoch CL, Spatafora JW: A phylogenomic analysis of the Ascomycota. Fungal Genet Biol 2006, 43:715-725.
- 47. Hane JK, Lowe RG, Solomon PS, Tan KC, Schoch CL, Spatafora JW, Crous PW, Kodira C, Birren BW, Galagan JE, Torrianie SFF, McDonalde BA, Oliver RP: Dothideomycete plant interactions illuminated by genome sequencing and EST analysis of the wheat pathogen Stagonospora nodorum. Plant Cell 2007, 19:3347-3368.
- Hall C, Dietrich FS: The reacquisition of biotin prototrophy in Saccharomyces cerevisiae involved horizontal gene transfer, gene duplication and gene clustering. Genetics 2007, 177:2293-2307.
- Thompson JD, Higgins DG, Gibson TJ: CLUSTAL W: improving the sensitivity of progressive multiple sequence alignment through sequence weighting, position-specific gap penalties and weight matrix choice. Nucleic Acids Res 1994, 22:4673-4680.
- 50. Castresana J: Selection of conserved blocks from multiple alignments for their use in phylogenetic analysis. Mol Biol Evol