



HAL
open science

Ectopic expression of VvMybPA2 promotes proanthocyanidin biosynthesis in grapevine and suggests additional targets in the pathway

Nancy N. Terrier, Laurent Torregrosa, Agnes Ageorges, Sandrine S. Vialet, Clotilde Verries, Veronique V. Cheynier, Charles Romieu

► To cite this version:

Nancy N. Terrier, Laurent Torregrosa, Agnes Ageorges, Sandrine S. Vialet, Clotilde Verries, et al.. Ectopic expression of VvMybPA2 promotes proanthocyanidin biosynthesis in grapevine and suggests additional targets in the pathway. *Plant Physiology*, 2009, 149 (2), pp.1028-1041. 10.1104/pp.108.131862 . hal-02666928

HAL Id: hal-02666928

<https://hal.inrae.fr/hal-02666928>

Submitted on 31 May 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Ectopic Expression of VvMybPA2 Promotes Proanthocyanidin Biosynthesis in Grapevine and Suggests Additional Targets in the Pathway^{1[W][OA]}

Nancy Terrier*, Laurent Torregrosa, Agnès Ageorges, Sandrine Vialet, Clotilde Verriès, Véronique Cheyrier, and Charles Romieu

UMR SPO 1083 (N.T., A.A., S.V., C.V., V.C.) and UMR DIAPC 1097 (L.T., C.R.), Campus SupAgro-INRA, F-34060 Montpellier, France

Grapevine (*Vitis vinifera*) proanthocyanidins contribute to plant defense mechanisms against biotic stress and also play a critical role in organoleptic properties of wine. In grapevine berry, these compounds are mainly accumulated in exocarps and seeds in the very early stages of development. A previous study has already identified VvMybPA1 as the first transcription factor involved in the regulation of the proanthocyanidin pathway during seed development in grapevine. A novel Myb factor, VvMybPA2, which is described in this study, is in contrast mainly expressed in the exocarp of young berries and in the leaves. This transcription factor shows very high protein sequence homology with other plant Myb factors, which regulate flavonoid biosynthesis. Ectopic expression of either *VvMybPA1* or *VvMybPA2* in grapevine hairy roots induced qualitative and quantitative changes of the proanthocyanidin profiles. High-throughput transcriptomic analyses of transformed grapevine organs identified a large set of putative targets of the *VvMybPA1* and *VvMybPA2* transcription factors. Both genes significantly activated enzymes of the flavonoid pathway, including anthocyanidin reductase and leucoanthocyanidin reductase 1, the specific terminal steps in the biosynthesis of epicatechin and catechin, respectively, but not leucoanthocyanidin reductase 2. The functional annotation of the genes whose expression was modified revealed putative new actors of the proanthocyanidin pathway, such as glucosyltransferases and transporters.

Flavonoids are a family of plant secondary metabolites that comprise several groups of compounds (e.g. anthocyanins, flavonols, and flavan 3-ols) and accumulate in a wide variety of plant tissues, where they are involved in diverse functions. In particular, flavonols play a role in protection against UV radiation (Winkel-Shirley, 2002), proanthocyanidins (PAs; i.e. flavan 3-ol oligomers and polymers) protect plants against microbial attacks and fungal growth (Dixon et al., 2005), and anthocyanins of flowers and fruits attract pollinators and help to disseminate seeds (Grotewold, 2006).

Grapevine (*Vitis vinifera*) flavonoids are also of particular importance for wine quality: anthocyanins of red-skinned cultivars are responsible for the red wine color, while PAs or so-called “condensed tannins” confer its astringency to the wine.

In grapevine fruit, flavan 3-ols are present in skin and seed tissues (Kennedy et al., 2001) and to a lesser extent in the pulp (Mané et al., 2007; Verriès et al., 2008). In pericarp, flavan 3-ols accumulate mainly before *véraison* (i.e. induction of ripening; Kennedy et al., 2001; Verriès et al., 2008), whereas anthocyanin accumulation in skin is restricted to ripening. In vegetative organs, PA content constantly increases during leaf development, but the rate of synthesis decreases in old leaves (Bogs et al., 2005). Grapevine flavan 3-ols are found as monomers, oligomers, and polymers. The major flavan 3-ol monomers in grapes are (+)-catechin and its isomer, (–)-epicatechin, and to a lesser extent (–)-epicatechin 3-O-gallate, the gallic ester of (–)-epicatechin (Su and Singleton, 1969). Grapevine seed PAs are partly galloylated procyanidins, based on (+)-catechin and (–)-epicatechin units, with mean degrees of polymerization (mDP) lower than 10 (Prieur et al., 1994). Grapevine skin PAs have a higher mDP around 30. They contain (–)-epicatechin and (–)-epigallocatechin (trihydroxylated on the B ring) as their major extension units (EU) and are thus mixed procyanidin/prodelphinidin polymers with a few percentage of galloylated units (Souquet et al., 1996). Catechin is the major isomer found as free monomer or terminal unit (TU) in grape skin PAs (Souquet et al., 1996).

In plants, several structural genes of PA biosynthesis have already been identified. Genes encoding anthocyanidin reductase (ANR) and leucoanthocyanidin reductase (LAR) were shown to catalyze, respectively,

¹ This work was supported by the European Community (STREP project FLAVO-FOOD-CT-2004-513960).

* Corresponding author; e-mail terrier@supagro.inra.fr.

The author responsible for distribution of materials integral to the findings presented in this article in accordance with the policy described in the Instructions for Authors (www.plantphysiol.org) is: Nancy Terrier (terrier@supagro.inra.fr).

^[W] The online version of this article contains Web-only data.

^[OA] Open Access articles can be viewed online without a subscription.

www.plantphysiol.org/cgi/doi/10.1104/pp.108.131862

the synthesis of epicatechin and catechin as monomer (Devic et al., 1999; Tanner et al., 2003). In grapevine, two isogenes of *LAR* (*LAR1* and *LAR2*) and one gene of *ANR* have been identified (Bogs et al., 2005). Production of a *LAR1* *Escherichia coli* recombinant protein confirmed its role in catechin synthesis (Bogs et al., 2005). A *LAR2* yeast recombinant protein accepts the three leucoanthocyanidins (mono-, di-, and trihydroxylated on the B ring) as substrates (Pfeiffer et al., 2006). The expression of both *ANR* and *LAR* genes is high in young berries and then constantly decreases during fruit development. *LAR1* expression was almost restricted to seeds, whereas *LAR2* was expressed in both skin and seeds (Bogs et al., 2005). *ANR* transcripts and, to a lesser extent, *LAR2* transcripts were also detected in the pulp of green berries (Verriès et al., 2008). In vegetative organs, *ANR* was expressed at high levels throughout leaf development, *LAR2* expression was only detected in mature leaves, and *LAR1* was hardly expressed in leaves (Bogs et al., 2005).

Despite this recent progress in the knowledge of PA unit synthesis, the mechanisms involved in either polymerization or galloylation of units and the fine regulation of the spatiotemporal PA composition remain to be elucidated. At least six transcription factors, belonging to Myb, bHLH, WD40, WRKY, zinc finger, and MADS box proteins, were found to regulate PA biosynthesis in *Arabidopsis* (*Arabidopsis thaliana*). TT2, TT8, and TTG1, encoding Myb, bHLH, and WD40 proteins, respectively, were found to be associated in a transcriptional complex driving the expression of flavonoid structural genes (Lepiniec et al., 2006).

In grapevine, several Myb transcription factors controlling different branches of the flavonoid pathway have been identified: Myb5a for the general flavonoid pathway (Deluc et al., 2006), MybA1 (Kobayashi et al., 2002, 2004) and MybA2 (Bogs et al., 2007; Walker et al., 2007) for the anthocyanin pathway, and MybPA1 (Bogs et al., 2007) for the PA pathway. VvMybPA1 was shown to be able to promote both *VvANR* and *VvLAR1* and to complement *Arabidopsis tt2* mutants (Bogs et al., 2007). However, in the absence of a definitive argument from homologous reverse genetics, these data are still speculative.

The important number of ESTs for grapevine and the recent release of the grapevine genome sequence (Jaillon et al., 2007; Velasco et al., 2007) pave the way for a more exhaustive screening of the additional factors involved in the regulation of PA accumulation. We present here the identification and functional validation of VvMybPA2 as a new transcription factor for the regulation of PA biosynthesis. VvMybPA2 presents very high protein sequence homology with other plant Myb factors regulating the flavonoid pathway, including AtTT2. Its expression pattern fits well with PA accumulation during the development of grapevine fruit tissues. Functional properties of VvMybPA2 were studied and compared with those of VvMybPA1 through their ectopic expression in a homologous

system. The PA content of grapevine organs ectopically overexpressing either *VvMybPA1* or *VvMybPA2* was dramatically increased and showed noticeable composition changes. High-throughput comparative transcriptomic analysis revealed new putative targets of these Mybs in the PA pathway, possibly involved in gene regulation, PA modifications, and transport.

RESULTS

VvMybPA2 Encodes a Myb Domain Protein

Three ESTs, named EC939601, EC993595, and EC950762, were obtained by BLAST search in the *Vitis* database using AtTT2 as a query sequence. The sequences were assembled in a contig, and the full-length cDNA was amplified from cDNA of 3-week-postflowering Shiraz berries, a developmental stage corresponding to the maximum rate of PA biosynthesis. The sequence recovered, called *VvMybPA2* (accession no. EU919682), encodes a protein of 284 amino acid residues (Fig. 1) with a predicted molecular mass of 31.8 kD and a calculated pI of 5.45. In the grapevine PN 40024 genome sequence, this gene is located on chromosome 11 (Jaillon et al., 2007). In this region, gene GSVIVT00016470001 was predicted. However, the in silico predicted coding region of GSVIVT00016470001 is shorter than the *VvMybPA2* actual one experimentally observed. This difference is due to a failure in the automatic detection procedure of intron/exon boundaries used for the annotation of the whole genome.

A phylogenetic tree was constructed using the neighbor-joining method with the protein sequences of other plant Myb factors involved in the regulation of the phenylpropanoid and flavonoid pathways (Fig. 2), like phenylpropanoid volatiles (PhODO1; Verdonk et al., 2005), flavonols (AtMyb112, AtMyb11, and AtMyb12; Stracke et al., 2007), anthocyanins (VvMybA1 and VvMybA2 [Kobayashi et al., 2002], AtPAP1 and AtPAP2 [Borevitz et al., 2000], LeANT1 [Mathews et al., 2003], MdMyb10 [Espley et al., 2007], and MdMyb1 [Talos et al., 2006]), PAs (VvMybPA1 [Bogs et al., 2007] and AtTT2 [Nesi et al., 2001]), vacuolar pH (PH4; Quattrocchio et al., 2006), or Myb whose function is still undetermined (OsMyb3, PmMBF1). This tree reveals that VvMybPA2 is very distinct from regulators of the anthocyanin pathway and appears to be more closely related to AtTT2 than the other grapevine Myb factors identified so far. Analyses of the primary structure (motif 1; Fig. 1) revealed that the R2R3 repeat region of VvMybPA2 is highly homologous to other plant MYB factors and contains the residues of a conserved signature sequence, [D/E]Lx2[R/K]x3Lx6Lx3R, for interaction with bHLH proteins (Grotewold et al., 2000). The nine-residue consensus sequence V(I/V)R(T/P)(K/R)A(I/L/V)(R/K)C is present in both AtTT2 and OsMYB3 in the highly variable C-terminal region




Figure 1. Multialignment of *VvMybPA2* with related MYB proteins by ClustalW: *Arabidopsis* *AtMyb123* (TT2; Q9FJA2), rice (*Oryza sativa*) *OsMYB3* (BAA23339), and grapevine *VvMYB5b* (AAX51291) and *VvMybPA1* (CAJ90831). Identical amino acids are indicated in black, and similar amino acids are indicated in gray. The R2 and R3 repeats of the MYB domain, the motif participating in the interaction with bHLH proteins (motif 1), and the C-terminal conserved motif (motif 2) are indicated.

(Nesi et al., 2001; Stracke et al., 2001). It is noticeably conserved in *VvMybPA2* (motif 2; Fig. 1) while being absent from *VvMybPA1* and *VvMyb5a* (Deluc et al., 2006; Bogs et al., 2007).

Expression Profiling of *VvMybPA2*

The spatiotemporal expression of *VvMybPA2* was evaluated by real-time PCR with RNA isolated from several vegetative tissues and berries sampled at different stages of development. Figure 3A shows that *VvMybPA2* was highly expressed in very young berries shortly after anthesis. Then, transcript abundance decreased to a very low level after *véraison*. When berries were divided into exocarp, mesocarp, and seed at three stages of development, *VvMybPA2* expression was mostly restricted to the exocarp of very young berries (Fig. 3B). In vegetative organs, the maximum expression of *VvMybPA2* was detected in leaves, especially in younger ones (Fig. 3C).

Functional Characterization of *VvMybPA1* and *VvMybPA2*

In order to establish the function of *VvMybPA1* and *VvMybPA2*, each full-length cDNA driven by the 35S promoter was separately introduced into grapevine hairy roots. Hairy roots were screened by PCR for the presence of the hygromycin phosphotransferase gene from the pH2GW7 backbone, yielding seven positive independent transgenic lines from the 10 plants inoculated with each construct. Ectopic expression of

VvMybPA1 and *VvMybPA2* transcription factor led to PA accumulation. For each gene, the two lines with the highest PA content were selected as independent biological duplicates for more detailed phenotypic and transcriptomic analysis. Hairy roots without the *VvMybPA1/2* transgene were used as a wild-type control.

Flavonoid and Lignin Content in Grapevine Organs Overexpressing *VvMybPA1* and *VvMybPA2*

VvMybPA1- and *VvMybPA2*-expressing hairy roots contained around 8 mg PA g⁻¹ fresh weight, which represents a 5-fold greater accumulation than the levels found in control lines (Fig. 4). Whereas PAs of wild-type roots do not contain any B-ring trihydroxylated units, the percentage of epigallocatechin reached 4% and 5% in roots expressing *VvMybPA1* and *VvMybPA2*, respectively. The mDP exhibited a slight increase in *VvMybPA2*-expressing lines, with mean values of 16.7 (wild type = 11.6). The high heterogeneity between the two analyzed control lines prevents us from drawing any conclusion concerning the possible influence of the Myb factors on the galloylation level. Neither anthocyanin nor flavonol was detected.

Coloration of grapevine root organs with dimethylaminocinnamaldehyde (DMACA), which reacts with PAs to form a blue stain, clearly shows that PAs were not uniformly located in the hairy root tissues (Fig. 5). PAs were quite abundant in the youngest part of the roots (i.e. in the cell division area below the apex). In




Figure 2. Phylogenetic tree showing selected plant MYB transcription factors retrieved from public databases. The phylogenetic tree was constructed from the Clustal alignment using the neighbor-joining method in the MEGA4 package. The scale bar represents 0.05 substitutions per site. The GenBank accession numbers of the MYB proteins are as follows: AtMYB12 (NP_182268), AtMYB11 (NP_191820), AtMYB111 (NP_199744), ZmP (P27898), PhPH4 (AAY52377), VvMyb5b (AAX51291), VvMYB5a (AAS68190), VvMybPA1 (CAJ90831), PmMBF1 (AAA82943), ZmC1 (AAA33482), ZmPL (AAA19821), VvMybPA2 (EU919682), OsMYB3 (BAA23339), AtTT2 (Q9FJA2), BnTT2-1 (ABI13038), BnTT2-2 (ABI13039), BnTT2-3 (ABI13040), FaMYB1 (AAK84064), AmMIXTA (CAA55725), AtMyb4 (BAA21619), PhODO1 (AAV98200), AtGL1 (NP_189430), AtWER (AAF18939), MdMYB1 (ABK58136), MdMYB10 (ABB84753), AtPAP2 (NP_176813), AtPAP1 (NP_176057), LeANT1 (AAQ55181), VvMYBA1 (BAD18977), and VvMYBA2 (BAD18978).

the sectors accumulating high level of PAs, epidermis, endoderm, and vascular bundles were found to be the richest. Pericycle was less marked by the DMACA, and almost no coloration could be observed in the cortical parenchyma cells whatever the type of roots.

With the objective to evaluate whether the transformation induced some redirection in the phenolic me-

tabolism, preliminary analyses of lignin content and composition of the hairy root samples were performed by thioacidolysis (Lapierre et al., 1995) and are presented in Supplemental Table S1. The total yields of lignin-derived thioacidolysis guaiacyl and syringyl monomers were similarly low, whatever the samples (ranging between 8 and 14 $\mu\text{mol g}^{-1}$ dry weight), and the nonmethoxylated H monomers were recovered as trace components. This yield suggests that all of the hairy root samples have a similarly low lignin content.




Figure 3. Transcript levels of *VvMybPA2* during berry pericarp development (A), in different berry tissues at three developmental stages (B), and in different organs of vine plants (C). *Véraison* (V) is marked with the arrows. Gene expression was determined by real-time PCR and normalized with the expression of *EF1 α* . All data are means of three replicates, with error bars indicating sd.




Figure 4. PA contents and mean characteristics (mDP, percentage galloylation, percentage trihydroxylation) in grapevine organs over-expressing *VvMybPA1* (lines PA1-2A and PA1-7A) or *VvMybPA2* (lines PA2-1A and PA2-7A) versus the wild type (lines control-1 and control-2). Data are means \pm SD of triplicate extractions. Stars indicate that values are significantly different between control and transformed hairy roots ($P < 0.01$). FW, Fresh weight.

By contrast, the molar frequency of the monomethoxylated guaiacyl thioacidolysis monomers was found to be increased, particularly in the hairy roots of the *VvMybPA2*-expressing lines. This result suggests that

the formation of the guaiacyl lignin units might be affected by the transformation, whereas the total lignin content is not changed.

Expression Analysis of Putative Targets

In order to decipher the mechanisms that underlie the phenotypic changes observed in grapevine ectopically expressing *VvMybPA1* and *VvMybPA2*, we first performed real-time PCR experiments on previously known genes of grapevine PA metabolism (Fig. 6A). Overexpression of *VvMybPA1* and *VvMybPA2* transgenes was confirmed in the respective transformed lines. Some direct or indirect activation was found, as endogenous *VvMybPA1* was slightly induced by 35S:: *VvMybPA2*, while the converse was not true concerning *VvMybPA2*.

ANR transcripts were significantly increased by *VvMybPA1* and *VvMybPA2* overexpression. Similar results were observed with *LAR1*, except that its induction by *VvMybPA2* was slightly lower (+65% compared with wild-type organs) and not significant. *LAR2* expression was not significantly modified upon overexpression of the *VvMybPA* factors when compared with controls.

Global Transcriptome Response Analysis Induced by *VvMybPA1/2* Overexpression

Transcriptome analysis was performed with a 14 K microarray to identify new genes involved in PA biosynthesis and particularly the putative targets of *VvMybPA1* and *VvMybPA2*. The design corresponded to a comparison analysis of two couples of independent biological replicates of wild-type/over-expressing organs for both *VvMybPA1* and *VvMybPA2* (comprehensive data are available as Supplemental Materials and Methods S1).

A *t* test ($P < 0.01$ based on permutation) revealed that only 2% to 3% of the total number of oligonucleotides spotted on the array presented significant variations due to *VvMybPA1* and *VvMybPA2* overexpression (510 and 371, respectively). For *VvMybPA1*, 305 oligonucleotides presented an increase in their hybridization signal and 205 presented a decrease. In the case of *VvMybPA2*, the respective increase and decrease were 158 and 213 (Supplemental Materials and Methods S1). Among the transcripts whose abundance increased, 55 are common to the *VvMybPA1*- and *VvMybPA2*-overexpressing organs (Table I). Further analysis revealed that this set of 55 genes corresponds to only 51 unigenes, as four couples of oligonucleotides may hybridize the same transcript. Although significant, the induction ratios in organs ectopically expressing *VvMybPA* were rather low, as they ranged from 1.1 to 3.7.

Among the genes induced in common by *VvMybPA1* and *VvMybPA2*, 10 are linked to the PA pathway, including nine genes previously identified in grape-




Figure 5. Phenotype of grapevine hairy roots stained with DMACA. A, Wild-type hairy roots showing the distribution of blue dye formed with PA on main and secondary organs. Bar = 2 mm. B, Wild-type hairy root section. Bar = 250 μ m. C, *VvMybPA1*-expressing hairy root. Bar = 250 μ m. D, *VvMybPA2*-expressing hairy root. Bar = 250 μ m. DZ, Division zone; EZ, elongation zone; SR, secondary roots.

vine: Phe ammonia lyase, 4-coumarate:CoA ligase, chalcone synthase, two genes of flavanone-3-hydroxylase, flavonoid-3'-hydroxylase (F3'H), dihydroflavonol-4-reductase, leucoanthocyanidin dioxygenase (LDOX), ANR, and a multidrug and toxic compound extrusion (MATE) transporter (GSVIVP00018839001) exhibiting homology with TT12 (Debeaujon et al., 2001) that represents a new putative actor in the PA pathway in grapevine. Two isogenes encoding cinnamoyl-CoA reductase of the phenylpropanoid pathway were also identified. Both transcription factors led to the

induction of four genes linked to the metabolism of aromatic amino acids and shikimate (shikimate kinase, shikimate dehydrogenase, 3-deoxy-D-arabinoheptulosonate 7-phosphate synthase, and prepheate dehydratase), precursors of the flavonoid pathway. In addition to genes encoding enzymes already annotated as actors in the flavonoid pathway, the experiments revealed several genes linked with sugar metabolism: two glucosyltransferases (GSVIVT00036656001 and GSVIVT00036670001), one Glc acyltransferase (GSVIVT00038626001), two sugar




Figure 6. Transcript levels of putative targets (A) and validation of microarray results (B) in grapevine organs overexpressing *VvMybPA1* (lines PA1-2A and PA1-7A) or *VvMybPA2* (lines PA2-1A and PA2-7A) versus the wild type (lines control-1 and control-2). Gene expression was determined by real-time PCR and normalized with the expression of EF1 α . All data are means \pm SD of three replicates. Stars indicate that expression levels are significantly different between wild-type and transformed hairy roots ($P < 0.01$).

Table 1. List of transcripts whose expression is induced after both VvMybPA1 and VvMybPA2 overexpression

Magnitudes of induction relative to wild-type controls are given. The *Vitis* transcript number refers to the genome model (Jaillon et al., 2007; <http://www.genoscope.cns.fr/externe/GenomeBrowser/Vitis/>) or to the DFCI tentative contig (<http://compbio.dfci.harvard.edu/tgi/cgi-bin/tgi/gimain.pl?gudb=grape>). The four right columns present best annotated matches after BLASTX analysis against the nr protein public database, species, putative protein functions, and match e values.

PA2/Wild-Type Ratio	PA1/Wild-Type Ratio	Probe Set Identifier	<i>Vitis</i> Transcript	Best Match	Species	Putative Function	E Value
3.6	2.21	Vv_10003904	GSVIVT00014419001	ABM67589	<i>Vitis vinifera</i>	Flavanone-3-hydroxylase 2	0.0
3.3	2.15	Vv_10003855	GSVIVT00036784001	P41090	<i>Vitis vinifera</i>	Flavanone-3-hydroxylase 1	0.0
2.82	1.68	Vv_10000003	GSVIVT00030258001	None			
2.8	3.74	Vv_10004449	GSVIVT00024561001	ACC63891	<i>Populus trichocarpa</i>	Phe ammonia lyase	0.0
2.8	2.51	Vv_10012881	TC64176	None			
2.76	1.98	Vv_10013426	GSVIVT00036670001	ABH03018	<i>Vitis labrusca</i>	Resveratrol/hydroxycinnamic acid <i>O</i> -glucosyltransferase	0.0
2.66	2.17	Vv_10000485	GSVIVT00014584001	P51110	<i>Vitis vinifera</i>	Dihydroflavonol-4-reductase	2e-157
2.63	2.81	Vv_10004167	GSVIVT00006341001	BAB84111	<i>Vitis vinifera</i>	Chalcone synthase 3	0.0
2.6	2.56	Vv_10009527	GSVIVT00014031001	P31687	<i>Glycine max</i>	4-Coumarate-CoA ligase 2	2e-173
2.37	2.04	Vv_10000352	GSVIVT00001063001	ABV82967	<i>Vitis vinifera</i>	Leucoanthocyanidin dioxygenase	3e-170
2.31	1.71	Vv_10000110	GSVIVT00012660001	BAE47626	<i>Lactuca sativa</i>	Cytochrome <i>b₆/f</i> complex subunit 4	3e-61
2.23	1.76	Vv_10004634	GSVIVT00036656001	ABH03018	<i>Vitis labrusca</i>	Resveratrol/hydroxycinnamic acid <i>O</i> -glucosyltransferase	0.0
2.15	1.47	Vv_10003799	GSVIVT00001815001	BAF03494	<i>Populus nigra</i>	XRCC4 homolog	9e-44
2.08	1.98	Vv_10004484	GSVIVT00015738001	AAL67601	<i>Oryza sativa</i>	Cinnamoyl-CoA reductase	8e-77
2.07	2.72	Vv_10000926	GSVIVT00038153001	NP_196974	<i>Arabidopsis thaliana</i>	Cinnamoyl-CoA reductase-related	3e-59
2.03	2.27	Vv_10003778	GSVIVT00005344001	CAD91911	<i>Vitis vinifera</i>	Anthocyanidin reductase	0.0
2.03	2.61	Vv_10002321	GSVIVT00018839001	NP_191462	<i>Arabidopsis thaliana</i>	TT12, MATE-type transporter	4e-169
2	2.26	Vv_10001009	GSVIVT00025894001	AAD29806	<i>Arabidopsis thaliana</i>	Disease resistance response protein	2e-16
1.93	1.89	Vv_10001107	GSVIVT00038626001	AAF64227	<i>Solanum pennellii</i>	Glc acyltransferase	2e-114
1.91	1.57	Vv_10013578	GSVIVT00025888001	NP_187420	<i>Arabidopsis thaliana</i>	Arogenate dehydratase/prephenate dehydratase	1e-116
1.89	1.92	Vv_10007208	GSVIVT00016217001	CAI54278	<i>Vitis vinifera</i>	Flavonoid-3'-hydroxylase	0.0
1.85	1.42	Vv_10003899	TC59772	None			
1.81	1.56	Vv_10001147	GSVIVT00015196001	ABA54865	<i>Fagus sylvatica</i>	3-Deoxy-D-arabino-heptulosonate 7-phosphate synthase	0.0
1.74	2.02	Vv_10011954	GSVIVT00023340001	ABK94803	<i>Populus trichocarpa</i>	Unknown protein	3e-112
1.74	2.21	Vv_10013105	GSVIVT00029573001	AAF02155	<i>Arabidopsis thaliana</i>	Unknown protein	4e-17
1.69	1.67	Vv_10008665	GSVIVT00008901001	ABA54867	<i>Fagus sylvatica</i>	3-Dehydroquininate dehydratase/shikimate 5-dehydrogenase	4e-159
1.68	2.02	Vv_10003316	GSVIVT00006090001	ABN09011	<i>Medicago truncatula</i>	Sugar transporter superfamily	2e-131
1.66	1.77	Vv_10004589	GSVIVT00012147001	NP_176734	<i>Arabidopsis thaliana</i>	Allyl alcohol dehydrogenase, putative	4e-143
1.58	1.59	Vv_10004860	GSVIVT00013718001	NP_565423	<i>Arabidopsis thaliana</i>	Unknown protein	7e-21
1.57	1.94	Vv_10010085	GSVIVT00005432001	ABA54868	<i>Fagus sylvatica</i>	Shikimate kinase	6e-69
1.56	1.51	Vv_10002822	GSVIVT00017259001	BAE71196	<i>Trifolium pratense</i>	Hydroxymethylglutaryl-CoA lyase	2e-176
1.48	1.52	Vv_10002309	GSVIVT00005204001	AAL36369	<i>Arabidopsis thaliana</i>	Receptor kinase	0.0
1.46	1.96	Vv_10002140	GSVIVT00015239001	AAT77693	<i>Vitis vinifera</i>	Hexose transporter HT2	0.0

(Table continues on following page.)

Table 1. (Continued from previous page.)

PA2/Wild-Type Ratio	PA1/Wild-Type Ratio	Probe Set Identifier	Vitis Transcript	Best Match	Species	Putative Function	E Value
1.46	1.42	Vv_10000937	GSVIVT00032373001	NP_567869	<i>Arabidopsis thaliana</i>	Electron carrier	2e-145
1.44	1.48	Vv_10000117	TC68399	CAL07988	<i>Platanus × acerifolia</i>	Translation elongation factor 1 α	9e-5
1.38	1.22	Vv_10002123	GSVIVT00005969001	AAB37756	<i>Solanum tuberosum</i>	Kinesin heavy chain-like protein	0.0
1.37	1.48	Vv_10004956	GSVIVT00017469001	NP_563818	<i>Arabidopsis thaliana</i>	Strictosidine synthase family protein	8e-146
1.37	1.46	Vv_10000360	GSVIVT00030469001	ABW34393	<i>Vitis vinifera</i>	R2R3 Myb transcription factor C2 repressor motif protein	4e-130
1.36	1.42	Vv_10009241	GSVIVT00024556001	NP_565668	<i>Arabidopsis thaliana</i>	Unknown protein	4e-95
1.31	1.52	Vv_10008052	GSVIVT00020463001	NP_193106	<i>Arabidopsis thaliana</i>	Purple acid phosphatase	0.0
1.31	1.3	Vv_10009804	GSVIVT00024607001	ABN08215	<i>Medicago truncatula</i>	Zinc finger (CCCH-type) family protein	0.0
1.31	1.32	Vv_10000047	GSVIVT00033143001	NP_199757	<i>Arabidopsis thaliana</i>	ATP-citrate lyase B-2	0.0
1.28	1.15	Vv_10004398	GSVIVT00034002001	AAK55470	<i>Oryza sativa</i>	Phosphoinositide kinase	3e-150
1.26	1.2	Vv_10004784	GSVIVT00014475001	ABC68397	<i>Glycine max</i>	Cytochrome P450 monooxygenase CYP83E8	6e-156
1.26	1.32	Vv_10013455	GSVIVT00026033001	AAR88099	<i>Arabidopsis thaliana</i>	Glutamate receptor ion channel	0.0
1.26	1.24	Vv_10000645	GSVIVT00031453001	NP_568860	<i>Arabidopsis thaliana</i>	CBL-interacting protein kinase	1e-144
1.25	1.42	Vv_10004922	GSVIVT00031779001	ABW38332	<i>Fragaria × ananassa</i>	Cytosolic Fru-1,6-bisphosphatase	1e-176
1.24	1.24	Vv_10010519	GSVIVT00029310001	AAF29605	<i>Pisum sativum</i>	Gibberellin c20-oxidase	6e-06
1.20	1.08	Vv_10004613	GSVIVT00001461001	NP_179081	<i>Arabidopsis thaliana</i>	Vacuolar sorting receptor	0.0
1.16	1.23	Vv_10009243	GSVIVT00018590001	NP_179660	<i>Arabidopsis thaliana</i>	Glycosyl hydrolase family 5 protein	1e-180
1.14	1.14	Vv_10010645	GSVIVT00034434001	NP_001117624	<i>Arabidopsis thaliana</i>	Coatmer protein complex, subunit β 2	0.0

transporters, and a Fru-1,6-bisphosphatase. Another striking category contains seven genes related to signaling mechanisms, including a Myb factor, a calcineurin B-like (CBL) protein, and a CBL-interacting protein kinase. All of these genes represent the most induced genes, and for some of them this induction was confirmed by real-time PCR (Fig. 6B). Genes induced to a lower extent have miscellaneous or unknown functions, and some of them do not even exhibit homologies with other genes of the plant kingdom.

A total of 828 genes were found significantly differentially expressed between hairy roots overexpressing *VvMybPA1* and *VvMybPA2*. Among them, 59 are specifically induced in 35S::*VvMybPA1* transformants, such as a chalcone synthase isogene, another MATE transporter, while 21 are more specific for *VvMybPA2*, such as another glucosyltransferase, GSVIVT00013771001 (Fig. 6B; Supplemental Materials and Methods S1).

DISCUSSION

VvMybPA1 and *VvMybPA2* Regulate the PA Pathway in Grapevine Berries

PA synthesis in grapevine berries is restricted to the early stages of development in skin and seed and in pulp to a certain extent (Verriès et al., 2008). Moreover, PAs of these different tissues exhibit noticeable structural differences whose biological determinism is unknown, although it is probably linked to the differential regulation of F3'H, flavonoid 3',5'-hydroxylase (F3'5'H), ANR, LAR1, and LAR2 isogenes in a precise time- and tissue-specific manner. The data presented here indicate that at least two transcription factors belonging to the Myb family are involved in the control of PA content in grapevine.

VvMybPA1 was previously identified by Bogs et al. (2007). *VvMybPA2*, which was identified here, appears to be the closest ortholog of AtTT2 identified so

far in grapevine. R2R3 Myb factors share very limited homology in their C-terminal regions, which could be related to their target specificity. A sequence conserved between AtTT2 and OsMYB3 and more recently confirmed in *Brassica napus* putative orthologs of TT2 was identified in this highly variable C-terminal region (Nesi et al., 2001; Stracke et al., 2001; Wei et al., 2007). This motif is also present in VvMybPA2 but is not strictly conserved in the VvMybPA1 sequence (Bogs et al., 2007).

We demonstrated here that the spatiotemporal expression of *VvMybPA2*, restricted to the skin of young berries and leaves, is compatible with a putative role in the regulation of PA biosynthesis. The *VvMybPA2* expression pattern argues in favor of a preferential role in PA synthesis and accumulation in berry skins when compared with *VvMybPA1*. *VvMybPA1* is expressed before *véraison* and actually correlates with PA accumulation in seeds (Bogs et al., 2007). Its involvement in the accumulation of PA accumulation in skin was ruled out, on the basis of its very low expression level in this tissue. Moreover, according to some authors, *VvMybPA1* showed an unexpected maximum of expression 2 weeks after *véraison* when PA accumulation was stopped (Bogs et al., 2007; Deluc et al., 2007). Conversely, the profile of *VvMybPA2* is consistent with a critical function in the accumulation of PAs in berry skin. It was hypothesized that the different branches of the flavonoid and phenylpropanoid pathways are specified by Myb proteins inside the activating complex (Zhang et al., 2003). However, Deluc et al. (2006), expressing *VvMyb5a* transcription factor in tobacco (*Nicotiana tabacum*), showed that the activation of diverse branches of the flavonoid and phenylpropanoid pathways in the different parts of the transgenic plants could also be modulated by bHLH and WD40 as members of the transcriptional machinery. Nevertheless, this transcription factor did not trigger anthocyanin accumulation in natural circumstances, being expressed in berries at green stage, and its actual function in grapevine remains to be clarified. We show here that both VvMybPA1 and the new VvMybPA2 actually act as specific positive regulators of PA biosynthesis when stably expressed in grapevine, alleviating the hazard of artifacts upon ectopic expression in heterologous plants, as suspected for VvMyb5a (Deluc et al., 2006). Both transcription factors described here appeared as extremely specific to the PA pathway, since no shift toward lignin, anthocyanin, and flavonol production appeared in long-term experiments.

***VvMybPA1* and *VvMybPA2* Overexpression Give Similar Results**

In order to clarify the specific roles of VvMybPA1 and VvMybPA2, we observed the respective phenotypic and transcriptomic variations induced by their ectopic overexpression in grapevine. Both genes were able to modify PA contents and their subunit composition. When overexpressed in hairy roots, *VvMybPA1*

and *VvMybPA2* actually triggered de novo B ring trihydroxylation, and a slightly increased mDP was observed in the case of *VvMybPA2*-expressing lines, which is consistent with the composition of skin PAs. The level of trihydroxylation of transgenic hairy root PAs was similar to that of Maccabeo berry skin (3.5%) and higher than in seeds (0%; Souquet et al., 2006). In this respect, the increase of trihydroxylated units observed here upon the overexpression of *VvMybPA1* is consistent with its ability to activate the promoter of F3'5'H (Bogs et al., 2007). However grapevine seed PAs do not contain any (epi)gallo catechin unit, which may suggest a pretranscriptional or posttranscriptional negative regulation of F3'5'H in the seeds. Despite exhibiting an increase upon ectopic expression of *VvMybPA2*, the PA chain length remained three times lower than the mDP of approximately 50 encountered inside the skin of Maccabeo plants from the field (Souquet et al., 2006). More generally, we did not succeed in reproducing the large structural differences among seed and skin tannins upon differential ectopic expression of *VvMybPA1* and *VvMybPA2*.

This PA accumulation was found restricted to some tissues despite the use of a 35S promoter. At the transcription level, several causes could explain the deficit in PA accumulation in some tissues and cells: (1) lack of members of the activating complex, (2) absence of transcription factors in the role of activators, or (3) expression of inhibitors. In Arabidopsis, Nesi et al. (2001), observing that ectopic expression of TT2 activated TT8, BAN, or TT12 without subsequent PA accumulation, hypothesized the lack of activation of earlier genes of the pathway. In grapevine cell suspensions, cotransformation with a bHLH factor was found mandatory for the transient activation of the *LDOX* promoter by VvMybPA1 (Bogs et al., 2007). In our experiments, the sole overexpression of *VvMybPA1* or *VvMybPA2* proved to be sufficient to enhance the *LDOX* expression. This means that either the partners forming the PA regulatory complex were already present in the tissues of the roots accumulating PAs or that long-term overexpression is necessary to induce all partners required for full activation of the PA pathway. The latter hypothesis appears rather unlikely, since no induction of any bHLH or WD40 transcription factors was found associated with the ectopic accumulation of PAs in grapevine. However, it must be kept in mind that the 14 K oligoarray used in this study may probe only half of the *Vitis* transcriptome, according to Jaillon et al. (2007).

As the phenotypes of transgenic grapevine organs expressing *VvMybPA1* or *VvMybPA2* are quite similar, the reason for the existence of two distinct transcription factors with apparent redundant function can be questioned. A similar functional redundancy was described in Arabidopsis, in which three different Myb factors controlling the flavonol pathway were identified, each of them exhibiting a specific tissue expression pattern (Stracke et al., 2007). Recently, Matus et al. (2008) classified the 108 members of the grape R2R3

Myb family in terms of their genomic structures and similarity to their putative Arabidopsis orthologues. Eight genes, including *VvMybPA2* but neither *VvMybPA1* nor *VvMyb5a*, were designated as putative candidates for PA pathway regulators. Further work is needed to better understand the putative functions of these regulators in the tissue-specific regulation of the PA pathway.

Despite the phenotype similarities, the high-throughput transcriptomic analysis presented here revealed that several targets were specifically induced by each Myb factor. However, either the genes identified were isogenes of already identified targets, like the identified glucosyltransferase specifically induced in *VvMybPA2* transgenic lines, suggesting here again functional redundancy (or very subtle differences in their catalytic properties), or the level of induction for these genes was remarkably low.

This experiment also raises the question of the interactions between transcription factors. A first result emerged from this study, where overexpression of *VvMybPA2* resulted in the accumulation of *VvMybPA1* transcripts, suggesting that *VvMybPA2* signal acts upstream to *VvMybPA1*.

Expected and Unexpected Targets, and Absence of Expected Targets

Enzymes of the Flavonoid Pathway

Being the final result of a transcriptomic screening after overexpression of two different transcription factors inducing PA accumulation, the list of the 51 induced transcripts is very likely to contain specific isogenes involved in the PA pathway. Microarray analysis revealed the induction of genes from Phe ammonia lyase to *ANR*. Rather low expression ratios were observed after overexpression of both Myb factors. This could result from the preactivation of the PA pathway in wild-type hairy roots, which already contain significant amounts of PAs. In addition, we observed that despite the use of a constitutive promoter, only specific tissues were able to ectopically accumulate PAs. Consequently, the sensibility of the analysis is probably hampered by the low number of cells inside the organs affected by PA accumulation. Some genes belonging to this pathway were not detected. For example, the only copy of chalcone isomerase in the genome is induced in both types of transformants, but its induction reaches a significant threshold only under *VvMybPA1* overexpression. In addition, some genes could not be monitored due to the absence of a probe, like *F3'5'H*. The induction of *F3'5'H* seems necessary to synthesize the trihydroxylated units observed in transgenic roots. Unfortunately, the grapevine genome contains 10 genes coding for putative *F3'5'H* (Jaillon et al., 2007; Velasco et al., 2007), which precluded the design of isogene-specific 70-mer-long probes.

Neither the overexpression of *VvMybPA1* nor that of *VvMybPA2* resulted in a significant induction of *LAR2*.

Bogs et al. (2007) reported that *VvMybPA1* was able to activate the promoters of *LAR1*, as confirmed here, but they did not address *LAR2* activation. The proportion of catechin as TU or free monomer was not impaired when compared with the wild type, probably due to the activity of *LAR1* (data not shown). Disturbing results concerning particular *LAR* isogenes are reported in the literature on other plants (Pang et al., 2007; Paolucci et al., 2007). Similarly, ectopic expression of the maize (*Zea mays*) *bHLH* flavonoid regulator *Sn* in *Lotus corniculatus* resulted in an increase of PA content and induction of *ANR* and *LAR1* transcripts, while *LAR2* transcription remained unaffected (Paolucci et al., 2007). They failed to detect any catechin synthesis after the heterologous production of the *LAR2* enzyme (Paolucci et al., 2007). PAs of *Medicago truncatula* seed coat contain insignificant amounts of catechin as TU or EU, despite the presence of *LAR* transcript in this tissue and the ability of the heterologous *LAR* protein to synthesize catechin in vitro (Pang et al., 2007). These authors also reported that transgenic tobacco plants overexpressing *MtLAR* did not exhibit changes in their PA or catechin content. Taken together, these results indicate that *LAR* isogenes occupy a particular place in the PA pathway, from the functional and transcriptional regulation points of view, and the precise role of *LAR2* remains to be elucidated.

In addition to the induction of genes previously shown to be involved in *sensu stricto* PA biosynthesis, screenings revealed the induction of one (GSVIVT00018839001) of the 65 MATE-type transporters of the grapevine genome (C. Gomez, personal communication). With 70% amino acid homology, this gene appears to be the closest homolog of *TT12*, which was shown to be critical for PA accumulation in the Arabidopsis seed testa (Debeaujon et al., 2001). The protein encoded by *TT12* was found located on the tonoplast, and in vitro experiments showed that *TT12* transports the anthocyanin cyanidin-3-O-glucoside (Marinova et al., 2007). A glutathione S-transferase (Kitamura et al., 2004) and a AHA10-like H^+ -ATPase (Baxter et al., 2005) have been reported to be involved in PA storage, but no grapevine homolog of these transcripts has been identified in our microarray screening.

Several genes were described as systematically expressed concomitantly with anthocyanin accumulation, like UDP-Glc:flavonoid 3-O-glucosyltransferase (Boss et al., 1996) or a putative methyl transferase and a particular isogene of glutathione S-transferase (Ageorges et al., 2006). Other experiments were performed by our group with the aim to ectopically activate anthocyanin accumulation in hairy roots with the *VIMybA1* gene identified by Kobayashi et al. (2002). It resulted in anthocyanin accumulation and the induction of a set of genes specific for the anthocyanin pathway, without any modification of their PA content or of the expression level of the PA biosynthetic genes, *ANR*, *LAR1*, and *LAR2* (Cutanda-Perez

et al., 2009). On the other hand, neither the anthocyanin biosynthetic genes nor flavonol synthase were induced in the hairy roots overexpressing *VvMybPA1* and *VvMybPA2* genes, confirming their specific role in the regulation of the PA pathway and an activation spectrum different from that of *VIMybA1*.

Signal Transduction

Overexpression of both *VvMybPA1* and *VvMybPA2* also induced the expression of a set of genes associated with signaling. A homolog of Arabidopsis Myb4 was ectopically induced in grapevine organs. AtMyb4 was shown to act as a repressor of the phenylpropanoid pathway (Jin et al., 2000). Its induction as a consequence of *VvMybPA1* and *VvMybPA2* overexpression could be considered as a way to counterbalance their attempt to avoid the bolting of the PA pathway.

CBL proteins and their target kinases, CBL-interacting protein kinases, have often been described as functioning in complex in the response to abiotic stresses (Baticic and Kudla, 2004). The simultaneous induction of these genes may result from the stress experienced by the plant after ectopic expression of the *VvMybPA* or from a sudden increase of intracellular PA concentration.

Genes Linked to Sugar Metabolism

Several genes related to sugar metabolism appeared as potentially driven by *VvMybPA* factors. Among them, the induction of a hexose transporter called *VvHT2* (Fillion et al., 1999) was unexpected. Fillion et al. (1999) described its expression maximum 1 week after *véraison*, whereas Hayes et al. (2007) reported a constant decrease of expression during berry development. However, the in planta role of *VvHT2* remains to be characterized. Incidentally, a protein from *Lotus japonica* annotated as “Suc transporter” and located at the tonoplast proved to be capable of transporting phenyl glucosides (Reinders et al., 2008). A possible involvement of genes annotated as “sugar transporter” in the transport of a glucosylated intermediate of the pathway cannot be excluded.

The two closest identified homologs of GSVIVT00038626001, a putative Glc acyltransferase, are (1) *DkSCPL1*, a gene of persimmon (*Diospyros kaki*) identified through a suppression subtractive hybridization between fruits differing in their PA content (Ikegami et al., 2007), and (2) a *Solanum pennellii* acyltransferase catalyzing the formation of diacylglycerol (Li and Steffens, 2000). These proteins could act as Glc acyltransferases that use 1-O- β -acetyl Glc esters as acyl donors (Milkowski and Strack, 2004).

Several glucosyltransferases were also identified. Two of them (GSVIVT00036670001 and GSVIVT00036656001) are induced by both transcription factors, whereas another one (GTGSVIVT00013771001) appears more specifically driven by *VvMybPA2*. GSVIVT00036656001 was already identified as a

bifunctional resveratrol/hydroxycinnamic acid glucosyltransferase in *Vitis labrusca* (Hall and DeLuca, 2007). This protein forms Glc esters with phenolic acids at low pH (5–7) and O-glucosides of resveratrol or flavonols at higher pH (7–9) at a much lower rate. Here, only trace amounts of transresveratrol and of the O-glucosides of cis- and trans-resveratrol were detected without any difference between wild-type and *VvMybPA*-overexpressing organs (data not shown). Recently, a transcriptomic screening after ectopic expression of *TT2* in *M. truncatula* hairy roots identified a glucosyltransferase able to catalyze the formation of epicatechin 3'-glucoside (Pang et al., 2008). However, this protein is quite divergent from the glucosyltransferases identified in this work, with only 26% to 28% identity. The occurrence of glycosylated PAs was described in other species (Xie and Dixon, 2005), but they were never encountered in grapevine or in the hairy roots described here. The absence of glucosylated PA in *VvMybPA*-overexpressing hairy roots makes it unlikely that these GT isogenes are involved in PA glucosylation. The actual biological functions of Glc acyltransferase and/or glucosyltransferases in grapevine PA biosynthesis remain to be determined and should be further investigated.

Original PA composition with respect to model plants, recent developments of new genomic tools, and the availability of reverse genetics technologies allow grapevine to be considered as a valuable model for the study of PA biosynthesis. We demonstrated here that the identification of a new specific transcription factor associated with homologous transformation allowed the determination of a specific network of genes associated with the activation of PA accumulation. This network was shown to involve several already annotated genes and revealed new putative actors.

MATERIALS AND METHODS

Plant Material and Nucleic Acid Extraction

Organs (berries, leaves, tendrils, and roots) from grapevine (*Vitis vinifera* ‘Shiraz’) plants grown in the SupAgro-INRA vineyard in Montpellier, France, were collected at several developmental stages. Young leaves corresponded to leaves explanted from the third node below the shoot tips, with a mean weight of 0.3 g. Old leaves corresponded to fully expanded leaves sampled from the mature shoot part, with a mean weight of 2.8 g. Eight- to 10-week-old plantlets of cv Maccabeu grapevine propagated onto half-strength Murashige and Skoog medium were used for *Agrobacterium tumefaciens* transformation procedures.

After sampling, hairy roots and plant organs were rapidly frozen in liquid nitrogen and then ground to a fine powder with a Dangoumau blender (Dangoumill 300) and stored at -80°C until use.

DNA was extracted from 50 mg of frozen tissue using the DNA Plant Mini kit (Qiagen). Total RNA was extracted using the RNeasy Plant Mini kit (Qiagen) following the manufacturer’s instructions, starting from 200 mg of tissue.

Cloning and Vectors

The coding regions of *VvMybPA1* and *VvMybPA2* were amplified from young Shiraz pericarp cDNA with high-fidelity Taq polymerase (Advantage-

HF 2 PCR kit; Clontech) using the forward primers 5'-CACCATGGGCA-GAGCACCTTGTGT-3' and 5'-CACCATGGGAAGAAGACCTTGCTG-3' and the reverse primers 5'-TTAAATGAGTAGTGATTCGGC-3' and 5'-CTA-TGGGACTTGATTATTTTC-3' for VvMybPA1 and VvMybPA2, respectively. The amplicons were directionally cloned in pENTR/D-TOPO (Invitrogen Life Technologies) according to the manufacturer's instructions. The sequences of the positive clones were confirmed following transformation of One Shot competent *Escherichia coli* (Invitrogen Life Technologies) and LR recombination in the binary vector pH2GW7 (Karimi et al., 2002) to yield the 35S VvMybPAs constructs. The sequence of *VvMybPA2* was deposited in GenBank under accession number EU919682.

Recombinant plasmids were electroporated into *Agrobacterium rhizogenes* strain A4 introduced from Collection Française de Bactéries Phytopathogènes (<http://www-intranet.angers.inra.fr/cfbp/>).

Sequence Analysis

Full-length amino acid sequences of Myb factors from several species were retrieved from public databases. Alignments were performed with the ClustalW2 algorithm with default parameters (Thompson et al., 1994). The phylogenetic tree was constructed from the Clustal alignment using the neighbor-joining method in the MEGA4 package (Kumar et al., 2004).

Real-Time PCR

The RNA was accurately quantified with Ribogreen reagent (Molecular Probes). A triplicate reverse transcription was performed on 500 ng of total RNA from each developmental stage using the SuperScript II reverse transcription kit (Invitrogen) according to the manufacturer's instructions. Triplicate reverse transcriptions for PCR were pooled to minimize the heterogeneity of the reverse transcription reaction efficiency. Specific oligonucleotide primer pairs were designed with Primer3 software except for LDOX (Bogs et al., 2005) and for dihydroflavonol-4-reductase (Jeong et al., 2004) and are listed in Supplemental Table S2. Specific annealing of the oligonucleotides was controlled on dissociation kinetics performed at the end of each PCR run. The efficiency of each primer pair was measured on a PCR product or plasmid serial dilutions. PCR was performed in triplicate on 1 μ L of cDNA from each sample using the model 7300 Sequence Detection System (Applied Biosystems) and the Power SYBR-Green PCR Master kit (Applied Biosystems Applied France). EF1 α was used as a reference for normalization of gene expression (Terrier et al., 2005; Reid et al., 2006). The difference between the cycle threshold (Ct) of the target gene and EF1 α was used to obtain the normalized expression of the target gene, calculated as $2^{\text{exp} - (\text{Ct}_{\text{target}} - \text{Ct}_{\text{EF1}\alpha})}$.

Genetic Transformation Procedures

Induction and culture of transgenic hairy roots in grapevine were performed as described by Torregrosa and Bouquet (1997) with the following modifications: (1) *A. rhizogenes* strain A4 was used as vector; (2) bacterial cultures were grown on semisolid MGL/B (5 g L⁻¹ tryptone, 5 g L⁻¹ mannitol, 2.5 g L⁻¹ yeast extract, 1 g L⁻¹ Glu, 250 mg L⁻¹ K₂HPO₄, 100 mg L⁻¹ NaCl, 100 mg L⁻¹ MgSO₄·7H₂O, and 5 μ g L⁻¹ biotin at pH 7) medium (Torregrosa et al., 2002) supplemented with 50 μ g mL⁻¹ spectinomycin and then suspended with half-strength Murashige and Skoog liquid medium (optical density at 600 nm = 0.3) in the presence of 100 μ M acetosyringone for inoculation; and (3) roots tips extracted from explants were cultured on modified LG0 medium (LG0 as described by Torregrosa and Bouquet [1997] added with 250 mg L⁻¹ casein hydrolysate and solidified 5 g L⁻¹ Phytigel [Sigma] with 200 μ g mL⁻¹ both augmentin and cefotaxime [Duchefa] to prevent bacterial growth). Stable hairy root lines were subcultured every 4 to 5 weeks on the same medium until use.

Phenotyping the Transformed Hairy Roots

Extraction and HPLC Methods

PA extractions were performed as described by Verriès et al. (2008). During 1 h, 100 mg of frozen and powdered sample was mixed with 750 μ L of the extraction solution (acetone:water [70:30, v/v] containing 0.05% trifluoroacetic acid) and 50 μ L of an internal standard solution (*p*-hydroxy methyl ester, 3 g L⁻¹ in methanol). *p*-Hydroxy methyl ester was chosen because of its

stability during extraction and phloroglucinolysis reactions, its retention time being quite different from those of PA units in our HPLC conditions, and its absorbance spectrum. Samples were centrifuged (13,000 g, 15 min, 4°C), and the supernatants were recovered. Immediately, 200 μ L of each supernatant was dried under vacuum at 35°C for 2 h (Genevac) and resuspended in 100 μ L of reagent solution (0.25 g of phloroglucinol, 0.05 g of ascorbic acid, and 5 mL of acidified methanol [0.2 N HCl]) for acid-catalyzed degradation in the presence of excess phloroglucinol. After incubation (50°C, 20 min), the reaction was stopped by adding 100 μ L of sodium acetate buffer (200 mM, pH 7.5). Samples were then centrifuged before injection into the HPLC system. HPLC analyses were performed as described by Verriès et al. (2008). Results were expressed in milligrams per gram fresh weight. Polymer length was estimated by the mDP, calculated as the ratio between the sum of EU (phloroglucinol adducts) and TU and the sum of TU, including free monomers. The mDP, the percentage of galloylation, and the percentage of epigallocatechin units (or trihydroxylation) were calculated on a molar basis.

The lignin content and composition of the hairy root samples were evaluated by thioacidolysis, as described previously, on 5 to 10 mg of freeze-dried sample (Lapierre et al., 1995; Nakashima et al., 2008).

Coloration with DMACA

DMACA reacts quite specifically with flavan 3-ol monomers and PAs to form a blue chromophore (Feucht et al., 2004). The presence and localization of PAs in hairy roots were detected by staining the tissues with the DMACA solution (1% DMACA and 1% 6 N HCl in methanol) for 15 min. The roots were then rinsed with distilled water, and blue staining was visualized with a microscope and captured using a digital camera. Figures were formatted and assembled with Adobe Photoshop 7.0.

Microarray Experiments

We used the Qiagen Operon Array-Ready Oligo Set for the Grape Genome Version 1.0 containing 14,562 70-mer probes representing 14,562 transcripts from The Institute for Genomic Research (TIGR) Grape Gene Index, release 3. Oligonucleotides were reannotated using the DFCI Grape Gene Index, release 5.0 (June 21, 2006; <http://compbio.dfci.harvard.edu/tgi/cgi-bin/tgi/gimain.pl?gudb=grape>). Oligonucleotides were spotted on mirror slides, and the probes were labeled with Cy3 and Cy5 dyes with the Amino Allyl MessageAmp II aRNA kit (Ambion). The experiment was performed using eight slides (two biological replicates for each condition and a dye swap, detailed in Supplemental Materials and Methods S1). Hybridized microarrays were simultaneously scanned for Cy3- and Cy5-labeled probes with an Axon Genepix 4000B scanner.

Statistical Treatments

Regarding microarray experiments, data from both channels corresponding to Cy3- and Cy5-labeled probes were normalized using the Lowess algorithm in the Microarray Data Analysis System at TIGR. Data from all of the slides were log₂ transformed and normalized (centered on 0, variance equalized to 1); those data are available in Supplemental Materials and Methods S1. The significance was calculated at the 0.01 level by permutation *t* test in Multiexperiment Viewer from TIGR.

The significance (*P* < 0.01) of the results from biochemical analysis and real-time PCR was statistically assessed with a permutation *t* test in the Past software.

Sequence data from this article can be found in the GenBank/EMBL data libraries under accession number EU919682.

Supplemental Data

The following materials are available in the online version of this article.

Supplemental Table S1. Lignin content, subunit composition, and proportion (guaiacyl and syringyl lignin-derived monomers) in grapevine organs overexpressing *VvMybPA1* or *VvMybPA2* compared with controls.

Supplemental Table S2. Primers used for real-time PCR and expected sizes of the amplified fragments.

Supplemental Materials and Methods S1. terriersupS2.xls contains experimental design ("design"; sheet 1), complete hybridization results \log_2 transformed and normalized ("totlognorm"; sheet 2), list of genes for which transcript accumulation is significantly affected by ectopic overexpression of *VvMybPA1* ("PA1signif"; sheet 3) or *VvMybPA2* ("PA2signif"; sheet 4), list of genes for which transcript accumulation is significantly different between ectopic overexpression of *VvMybPA1* and *VvMybPA2* ("PA2diffPA1"; sheet 5), list of genes specifically induced by ectopic overexpression of *VvMybPA1* ("PA1spec"; sheet 6), and list of genes specifically induced by ectopic overexpression of *VvMybPA2* ("PA2spec"; sheet 6).

ACKNOWLEDGMENTS

The assistance of Prof. Catherine Lapierre and of Mrs. Brigitte Pollet (UMR 206 Chimie Biologique, AgroParisTech-INRA, Agroparistech Centre de Grignon) is sincerely acknowledged for the thioacidolysis experiments and their interpretation.

Received October 29, 2008; accepted December 16, 2008; published December 19, 2008.

LITERATURE CITED

- Ageorges A, Fernandez L, Violet S, Merdinoglu D, Terrier N, Romieu C (2006) Four specific isogenes of the anthocyanin metabolic pathway are systematically co-expressed with the red colour of grape berries. *Plant Sci* **170**: 372–383
- Batistic O, Kudla J (2004) Integration and channeling of calcium signaling through the CBL calcium sensor/CIPK protein kinase network. *Planta* **219**: 915–924
- Baxter IR, Young JC, Armstrong G, Foster N, Bogenschutz N, Cordova T, Peer WA, Hazen SP, Murphy AS, Harper JF (2005) A plasma membrane H^+ -ATPase is required for the formation of proanthocyanidins in the seed coat endothelium of *Arabidopsis thaliana*. *Proc Natl Acad Sci USA* **102**: 2649–2654
- Bogs J, Downey M, Harvey J, Ashton A, Tanner G, Robinson S (2005) Proanthocyanidin synthesis and expression of genes encoding leucoanthocyanidin reductase and anthocyanidin reductase in developing grape berries and grapevine leaves. *Plant Physiol* **139**: 652–663
- Bogs J, Jaffe FW, Takos AM, Walker AR, Robinson SP (2007) The grapevine transcription factor VvMYBPA1 regulates proanthocyanidin synthesis during fruit development. *Plant Physiol* **143**: 1347–1361
- Borevitz J, Xia Y, Blount J, Dixon R, Lamb C (2000) Activation tagging identifies a conserved MYB regulator of phenylpropanoid biosynthesis. *Plant Cell* **12**: 2383–2393
- Boss PK, Davies C, Robinson SP (1996) Analysis of the expression of anthocyanin pathway genes in developing *Vitis vinifera* L. cv Shiraz grape berries and the implications for pathway regulation. *Plant Physiol* **111**: 1059–1066
- Cutanda-Perez M, Ageorges A, Gomez C, Violet S, Terrier N, Romieu C, Torregrosa L (2009) Ectopic expression of VmybA1 in grapevine activates a narrow set of genes involved in anthocyanin synthesis and transport. *Plant Mol Biol* (in press)
- Debeaujon I, Peeters AJ, Léon-Kloosterziel KM, Koornneef M (2001) The TRANSPARENT TESTA12 gene of *Arabidopsis* encodes a multidrug secondary transporter-like protein required for flavonoid sequestration in vacuoles of the seed coat endothelium. *Plant Cell* **13**: 853–871
- Deluc L, Barrieu F, Marchive C, Lauvergeat V, Decendit A, Richard T, Carde JP, Merillon JM, Hamdi S (2006) Characterization of a grapevine R2R3-MYB transcription factor that regulates the phenylpropanoid pathway. *Plant Physiol* **140**: 499–511
- Deluc LG, Grimplet J, Wheatley MD, Tillett RL, Quilici DR, Osborne C, Schooley DA, Schlauch KA, Cushman JC, Cramer GR (2007) Transcriptomic and metabolite analyses of Cabernet Sauvignon grape berry development. *BMC Genomics* **8**: 429
- Devic M, Guillemint J, Debeaujon I, Bechtold N, Bensaude E, Koornneef M, Pelletier G, Delseny M (1999) The BANYULS gene encodes a DFR-like protein and is a marker of early seed coat development. *Plant J* **19**: 387–398
- Dixon RA, Xie DY, Sharma SB (2005) Proanthocyanidins: a final frontier in flavonoid research? *New Phytol* **165**: 9–28
- Espley R, Hellens R, Putterill J, Stevenson D, Kuty-Amma S, Allan A (2007) Red coloration in apple fruit is due to the activity of the MYB transcription factor, MdMYB10. *Plant J* **49**: 414–427
- Feucht W, Treutter D, Polster J (2004) Flavanol binding of nuclei from tree species. *Plant Cell Rep* **22**: 430–436
- Fillion L, Ageorges A, Picaud S, Coutos-Thévenot P, Lemoine R, Romieu C, Delrot S (1999) Cloning and expression of a hexose transporter gene expressed during the ripening of grape berry. *Plant Physiol* **120**: 1083–1094
- Grotewold E (2006) The genetics and biochemistry of floral pigments. *Annu Rev Plant Biol* **57**: 761–780
- Grotewold E, Sainz MB, Tagliani L, Hernandez JM, Bowen B, Chandler VL (2000) Identification of the residues in the Myb domain of maize C1 that specify the interaction with the bHLH cofactor R. *Proc Natl Acad Sci USA* **97**: 13579–13584
- Hall D, De Luca V (2007) Mesocarp localization of a bi-functional resveratrol/hydroxycinnamic acid glucosyltransferase of Concord grape (*Vitis labrusca*). *Plant J* **49**: 579–591
- Hayes MA, Davies C, Dry IB (2007) Isolation, functional characterization, and expression analysis of grapevine (*Vitis vinifera* L.) hexose transporters: differential roles in sink and source tissues. *J Exp Bot* **58**: 1985–1997
- Ikegami A, Eguchi S, Kitajima A, Inoue K, Yonemori K (2007) Identification of genes involved in proanthocyanidin biosynthesis of persimmon (*Diospyros kaki*) fruit. *Plant Sci* **172**: 1037–1047
- Jaillon O, Aury JM, Noel B, Policriti A, Clepet C, Casagrande A, Choisne N, Aubourg S, Vitulo N, Jubin C, et al (2007) The grapevine genome sequence suggests ancestral hexaploidization in major angiosperm phyla. *Nature* **449**: 463–467
- Jeong ST, Goto-Yamamoto N, Kobayashi S, Esaka M (2004) Effects of plant hormones and shading on the accumulation of anthocyanins and the expression of anthocyanin biosynthetic genes in grape berry skins. *Plant Sci* **167**: 247–252
- Jin H, Cominelli E, Bailey P, Parr A, Mehrtens F, Jones J, Tonelli C, Weisshaar B, Martin C (2000) Transcriptional repression by AtMYB4 controls production of UV-protecting sunscreens in *Arabidopsis*. *EMBO J* **19**: 6150–6161
- Karimi M, Inze D, Depicker A (2002) Gateway vectors for Agrobacterium-mediated plant transformation. *Trends Plant Sci* **7**: 193–195
- Kennedy JA, Hayasaka Y, Vidal S, Waters EJ, Jones GP (2001) Composition of grape skin proanthocyanidins at different stages of berry development. *J Agric Food Chem* **49**: 5348–5355
- Kitamura S, Shikazono N, Tanaka A (2004) TRANSPARENT TESTA 19 is involved in the accumulation of both anthocyanins and proanthocyanidins in *Arabidopsis*. *Plant J* **37**: 104–114
- Kobayashi S, Goto-Yamamoto N, Hirochika H (2004) Retrotransposon-induced mutations in grape skin color. *Science* **304**: 982
- Kobayashi S, Ishimaru M, Hiraoka K, Honda C (2002) Myb related genes of the Kyoho grape (*Vitis labruscana*) regulate anthocyanin biosynthesis. *Planta* **215**: 924–933
- Kumar S, Tamura K, Nei M (2004) MEGA3: integrated software for molecular evolutionary genetics analysis and sequence alignment. *Brief Bioinform* **5**: 150–163
- Lapierre C, Pollet B, Rolando R (1995) New insights into the molecular architecture of hardwood lignins by chemical degradation methods. *Res Chem Intermediat* **21**: 397–412
- Lepiniec L, Debeaujon I, Routaboul JM, Baudry A, Pourcel L, Nesi N, Caboche M (2006) Genetics and biochemistry of seed flavonoids. *Annu Rev Plant Biol* **57**: 405–430
- Li AX, Steffens JC (2000) An acyltransferase catalyzing the formation of diacylglycerol is a serine carboxypeptidase-like protein. *Proc Natl Acad Sci USA* **97**: 6902–6907
- Mané C, Souquet JM, Olle D, Verriès C, Veran F, Mazerolles G, Cheyrier V, Fulcrand H (2007) Optimization of simultaneous flavanol, phenolic acid, and anthocyanin extraction from grapes using an experimental design: application to the characterization of champagne grape varieties. *J Agric Food Chem* **55**: 7224–7233
- Marinova K, Pourcel L, Weder B, Schwarz M, Barron D, Routaboul JM, Debeaujon I, Klein M (2007) The *Arabidopsis* MATE transporter TT12 acts as a vacuolar flavonoid/ H^+ -antiporter active in proanthocyanidin-accumulating cells of the seed coat. *Plant Cell* **19**: 2023–2038
- Mathews H, Clendennen S, Caldwell C, Liu XL, Connors K, Matheis N, Schuster D, Menasco D, Wagoner W, Lightner J, Ry Wagner D (2003)

- Activation tagging in tomato identifies a transcriptional regulator of anthocyanin biosynthesis, modification, and transport. *Plant Cell* **15**: 1689–1703
- Matus JT, Aquea F, Arce-Johnson P** (2008) Analysis of the grape MYB R2R3 subfamily reveals expanded wine quality-related clades and conserved gene structure organization across *Vitis* and *Arabidopsis* genomes. *BMC Plant Biol* **8**: 83
- Milkowski C, Strack D** (2004) Serine carboxypeptidase-like acyltransferases. *Phytochemistry* **65**: 517–524
- Nakashima J, Chen F, Jackson L, Shadle G, Dixon RA** (2008) Multi-site genetic modification of monolignol biosynthesis in alfalfa (*Medicago sativa*): effects on lignin composition in specific cell types. *New Phytol* **179**: 738–750
- Nesi N, Jond C, Debeaujon I, Caboche M, Lepiniec L** (2001) The *Arabidopsis* TT2 gene encodes an R2R3 MYB domain protein that acts as a key determinant for proanthocyanidin accumulation in developing seed. *Plant Cell* **13**: 2099–2114
- Pang Y, Peel GJ, Sharma SB, Tang Y, Dixon RA** (2008) A transcript profiling approach reveals an epicatechin-specific glucosyltransferase expressed in the seed coat of *Medicago truncatula*. *Proc Natl Acad Sci USA* **105**: 14210–14215
- Pang Y, Peel GJ, Wright E, Wang Z, Dixon RA** (2007) Early steps in proanthocyanidin biosynthesis in the model legume *Medicago truncatula*. *Plant Physiol* **145**: 601–615
- Paolucci F, Robbins MP, Madeo L, Arcioni S, Martens S, Damiani F** (2007) Ectopic expression of a basic helix-loop-helix gene transactivates parallel pathways of proanthocyanidin biosynthesis: structure, expression analysis, and genetic control of leucoanthocyanidin 4-reductase and anthocyanidin reductase genes in *Lotus corniculatus*. *Plant Physiol* **143**: 504–516
- Pfeiffer J, Kuhnel C, Brandt J, Duy D, Punyasiri P, Forkmann G, Fischer T** (2006) Biosynthesis of flavan 3-ols by leucoanthocyanidin 4-reductases and anthocyanidin reductases in leaves of grape (*Vitis vinifera* L.), apple (*Malus domestica* Borkh.) and other crops. *Plant Physiol Biochem* **44**: 323–334
- Prieur C, Rigaud J, Cheynier V, Moutounet M** (1994) Oligomeric and polymeric procyanidins from grape seeds. *Phytochemistry* **36**: 781–784
- Quattrocchio F, Verweij W, Kroon A, Spelt C, Mol J, Koes R** (2006) PH4 of *Petunia* is an R2R3 MYB protein that activates vacuolar acidification through interactions with basic-helix-loop-helix transcription factors of the anthocyanin pathway. *Plant Cell* **18**: 1274–1291
- Reid KE, Olsson N, Schlosser J, Peng F, Lund ST** (2006) An optimized grapevine RNA isolation procedure and statistical determination of reference genes for real-time RT-PCR during berry development. *BMC Plant Biol* **14**: 6–27
- Reinders A, Sivitz AB, Starker CG, Gantt JS, Ward JM** (2008) Functional analysis of LjSUT4, a vacuolar sucrose transporter from *Lotus japonicus*. *Plant Mol Biol* **68**: 289–299
- Souquet JM, Cheynier V, Brossaud F, Moutounet M** (1996) Polymeric proanthocyanidins from grape skins. *Phytochemistry* **43**: 509–512
- Souquet JM, Veran F, Mané C, Cheynier V** (2006) Optimization of extraction conditions on phenolic yields from the different parts of grape clusters: quantitative distribution of their proanthocyanidins. In F Daayf, A El Hadrami, L Adam, GM Ballance, eds, XXIII International Conference on Polyphenols, Winnipeg, Manitoba, Canada. Groupe Polyphénols, Bordeaux, France, pp 245–246
- Stracke R, Ishihara H, Huep G, Barsch A, Mehrstens F, Niehaus K, Weisshaar B** (2007) Differential regulation of closely related R2R3-MYB transcription factors controls flavonol accumulation in different parts of the *Arabidopsis thaliana* seedling. *Plant J* **50**: 660–677
- Stracke R, Werber M, Weisshaar B** (2001) The R2R3-MYB gene family in *Arabidopsis thaliana*. *Curr Opin Plant Biol* **4**: 447–456
- Su C, Singleton V** (1969) Identification of three flavan-3-ols from grapes. *Phytochemistry* **8**: 1553–1558
- Takos A, Jaffe F, Jacob S, Bogs J, Robinson S, Walker A** (2006) Light-induced expression of a MYB gene regulates anthocyanin biosynthesis in red apples. *Plant Physiol* **142**: 1216–1232
- Tanner GJ, Francki KT, Abrahams S, Watson JM, Larkin PJ, Ashton AR** (2003) Proanthocyanidin biosynthesis in plants: purification of legume leucoanthocyanidin reductase and molecular cloning of its cDNA. *J Biol Chem* **278**: 31647–31656
- Terrier N, Glissant D, Grimplet J, Barrieu B, Abbal A, Couture C, Ageorges A, Atanassova R, Léon C, Renaudin JP, et al** (2005) Isogene specific oligo arrays reveal multifaceted changes in gene expression during grape berry (*Vitis vinifera* L.) development. *Planta* **222**: 832–847
- Thompson JD, Higgins DG, Gibson TJ** (1994) CLUSTAL W: improving the sensitivity of progressive multiple sequence alignment through sequence weighting, position-specific gap penalties and weight matrix choice. *Nucleic Acids Res* **22**: 4673–4680
- Torregrosa L, Bouquet A** (1997) *Agrobacterium tumefaciens* and *Agrobacterium rhizogenes* co-transformation to obtain grapevine hairy roots producing the coat protein of grapevine chrome mosaic nepovirus. *Plant Cell Tissue Organ Cult* **49**: 53–62
- Torregrosa L, Verriès C, Tesniere C** (2002) Grapevine (*Vitis vinifera* L.) promoter analysis by biolistic mediated transient transformation of cell suspensions. *Vitis* **41**: 27–32
- Velasco R, Zharkikh A, Troggo M, Cartwright DA, Cestaro A, Pruss D, Pindo M, Fitzgerald LM, Vezzulli S, Reid J, et al** (2007) A high quality draft consensus sequence of the genome of a heterozygous grapevine variety. *PLoS One* **19**: e1326
- Verdonk JC, Haring MA, van Tunen AJ, Schuurink RC** (2005) ODOR-ANT1 regulates fragrance biosynthesis in petunia flowers. *Plant Cell* **17**: 1612–1624
- Verriès C, Guiraud JL, Souquet JM, Violet S, Terrier N, Ollé D** (2008) Validation of an extraction method on whole pericarp of grape berry (*Vitis vinifera* L. cv. Shiraz) to study biochemical and molecular aspects of flavan-3-ol synthesis during berry development. *J Agric Food Chem* **56**: 5896–5904
- Walker AR, Lee E, Bogs J, McDavid DAJ, Thomas MR, Robinson SP** (2007) White grapes arose through the mutation of two similar and adjacent regulatory genes. *Plant J* **49**: 772–785
- Wei YL, Li JN, Lu J, Tang ZL, Pu DC, Chai YR** (2007) Molecular cloning of Brassica napus TRANSPARENT TESTA 2 gene family encoding potential MYB regulatory proteins of proanthocyanidin biosynthesis. *Mol Biol Rep* **34**: 105–120
- Winkel-Shirley B** (2002) Biosynthesis of flavonoids and effects of stress. *Curr Opin Plant Biol* **5**: 218–223
- Xie DY, Dixon RA** (2005) Proanthocyanidin biosynthesis: still more questions than answers? *Phytochemistry* **66**: 2127–2144
- Zhang F, Gonzalez A, Zhao M, Payne CT, Lloyd A** (2003) A network of redundant bHLH proteins functions in all TTG1-dependent pathways of *Arabidopsis*. *Development* **130**: 4859–4869