



HAL
open science

Estimating the greenhouse gas fluxes of European grasslands with a process-based model: 1. Model evaluation from in situ measurements

Nicolas Vuichard, Jean-François J.-F. Soussana, Philippe Ciais, Nicolas Viovy, Christof Ammann, Pierluigi Calanca, John Clifton-Brown, Jürg Fuhrer, Mike Jones, Cécile Martin

► To cite this version:

Nicolas Vuichard, Jean-François J.-F. Soussana, Philippe Ciais, Nicolas Viovy, Christof Ammann, et al.. Estimating the greenhouse gas fluxes of European grasslands with a process-based model: 1. Model evaluation from in situ measurements. *Global Biogeochemical Cycles*, 2007, 21 (1 - Art. No GB1004), pp.1-14. 10.1029/2005GB002611 . hal-02667117

HAL Id: hal-02667117

<https://hal.inrae.fr/hal-02667117>

Submitted on 20 Jul 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Copyright

Estimating the greenhouse gas fluxes of European grasslands with a process-based model:

1. Model evaluation from in situ measurements

Nicolas Vuichard,^{1,2} Jean-François Soussana,³ Philippe Ciais,¹ Nicolas Viovy,¹ Christof Ammann,⁴ Pierluigi Calanca,⁴ John Clifton-Brown,^{5,6} Jürg Fuhrer,⁴ Mike Jones,⁵ and Cécile Martin⁷

Received 2 September 2005; revised 20 July 2006; accepted 28 August 2006; published 19 January 2007.

[1] We improved a process-oriented biogeochemical model of carbon and nitrogen cycling in grasslands and tested it against in situ measurements of biomass and CO₂ and CH₄ fluxes at five European grassland sites. The new version of the model (PASIM) calculates the growth and senescence of aboveground vegetation biomass accounting for sporadic removals when the grassland is cut and for continuous removals when it is grazed. Limitations induced by high leaf area index (LAI), soil water deficits and aging of leaves are also included. We added to this a simple empirical formulation to account for the detrimental impact on vegetation of trampling and excreta by grazing animals. Finally, a more realistic methane emission module than is currently used was introduced on the basis of the quality of the animals' diet. Evaluation of this improved version of PASIM is performed at (1) Laqueuille, France, on grassland continuously grazed by cattle with two plots of intensive and extensive grazing intensities, (2) Oensingen, Switzerland, on cut grassland with two fertilized and nonfertilized plots, and (3) Carlow, Ireland, on grassland that is both cut and grazed by cattle during the growing season. In addition, we compared the modeled animal CH₄ emissions with in situ measurements on cattle for two grazing intensities at the grazed grassland site of Laqueuille. Altogether, when all improvements to the PASIM model are included, we found that the new parameterizations resulted into a better fit to the observed seasonal cycle of biomass and of measured CO₂ and CH₄ fluxes. However, the large uncertainties in measurements of biomass and LAI make simulation of biomass dynamics difficult to make. Also simulations for cut grassland are better than for grazed swards. This work paves the way for simulating greenhouse gas fluxes over grasslands in a spatially explicit manner, in order to quantify and understand the past, present and future role of grasslands in the greenhouse gas budget of the European continent.

Citation: Vuichard, N., J.-F. Soussana, P. Ciais, N. Viovy, C. Ammann, P. Calanca, J. Clifton-Brown, J. Fuhrer, M. Jones, and C. Martin (2007), Estimating the greenhouse gas fluxes of European grasslands with a process-based model: 1. Model evaluation from in situ measurements, *Global Biogeochem. Cycles*, 21, GB1004, doi:10.1029/2005GB002611.

¹Laboratoire des Sciences du Climat et de l'Environnement, CEA-CNRS, Gif-sur-Yvette, France.

²Now at Department of Forest Science and Environment, University of Tuscia, Viterbo, Italy.

³INRA, UR874 Grassland Ecosystem Research, Clermont-Ferrand, France.

⁴Agroscope Reckenholz-Tänikon, Research Station ART Air Pollution and Climate Group, Zurich, Switzerland.

⁵Botany Department, Trinity College of Dublin, Dublin, Ireland.

⁶Now at Plant Breeding and Genetics, IGER, Aberystwyth, UK.

⁷Unité de Recherches sur les Herbivores, INRA Theix, St. Genès Champanelle, France.

1. Introduction

[2] Cultivated grasslands cover 20% of the European continent and are distributed about equally between western Europe (80 Mha) and eastern Europe (60 Mha). Despite this contribution to land cover, very little is known about their greenhouse gas budget. Most of the grasslands in Europe are cultivated for feeding animals, either directly by grazing or indirectly by forage grass production. European grasslands are amongst the most productive in the world. The amount of grassland biomass harvested for forage production, a fraction of net primary productivity (NPP), is in the range of 2–8 t C ha⁻¹ yr⁻¹ [Corrall, 1988] which is similar to NPP of European temperate forests [Schulze, 2000]. To sustain a high productivity, European grasslands are gener-

ally supplied with N fertilizers and consequently they emit N_2O to the atmosphere [Jarvis *et al.*, 2001]. N_2O is a powerful greenhouse gas, about 300 times more powerful than CO_2 (per unit mass) on a 100-years time horizon [Intergovernmental Panel on Climate Change, 1995]. Typical N_2O emissions from grassland soils, converted into CO_2 equivalent sources on a 100-years time horizon [Bouwman, 1996] range between 0.1 and 1 t eq C $\text{ha}^{-1} \text{yr}^{-1}$ [Machefert *et al.*, 2002; Sozanska *et al.*, 2002]. Extrapolated to the entire area of European grasslands, this is equivalent to a CO_2 source of 0.14 GtC yr^{-1} , i.e., one sixth of the EU-15 fossil fuel emissions. European grasslands also sustain $\sim 15\%$ of the global ruminant population (150 millions of cows and 150 millions of sheep) (FAOSTAT data, 2004, available at faostat.fao.org). Animals are direct sources of CO_2 via their metabolic activity. A cow respire for example about 1 tC yr^{-1} , a flux which bypasses the soil and vegetation respiration but needs to be accounted for in closing the carbon budget of grassland ecosystems. Grazing animals emit CH_4 to the atmosphere (23 times more powerful than CO_2 [Intergovernmental Panel on Climate Change, 1995]), typically 0.05 to 0.25 t CH_4 per animal and per year (0.3 to 1.5 t eq C [Vermorel, 1995]). In addition, they have an impact on the cycling of C and N in pastures via grazing, urine and faecal returns, and mechanical disturbance (trampling). Overall, compared to the CO_2 fluxes in terms of radiative forcing, the fluxes of CH_4 and N_2O over grasslands are so important that they need to be accounted for in any evaluation of the European grasslands greenhouse gas budget [Soussana *et al.*, 2004].

[3] The greenhouse gas budget of European grasslands is highly uncertain as there have been very few direct measurements of the fluxes with a sufficiently long-term continuity. However, a network of nine new sites was established as part of the GreenGrass project in 2002 (European Commission DG Research Vth Framework Programme, contract EVK2-CT2001-00105). There are also few continental-scale model-derived estimates of greenhouse gas budget of grasslands. Vleeshouwers and Verhagen [2002], further quoted by Janssens *et al.* [2003] developed a semi-empirical model of land use that induced soil carbon disturbances to the European continent, and they inferred a carbon sink of 101 TgC yr^{-1} over grasslands (0.52 tC $\text{ha}^{-1} \text{yr}^{-1}$) with uncertainties of similar magnitude to the mean. Regarding N_2O fluxes, Freibauer [2003] and Boeckx and VanCleemput [2001] estimated emissions of agricultural land using emission factors. They obtained a total N_2O emission of 442 ± 116 Gg $\text{N}_2\text{O-N}$ yr^{-1} (0.056 Gt-eq $\text{CO}_2\text{-C}$), of which 165 ± 48 Gg $\text{N}_2\text{O-N}$ yr^{-1} was emitted by grasslands. Other studies were made by Brown *et al.* [2002] for the United Kingdom and by de Vries *et al.* [2003] for the Netherlands which yielded national average estimates. Methane emissions due to animal enteric fermentation have been mainly determined at the animal level (Murray *et al.* [2001] for sheep; Vermorel [1995] for cattle) and then upscaled to the regional or continental level using statistical information on animal numbers by category [Crutzen *et al.*, 1986].

[4] These studies have no doubt delivered improved quantification of individual components of grasslands

greenhouse gas budget but they do not allow a comparison of sources and sinks of CO_2 with sources of CH_4 and of N_2O . To our knowledge, only the recent study of Freibauer [2003] has combined flux estimates for N_2O , CH_4 and CO_2 . In that study, regional budgets are calculated using emission factors for each gas and the only flux component of the CO_2 budget is the one related to soil disturbance on farmed organic soils: a process which always was assumed to act as a net source to the atmosphere. When upscaled using emission factors, the fluxes of each gas are decoupled from each other and the process-related interactions between them cannot be properly accounted. For instance, increasing N fertilization leads to higher N_2O emissions, but there is a concurrent stimulation of NPP, which fosters CO_2 uptake. Current approaches lack consistent handling of management effects on greenhouse gas emissions from grasslands and therefore hinder any reliable predictions under future climate and management changes.

[5] In this paper we develop a new modeling approach with the objective of quantifying how climate and management conditions determine greenhouse gas emissions for grasslands in a spatially and temporally explicit manner. Several process-based models of grasslands have been developed [Sheehy and Johnson, 1988], and applied at different scales from plot [Gilmanov *et al.*, 1997; Riedo *et al.*, 1999] to region [Mougin *et al.*, 1995; Schapendonk *et al.*, 1998]. Some of these models describe vegetation for an agronomic purpose or for an economic one by predicting the amount of harvested forage grass [Riedo *et al.*, 1999; Schapendonk *et al.*, 1998]. Others account for the fate of carbon and nitrogen in the soils [Parton *et al.*, 1988]. Few simulation models are designed to simulate simultaneously CO_2 fluxes exchanged with vegetation, soil and animals and the atmosphere, N_2O emissions produced by soil processes and CH_4 emissions due to animals. We improve a process-based model of grassland biogeochemistry called PASIM [Riedo *et al.*, 1998]. The PASIM model was evolved from the Hurley-Pasture Model [Thornley, 1998] and has been to date only tested at Swiss grassland sites harvested for forage production, for simulating vegetation variables [Riedo *et al.*, 2000] and N_2O emissions [Schmid *et al.*, 2001a, 2001b]. Here we enhance the model and apply it to grazed sites and include CH_4 emissions by animals. We selected three temperate grassland sites (of which two compare two treatments) used within the GreenGrass project, located in Ireland, France and Switzerland, which represent contrasting soil and climate conditions for both grazing and cutting management schemes. Two of these sites include high- and low-nitrogen fertilizer applications regimes. We evaluate the PASIM model against in situ NEE measurements made on a continuous basis by the eddy covariance technique, and against shoot biomass and leaf area index measurements. In addition, we evaluate the modeled CH_4 emissions against flux data from Laqueuille.

[6] In this paper, we first describe the model and new improvements; second, we describe the different sites and data sets; and third, the modeled and observed data were compared to analyze the model's performance under different management regimes. In a companion paper, we will




Figure 1. Flow diagram of PASIM model. The grey boxes represent management drivers, treated as boundary conditions for the model.

apply the same model to upscale the greenhouse gas budget of grasslands for the European continent.

2. Methods

2.1. Grassland Model

2.1.1. Key Processes

[7] PASIM is a process-based grassland biogeochemical model derived by *Riedo et al.* [1998] from the Hurley Pasture Model (HPM) [Thornley, 1998]. The program contains 8000 lines and it is written in ACSL language [Mitchell and Gauthier Associates, 1993]. As for other advanced biogeochemical models, PASIM simulates the carbon and nitrogen cycles, the latter having been improved by *Schmid et al.* [2001b], who added a detailed description of the nitrification and denitrification processes leading to nitrous oxide emissions. In PASIM, the photosynthesis model is a multilayered one based on a nonrectangular hyperbola [Riedo et al., 1998]. This model takes into account sun and shaded leaf fractions. Carbon assimilated by photosynthesis on a time step of 30 minutes is allocated dynamically to 1 root and 3 different shoot biomass compartments (each of those being dissociated in 4 age compartments), or is respired by autotrophic processes (Figure 1). Aboveground biomass is either cut for yield production, grazed by animals or accumulates on the soil surface owing to death and detachment. Typically 3 to 5% of the total carbon ingested by animals is emitted back to the atmosphere as CH₄. The nitrogen cycle in PASIM considers three different types of N inputs to the soil via atmospheric deposition, fertilizer addition, and symbiotic fixation by

clover. A fraction of the added soil nitrogen is available for plant growth and subsequently taken up by roots, whereas another fraction of soil nitrogen is lost through leaching, volatilization and nitrification/denitrification, the latter processes being responsible for N₂O gas emission to the atmosphere. Three distinct management practices are treated as boundary conditions to the model: (1) grazing intensity estimated from the cattle stocking density (animals ha⁻¹), (2) harvesting as controlled by the prescribed timing of cuts and amount of harvested biomass, and (3) timing and amount of N fertilizer applications. We have used the PASIM model in its standard configuration [Riedo et al., 1998; Schmid et al., 2001b], with the same parameter values as in Riedo for all sites, except for the key changes described below (see also Appendix A).

2.1.2. Effects of Animals' Diet on CH₄ Emissions

[8] Methane emissions by animals were previously calculated as a fixed proportion of the ingested carbon. In reality, the main factors responsible for CH₄ production are not only the amount but also the quality of the diet [Blaxter and Clapperton, 1965]. We thus introduced a quality factor in the equation governing CH₄ emission from cattle. Thus we let the CH₄ production (hereafter called m) depend on the DNDFI, the amount of digestible fiber in the animal's intake (hereafter called q). Regression model of *Pinarès-Patino et al.* [2007] gives

$$m = aq + b, \quad (1)$$

where m is the daily methane emission per kg of animal live weight (gCH₄ kg⁻¹ day⁻¹), q is the amount of digestible

Table 1. Proportion of Fiber in Ingested Biomass (NDF) and Fraction of Digestible Fiber in the Total Ingested Fiber (DNDF) for Lamina, Sheath, or Ear Tissues of Different Ages

	DNDF				NDF
	Age 1	Age 2	Age 3	Age 4	
Lamina	0.92	0.82	0.76	0.74	60
Sheath	0.84	0.65	0.53	0.50	70
Ear	0.76	0.48	0.30	0.26	80

fiber in the intake (kg day^{-1}). The slope and intercept values are $a = 0.045 \text{ gCH}_4 \text{ kg}^{-1} \text{ kg}^{-1}$ and $b = 0.287 \text{ gCH}_4 \text{ kg}^{-1} \text{ day}^{-1}$. The intake q can further be expressed as the product

$$q = dfi, \quad (2)$$

where i is the amount of organic dry-matter biomass ingested per animal per day ($\text{kg DM day}^{-1} \text{ animal}^{-1}$), f is the fraction of fiber in the intake, often called NDF (dimensionless) and d is the fraction of digestible fiber in total fibers, often called DNDF (dimensionless). In PASIM, we write f and d as the weighted means of the different leaf, sheath and ear pools of shoot biomass. This is expressed by

$$f = \frac{\sum_i w_i f_i}{\sum_i w_i} \quad (3)$$

$$d = \frac{\sum_i \sum_{j=1..4} w_{ij} d_{ij}}{\sum_i \sum_{j=1..4} w_{ij}}, \quad (4)$$

where w_{ij} is the amount of the pool i (age j) ingested by the animal, and d_{ij} is the fraction of digestible fibers for pool i (age j) defined according to Groot [1999]. The values of these parameters are given in Table 1.

2.1.3. Water and LAI Limitations on Vegetation Dynamics

[9] Vegetation growth is simulated for four biomass compartments of different ages. Biomass flows from a compartment to the next one at a turnover rate k . Thus biomass in the i^{th} age compartment (w_i) follows a first-order differential equation given by

$$\frac{dw_i}{dt} = w_i(1 - k) + kw_{i-1}. \quad (5)$$

[10] In the first age compartment, however, the input flux corresponds to the allocation of assimilates. In HPM, $k = k_{20} f_T / f_W$, where f_T is a dimensionless stress factor relative to temperature, f_W a stress factor relative to soil water deficit and k_{20} the turnover rate at 20°C without any stress. In the initial version of PASIM, f_W was arbitrarily set to 1, thus ignoring water limitations effects on vegetation age dynamics. We introduce a simple formulation of f_W given by

$$f_W = 0.5(1 + f_{Wp}), \quad (6)$$

where f_{Wp} is the water stress effect on photosynthesis, and varies between 0 and 1 [Riedo *et al.*, 1998]. With that new formulation, the turnover rate of vegetation exposed to strong water deficit conditions ($f_{Wp} \approx 0$) reaches up to twice the value in normal conditions.

[11] In addition, we also impose a limitation of vegetation turnover rates driven by leaf area index (LAI) values. As suggested by Bouman *et al.* [1996] in the LINGRA model, “with increasing LAI, the deeper layers of the crop become shaded, the low light intensities initiate remobilization of nitrogen from the shaded leaves and the leaves go through a stage of rapid senescence.” The LAI limitation factor f_L , acts to increase the turnover rate at high LAI value. As in the work by Bouman *et al.* [1996], we assume $f_L = 1$ for $\text{LAI} < 4$, and f_L increasing linearly with LAI to reach a maximum value of 2.5 for $\text{LAI} = 8$. The expression of f_L writes

$$f_L = \max \left[1, 0.5 \left(\min(8, \text{LAI}) \frac{3}{4} - 1 \right) \right]. \quad (7)$$

[12] The effects of high LAI and of soil water deficit in increasing turnover rates (shortening turnover times) are not considered to be additive. In presence of both water and LAI limitations, the value of k is given by

$$k = k_{20} f_T \max \left(\frac{1}{f_W}, f_L \right). \quad (8)$$

2.1.4. Senescence Impacts on Vegetation Dynamics

[13] Both growth and senescence are governed by the allocation of assimilates and by the turnover rate k of the vegetation age cohorts. During mortality periods, when no new assimilates get allocated to the tissues, the mass of each age cohort follows an exponential decay with time. This formulation causes a delay in the mortality of each compartment and of the plant system as a whole. The half-life of a one-compartment system is $\ln(2)/k$. Contrary to what we know on plant physiology [Thornley, 1998], with a cascade of 4 compartments of same turnover k , the half-life of the system of pools is not $4 * \ln(2)/k$. For example, for $k = 0.05 \text{ day}^{-1}$ (the nominal value in PASIM at 20°C and without any stress), the half-life of the 4-compartment system has been defined by numerical integration to 73.5 days, instead of 55.4 days if using the $4 * \ln(2)/k$. Thus the more age dependent the pools, the longer the turnover rate of the system will be. To correct for this bias, we introduce in the model a senescence term, which accelerates the turnover when the age of a compartment is above a certain threshold. This senescence term $k_{S,i}$ is added to the turnover k of each compartment i , according to the following formula suggested by Krinner *et al.* [2005],

$$k_{S,i} = \min \left(1, \frac{1}{a_{C,i} (a_{C,i}/a_i)^4} \right) \quad \text{if } a_i > a_{C,i}/2; \\ k_{S,i} = 0 \quad \text{if } a_i < a_{C,i}/2, \quad (9)$$

where a_i is the age of biomass in the compartment i and $a_{C,i}$ is a critical age equals to multiples of the half-life of the first




Figure 2. Biomass dynamics for a system of one age compartment (pluses), two age-compartments (circles), three age compartments (crosses) and four age compartments (triangles). The thick lines represent the biomass dynamic with the added senescence term ($k_{S,i}$, see text) and the thin ones represent that without. The curves were generated by a simple model using Scilab with a value of the turnover rate $k = 0.05 \text{ day}^{-1}$ and by injecting an arbitrary biomass input of 100 units on day 0 into the first age compartment and no input thereafter.

compartment: $a_{C,1} = \ln(2)/k$, $a_{C,2} = 2 * \ln(2)/k$, $a_{C,3} = 3 * \ln(2)/k$ and $a_{C,4} = 4 * \ln(2)/k$. Using a turnover rate value of $k = 0.05 \text{ day}^{-1}$, we obtain for example $a_{C,1}$, $a_{C,2}$, $a_{C,3}$ and $a_{C,4}$ values of respectively 13.9, 27.7, 41.6 and 55.4 days.

[14] The age of biomass in the compartment i , a_i is calculated at each step from the input of younger material issued from compartment $(i - 1)$ and the loss of material to $(i + 1)$ according to

$$\frac{da_i}{dt} = \frac{w_i a_i (1 - k_{tot,i}) + k_{tot,i-1} w_{i-1} a_{i-1}}{w_i (1 - k_{tot,i}) + k_{tot,i-1} w_{i-1}} + 1, \quad (10)$$

where $k_{tot,i}$ is the total turnover rate, including the senescence term of (9), as given by

$$k_{tot,i} = k + k_{S,i}. \quad (11)$$

[15] Figure 2 shows how adding a senescence term, which accelerates the turnover of aging biomass pools reduces the half-life of vegetation biomass. This change in the PASIM model prevents the half-life of the green material being overestimated, and avoids errors in the dynamics of carbon and nitrogen pools.

2.1.5. Grazing Impact on Vegetation Dynamics

[16] In the previous version of PASIM, the impact of defoliation and intake by animals was accounted for but the detrimental effects of trampling and of excretal returns (e.g., urine scorching) [Guthery and Bingham, 1996] was not. We simulate these extra processes by removing at each time step a fixed proportion k_G of the above ground biomass, which flows to the litter compartment. The value of the rate of removal k_G (day^{-1}) depends on animal stocking rate s

(Livestock Units per hectare and per day, $\text{LSU ha}^{-1} \text{ day}^{-1}$), and it is given by

$$k_G = gs, \quad (12)$$

where g is expressed in $(\text{LSU ha}^{-1})^{-1}$ and equals to 0.008, implying that an additional 0.8% of the aboveground herbage biomass is returned each day to litter for an instantaneous stocking rate of 1 LSU ha^{-1} .

2.2. Flux and Pools Data From Representative Sites

[17] The results of the model are compared to carbon fluxes and pools measurements from three grasslands, Oensingen (OEN), Laqueuille (LAQ) and Carlow (CAR) over the period mid-2002 to mid-2003. Two of these grasslands (OEN and LAQ) compare two experimental sites. Table 2 gives the timing and characteristics of grassland management for the five sites. Details on the site-specific parameters values may be found in auxiliary Table S1¹.

2.2.1. Oensingen, Switzerland (Cut)

[18] The OEN grassland has been newly sown in spring 2001 with grass and clover. It is intensively managed grassland, located in Switzerland ($47^\circ 17'N$, $07^\circ 44'E$) at 450 m a.s.l., with an annual mean temperature of 9°C and annual precipitation of 1100 mm yr^{-1} . The soil type on OEN is stagic Cambisol (eutric) with a soil organic matter content of 3.5%. We use data from two distinct plots characterized by contrasted N treatments, OEN-HN (for High Nitrogen) and OEN-LN (for Low Nitrogen). The two plots were managed by cutting. At OEN-HN, five cuts were applied in 2002, together with five fertilizer applications, one in the early spring and one after each cut, summing up to a total amount of 210 kg N ha^{-1} . In 2003, an amount of 60 kg N ha^{-1} fertilizer was applied in the early growing season. At OEN-LN, only three cuts were applied in 2002 and no fertilizer was supplied during the whole period.

2.2.2. Laqueuille, France (Grazed)

[19] The Laqueuille grassland, LAQ, is semi-natural grazed grassland located in central France ($45^\circ 38'N$, $02^\circ 44'E$) at an elevation of 1040 m a.s.l. with an annual mean temperature of 8°C and annual precipitation of 1000 mm yr^{-1} . These climate conditions are quite comparable with those of the Oensingen site. The LAQ grassland is an andosol developed on basaltic rocks. It has a loamy texture and a high soil organic matter content (18% with a C:N ratio of 10.5). As for OEN, two distinct plots have been established, characterized by contrasted N treatments and cattle grazing intensity. Those sites are called respectively LAQ-HN (high nitrogen and intensive grazing) and LAQ-LN (low nitrogen and extensive grazing). The two plots were managed by continuous grazing with heifers from day 141 until day 292 in 2002. The mean animal stocking rate during this grazing season reached 2.2 and 1.3 LSU ha^{-1} for LAQ-HN and for LAQ-LN, respectively. Both plots received 80 kg N ha^{-1} of organic

¹Auxiliary material data sets are available at <ftp://ftp.agu.org/apend/gb/2005gb002611>. Other auxiliary material files are in the HTML.

Table 2. Characteristics of the Management Patterns of the Five Grassland Sites Used in This Study To Evaluate the Model^a

Site	Fertilization			Time of Cuts, DOY	Time of Grazing, DOY	Mean Stocking Rate, LSU ha ⁻¹
	Time, DOY	Qty, kg N ha ⁻¹	Type ^b			
OEN-HN	71	35	AN	135	none	none
	142	45	LM	176		
	182	35	AN	227		
	231	73	LM	261		
	273	30	AN	343		
	77 ^c	57	LM			
OEN-LN	none	none	none	163	none	none
				227		
				270		
LAQ-HN	14	80	LM	none	141 → 292	2.2
	171	80	AN			
LAQ-LN	14	80	LM	none	141 → 292	1.3
CAR	105	153	AN	155	197 → 321	1.6
	162	50	AN			
	85 ^c	150	AN			

^aThese are the amount and nature of N fertilizer applications, the timing of cuts and the timing and stocking rates of grazing.

^bAbbreviations are as follows: LM, liquid manure; AN, ammonium nitrate.

^cThis is day of year 2003.

fertilizer on day 14 in 2002, before the start of the treatments, and the intensive plot received an additional 80 kg N ha⁻¹ of ammonium nitrate on day 171 in 2002 (see Table 2).

2.2.3. Carlow, Ireland (Cut and Grazed)

[20] The Carlow grassland, CAR, has been resown with grass and clover in 2001. It is an extensively cut and grazed grassland located in Ireland (52° 52'N, 06° 55'W) at 56 m a.s.l.. The annual mean temperature is 9.5°C and annual precipitation 820 mm yr⁻¹. The soil is a medium to loamy textured grey/brown podsollic soil (4 % OM, C:N ratio of 9.4). During 2002–2003, the CAR grassland has been used for cutting and grazing. In 2002, the CAR grassland received two nitrogen fertilizer applications for an annual amount of 200 kg N ha⁻¹ (Table 2). The main application of 150 kg N ha⁻¹ occurred in spring. The CAR site has been cut once on day 155 in 2002 and it was subsequently grazed between day 197 and day 321 with a mean animal stocking rate of 1.6 LSU ha⁻¹ (see Table 2).

2.2.4. In Situ Flux and Pool Measurements

[21] At each of the three sites, GreenGrass experimental researchers measured the Net Ecosystem Exchange of CO₂ (NEE) on a quasicontinuous basis by the eddy covariance method (for Oensingen [Ammann *et al.*, 2004]). These data are reported here on a daily basis. Further, at the grazed site LAQ, the CH₄ emissions by cows were quantified in situ by the SF₆ dual tracer method [Johnson *et al.*, 1994]. Unlike NEE, these measurements of CH₄ fluxes are noncontinuous, and they span over 4 weekly periods in 2002. The SF₆ dual tracer method shows higher animal-to-animal variation than the open-circuit hood calorimetry (Cal) method [Boadi *et al.*, 2002]. Thus CH₄ measurements on Laqueuille have been performed on 7 cows leading to standard deviation always lower than 20% of the mean value.

[22] At approximately monthly intervals during the growing season, the aboveground mass was measured by cutting quadrats at ground level. The harvested herbage was removed, a subsample taken for leaf area determination and

the remainder was oven dried and weighted (see auxiliary Text S1 for greater detail).

3. Results: Model Evaluation Against Observations

3.1. Setup of Site-Specific Simulations

[23] We define a control simulation called S0, with the former version of the model without any of the changes presented in section 2.1 (version 3.5 [Riedo *et al.*, 1998; Schmid *et al.*, 2001b]). Then we increment successively LAI and water deficit limitations (simulation S1), age-dependent senescence (S2) and trampling impacts (S3). The model is driven at each site by observed gap-filled hourly time series of temperature, precipitation, and downwelling shortwave radiation. The soil texture is prescribed from measurements made on each site. The biomass and soil pools are always initialized to values closed to their steady state equilibrium values after a 50 years spinup during which climate and management practice of year 2002 are repeatedly cycled. Then we calculate the fluxes and biomass evolution for the period from mid-2002 to mid-2003. We provide a detailed comparison of the model output with observed shoot biomass, LAI and NEE for the three sites and for both N treatments at Oensingen and Laqueuille. In addition, we evaluate the model results against site measurements of herbivore CH₄ emissions at Laqueuille. For herbivore methane emissions, we use the S3 version of the model, but with and without the changes in the methane production module (equations (1) to (4)).

3.2. Oensingen Carbon Simulation

[24] For the cut and highly fertilized grassland of Oensingen, OEN-HN, the phase and amplitude of NEE, LAI and shoot biomass are well simulated by the control version S0 (Figure 3) when the observed dates of grass harvest are prescribed. All the S0-S2 runs are equally close to the measured data (Table 3). The model-data correlation coef-




Figure 3. (a) Model-data comparison for the highly fertilized cut grassland of Oensingen, OEN-HN, of net ecosystem exchange in $\text{kgC-CO}_2 \text{ ha}^{-1} \text{ d}^{-1}$. (b) Same for LAI ($\text{m}^2 \text{ m}^{-2}$). (c) Same for shoot biomass (kg DM m^{-2}). Vertical dashed lines represent cut events. Each column correspond to the simulations with the initial version of the model (S0), to which are added water and LAI limitations on vegetation dynamics (S1), and senescence (S2).

ficients for NEE are $R_{S0}^2 = 0.67$, $R_{S2}^2 = 0.73$ (Figure 4 for S2, intercept of $-4.4 \text{ kgC ha}^{-1} \text{ d}^{-1}$ and slope of 0.71). The root-mean squared difference between model and data values are $\text{RMS}_{\text{wsh},S0} = 0.1$ and $\text{RMS}_{\text{wsh},S2} = 0.1 \text{ kg DM m}^{-2}$ for shoot biomass (26% and 22% of the mean observed shoot biomass, respectively). For LAI simulations, we obtain root mean squared differences of $\text{RMS}_{\text{LAI},S0} = 1.4$ and $\text{RMS}_{\text{LAI},S2} = 1.1 \text{ m}^2 \text{ m}^{-2}$ (50% and 40% of the mean observed shoot biomass, respectively). It can be seen in Figure 3 that the modeled NEE in both S0 and S1 has an over-early start of the net CO_2 uptake in 2003. This bias is reduced in S2 since the higher leaf mortality delays the growth of biomass in spring. Overall, we find that when N is not limiting and under intensive cutting, PASIM simulates well the carbon fluxes and biomass dynamic of the Oensingen grassland. In particular, the model captures very well the recovery of biomass after cutting (Figure 3). In contrast, when no fertilizers are applied at this site, we can see in Figure 5, that the simulation S0 grossly overestimates the LAI during the peak of the growing season, by up to $7 \text{ m}^2 \text{ m}^{-2}$ as compared with the data (Table 3, $\text{RMS}_{\text{LAI},S0} = 3.9$) while NEE and shoot biomass at harvest dates are simulated well (Table 3, $\text{RMS}_{\text{wsh},S0} = 0.1$). Accounting for water and high LAI limitations on biomass growth in S1 decreases the

maximum LAI values ($\text{RMS}_{\text{LAI},S1} = 3.6$) in the right direction, but still significantly overestimates the early season LAI. This is also reflected in the overestimated CO_2 uptake during the early season, up to May ($R_{S1}^2 = 0.59$). This model bias is reduced in S2 where LAI drops to 0.5 in January–February ($\text{RMS}_{\text{LAI},S2} = 2.0$), causing NEE to be close to zero during the winter ($R_{S2}^2 = 0.72$, Figure 4, intercept of $-5.3 \text{ kgC ha}^{-1} \text{ d}^{-1}$ and slope of 0.89). This is because the added senescence effect induces enough leaf mortality to bring down LAI in the winter (Figure 5). In parallel, the simulation S2 reduces significantly the shoot biomass at the first harvest dates to 0.6 kg DM m^{-2} (Figure 5 and Table 3, $\text{RMS}_{\text{wsh},S2} = 0.2 \text{ kg DM m}^{-2}$). Finally, in the S2 version, the model still largely overestimates LAI but not the shoot biomass on OEN-LN (low N), while it calculates consistent LAI and shoot biomass values on OEN-HN (high N). At low N and low LAI leaves appear to have a lower area per unit mass than predicted by the model, which induces an overestimation of LAI for the OEN-LN site.

3.3. Laqueuille Carbon Simulation

[25] For the high-nitrogen and intensively grazed plot of Laqueuille, LAQ-HN, the simulation S0 gives unrealistically high values of LAI, with values of up to around $20 \text{ m}^2 \text{ m}^{-2}$.




Figure 4. Scatterplots of the modeled versus measured NEE (in $\text{kgC-CO}_2 \text{ ha}^{-1} \text{ d}^{-1}$) for the five sites of this study for the last version of the model (S2 for Oensingen sites and S3 for the grazed sites).

Similarly, shoot biomass is overestimated by a factor of 9 ($\text{RMS}_{\text{WSH},\text{S0}} = 1.3 \text{ kg DM m}^{-2}$) when compared to the data (Figures 6b and 6c). In S0, the model-data correlation coefficient for NEE is also very low ($R_{\text{S0}}^2 = 0.01$). Accounting for LAI and water limitations on vegetation dynamics in the simulation S1 greatly improves the fit to the data, with a gain of a factor of 2 (Table 3). Further, adding senescence in simulation S2 produces more realistic LAI values, especially in winter times. The best fit to the data for NEE, shoot biomass and LAI is finally obtained when adding the effect of trampling in the S3 simulation (Figure 6 and Table 3), albeit

the model still overestimates LAI and shoot biomass from April to June. In the simulation S3, the NEE variability is reduced, in closer agreement with the flux data (Figure 4, $R_{\text{S3}}^2 = 0.09$, intercept of $1.0 \text{ kgC ha}^{-1} \text{ d}^{-1}$ and slope of 0.39). Note that the NEE of the grazed Laqueuille grassland is nearly flat during the growing season because of the continuous biomass consumption by cows, in contrast with Oensingen which has large intraannual NEE variations due to the cuts (compare Figure 5 and Figure 6). This major difference between grazed and cut NEE, which determines the European patterns of NEE




Figure 5. Same as Figure 3 for the nonfertilized cut grassland of Oensingen, OEN-LN.

Table 3. Evaluation of the Modeled Shoot Biomass (WSH), LAI and NEE at the Five Different Grasslands Versus In Situ Measurements^a

	OEN-HN	OEN-LN	LAQ-HN	LAQ-LN	CARL
		<i>WSH, kg DM m⁻²</i>			
Observed mean value	0.5	0.6	0.1	0.2	0.2
RMS S0	0.1	0.1	1.3	1.1	0.5
RMS S1	0.1	0.1	0.6	0.5	0.3
RMS S2	0.1	0.2	0.4	0.3	0.2
RMS S3	0.3	0.3	0.1
		<i>LAI, m² m⁻²</i>			
Observed mean value	2.9	2.4	1.3	1.9	1.8
RMS S0	1.4	3.9	19.2	16.9	9.6
RMS S1	1.4	3.6	6.7	5.3	6.1
RMS S2	1.1	2.0	5.0	3.9	3.9
RMS S3	1.8	2.0	2.1
		<i>R² NEE, %</i>			
S0	67	60	1	0	75
S1	67	59	3	4	78
S2	73	72	4	9	82
S3	9	10	83

^aRMS, root mean squared difference of model versus data; R², correlation coefficient of model versus NEE data.

(see companion paper) is correctly captured by the model. For the low-nitrogen and extensively grazed plot of LAQ-LN, the simulation S0 run has similar shortcomings to those at LAQ-HN (Figure 7), with overestimated LAI, shoot biomass and NEE variability during the growing season. We find that adding successively LAI and water limitations (S1 run),

winter senescence (S2 run) and trampling (S3 run) in the model improves the fit to the data at LAQ-LN. The above conclusions regarding the model performances at LAQ-HN can thus be generalized to the low-nitrogen extensive conditions of LAQ-LN. This is illustrated by comparing Figure 6 and Figure 7. The best fit to the site data here again is




Figure 6. Same as Figure 3 for the fertilized and intensively grazed grassland of Laqueuille, LAQ-HN. We added the results of Simulation S3 which accounts for animal trampling in reducing the growth of aboveground biomass. Horizontal thick lines represent the grazing period.




Figure 7. Same as Figure 6 for the low-fertilized and extensively grazed grassland of Laqueuille, LAQ-LN.

obtained in the simulation S3 giving $R_{S3}^2 = 0.10$ for NEE (Figure 4, intercept of $1.9 \text{ kgC ha}^{-1} \text{ d}^{-1}$ and slope of 0.40), RMS for LAI of $2 \text{ m}^2 \text{ m}^{-2}$ and RMS for shoot biomass of 0.3 kg DM m^{-2} .

3.4. Carlow Carbon Simulations

[26] For the cut and grazed grassland site of Carlow, CAR, we observe in Figure 8 that NEE is correctly simulated in the simulation S0 ($R_{S0}^2 = 0.75$). However, the simulation S0 grossly overestimates both LAI and shoot biomass values at CAR (Figure 8 and Table 3) with a RMS for LAI of $9.6 \text{ m}^2 \text{ m}^{-2}$ and a RMS for shoot biomass of 0.5 kg DM m^{-2} . New parameterizations introduced in S1-3 tend to successively improve the NEE simulation (Figure 8 and Table 3), leading to a model-data correlation coefficient of 0.83 for S3 (Figure 4, intercept of $6.1 \text{ kgC ha}^{-1} \text{ d}^{-1}$ and slope of 1.07). Shoot biomass and LAI are also improved in S2 and S3 simulations. However, during the period of grazing, these versions of the model still overestimate the observed LAI and biomass, by up to a factor of 2.

3.5. Laqueuille CH_4 Flux Simulation

[27] At the intensively grazed and fertilized grassland, LAQ-HN, the animal methane emissions are well simulated (Figure 9a). The thin dashes in this figure represent the estimate obtained by applying the regression model of section 2.1.2. applied directly to the DNDFI data. The main source of misfit between the simulated and observed CH_4 flux is caused by errors in the prescribed values of NDF,

DNDF or intake parameters as seen in Figures 9b–9d. The NDF is correctly simulated (Figure 9b) but the model is not able to represent the dynamics of the DNDF and intake (Figures 9c and 9d). At the extensively grazed and less fertilized grassland, LAQ-LN, more or less the same conclusions can be drawn that at LAQ-HN (Figure 10). This gives us confidence in the fact that the model can reproduce CH_4 emissions by grazing animals in the case of two contrasted N treatments, although the main driver (of the modeled CH_4 flux) remains the prescribed animal density. The DNDF data show a decrease from the second measurement period while the simulated DNDF increases at that period (Figure 10c). The intake is poorly simulated for the third measurement period (Figure 10d). In all cases, our new methane emission equations give better results than the former parameterizations. This is illustrated by comparing the dotted and dash-dotted lines for both LAQ-HN (Figure 9a) and LAQ-LN (Figure 10a).

4. Discussion and Concluding Remarks

[28] We first need to keep in mind that a very stringent evaluation of the model is hindered by rather large uncertainties on biomass measurements, typically with an error of 15%, due in part to spatial heterogeneity. Eddy covariance flux measurements, due to gaps and night time respiration problems are also accompanied by an uncertainty (which may reach 20% for day hourly data and 40% for night hourly data [Moncrieff *et al.*, 1996]). The model is not




Figure 8. Same as Figure 6 for the mixed cut and grazed grassland of Carlow, CAR. Vertical dashed lines represent cut events and horizontal thick lines grazing periods.

perfect either. Uncertainty analysis of the annual cumulated NEE (e.g., NEP) simulated by PASIM (using 50 version) has been performed by *Gottschalk et al.* [2007] for several European grasslands sites. They obtain a considerable variation of global uncertainty from site to site and between years (the standard deviation varying from 3% up to 100% of the mean NEP value) and conclude that the site specific combination and interaction of ecosystem parameters, driving forces and management make it impossible to define an absolute model output uncertainty. Regarding the simulated carbon balance of grasslands, it must also be kept in mind that all the runs are equilibrium calculations where the ecosystem is not disturbed. By construction the simulated long-term mean carbon balance, or Net Biome Productivity (NBP) equals zero. Thus the observed annual mean sink at LAQ-HN and LAQ-LN, or source at OEN-HN and OEN-LN cannot be simulated by PASIM. For this reason, we only discussed NEE variations during the year, but not the average annual NEE estimates.

[29] Generally, we found that (1) it is especially difficult to realistically simulate biomass dynamics, (2) the simulations for cut grasslands are better than for grazed ones, irrespective of the nitrogen treatments, and (3) accelerating the vegetation turnover during the growing season always improves the fit to the data. However, the model improvements benefit rather to shoot biomass and LAI than to NEE. Thus NEE improvements are loosely related to those made on LAI: for instance on LAQ-HN, the LAI is reduced by one order of magnitude among versions (from 20 to

2 m^2m^{-2} , Figure 6b) while model-data correlation coefficient for NEE does not vary by more than 9% (Figure 6a). This counterintuitive result is explained by the fact that full light interception is obtained at LAI values below $10 \text{m}^2\text{m}^{-2}$. Hence the high LAI values calculated in the S0 version do not increase photosynthesis as the additional leaf layers are fully shaded. Moreover, the N content of leaves increases with the number of the version used, as nutrients are concentrated in a smaller number of leaf layers, which tends to increase the photosynthesis per unit leaf area in the higher versions. As a result of these two processes, the annual GPP does not vary by more than 15% (data not shown) between the different versions of PASIM for grazed sites. The difference in the model performances between cut and grazed grasslands may be explained by the fact that sporadic cutting events enables the model to reset the vegetation variables to values close to zero and thus prevents the model from drifting away from the data after the onset of growth. In contrast, continuous defoliation by grazing has a weaker impact on vegetation than defoliation by cutting and it is more difficult to account for such a permanent disturbance in models. Moreover, grazing induces complex animal-vegetation interactions, which have an impact on sward level but are not simulated by PASIM. Because of the development of spatial heterogeneity in grazed paddocks, the herbage growth varies spatially which makes it more difficult to simulate using a model such as PASIM which assumes a spatial homogeneity. The correlations between simulated and observed NEE at the different sites confirm




Figure 9. (a) Model-data comparison of animals' daily CH_4 fluxes at the intensive LAQ-HN plot in $\text{gC-CH}_4 \text{ ha}^{-1} \text{ d}^{-1}$. (b) Proportion of fiber in the ingested biomass. (c) Proportion of digestible fiber in the total ingested fiber. (d) Animal intake ($\text{kg DM ha}^{-1} \text{ d}^{-1}$). The thin dashes mentioned as regression model are the CH_4 fluxes obtained when applying the empirical model of diet quality directly to the DNDFI data.




Figure 10. Same as Figure 9 but at the extensive LAQ-LN plot.

Table A1. Parameters and Variables of the Modified Equations

Variable	Definition	Unit	Value	Equation
m	Enteric methane emission per kg of live weight	$\text{gCH}_4 \text{ kg}^{-1} \text{ day}^{-1}$		1
q	Amount of digestible fiber in the intake	kg day^{-1}		1–2
d	Proportion of digestible fiber	-		2,4
f	Proportion of fiber in the intake	-		2,3
i	Amount of organic biomass ingested per animal and per day	$\text{kg DM day}^{-1} \text{ animal}^{-1}$		2
w	Shoot biomass	kg DM m^{-2}		3,4,10
k	Turnover rate of shoot biomass	day^{-1}		8,11
k_S	Turnover rate relative to senescence	day^{-1}		9
a_i	Age of biomass compartment of age i	day		9
$k_{\text{tot},i}$	Total turnover rate of shoot biomass	day^{-1}		10,11
k_G	Induced grazing impact	day^{-1}		12
Parameter				
a	Parameter of methane emission	$\text{gCH}_4 \text{ kg}^{-1} \text{ kg}^{-1}$	0.045	1
b	Parameter of methane emission	$\text{gCH}_4 \text{ kg}^{-1} \text{ day}^{-1}$	0.287	1
f_T	Parameter relative to temperature impact on k	-	[0–1]	8
f_W	Parameter relative to water deficit impact on k	-	[0–1]	6,8
f_{WP}	Parameter relative to water deficit impact on photosynthesis	-	[0–1]	6
f_L	Parameter relative to high LAI impact on k	-	[1–2.5]	7,8
k_{20}	Turnover rate of shoot biomass at 20°C	day^{-1}		8
$a_{C,i}$	Leaf critical age for biomass of age i	day		9
s	Instantaneous animal stocking rate	$\text{LSU ha}^{-1} \text{ day}^{-1}$		12
g	Parameter for defining k_G	$(\text{LSU ha}^{-1})^{-1}$	0.008	12

the fact that PASIM better simulates cut grasslands. The sawtooth shape of NEE due to harvest is well captured at both OEN and CAR cut sites, with an abrupt shift from CO_2 uptake to CO_2 release after harvest. In contrast, at the LAQ grazed site, the absence of cuts induces smaller amplitude NEE variations during the growing season, which the model does not capture very well.

[30] However, the poorer performances of the model at Laqueuille compared to other sites cannot only be attributed to grazing. The mismatch between data and model in early spring occurs even when no animals feed on the plot (Figures 6 and 7). Since weather conditions at LAQ are similar to those at OEN, a different response of the modeled NEE to temperature and precipitation cannot be inferred either. Laqueuille is a seminatural grassland whereas Oensingen is a newly sown one, a key difference explaining why the model parameterization of growth may not be adequate for Laqueuille. Since our ultimate goal is to use PASIM at the continental scale, the model must remain generic and flexible. As such, we did not attempt to adapt it for each type of grassland, mainly because sufficiently detailed regional information on grassland types is not currently available at the European level.

[31] Concerning animal CH_4 emissions, we found that the empirical model of Pinarès-Patino *et al.* [2007] can successfully be introduced in PASIM, yet with some difficulties to simulate some of the key input variables of the empirical model, such as the DNDF and intake. In PASIM, the intake is only driven by the shoot biomass averaged at the plot level. When sward spatial heterogeneity exists, the relation between intake and vegetation status may be much more complex and this may explain discrepancies between measurements and simulated intake. Moreover, to fully validate the model of Pinarès-Patino *et al.*, it will be necessary to test it across a large range of grasslands quality and with different types of animals.

[32] In conclusion, we have simulated grassland ecosystem fluxes for five contrasted grasslands with both grazing

and cutting managements, and with contrasted nitrogen fertilization applications. At each site, we evaluated the model against in situ continuous observations of eddy-covariance NEE, above ground shoot biomass and LAI. In addition, we compared the modeled animal CH_4 flux to direct measurements on cattle for two grazing intensities at the grazed grassland site of Laqueuille. We enhanced the original PASIM grassland model in order to include LAI, water limitation, and senescence effects on vegetation dynamics. In addition, we constructed a simple and empirical model of the detrimental impact on vegetation of trampling and excreta by grazing animals. Finally, a more realistic methane emission module was adapted to PASIM, on the basis of the quality of the animals' diet. Altogether, those new parameterizations resulted in a better fit to the observed carbon pools and to the measured CO_2 and CH_4 fluxes. The most important result is that the modified model is versatile enough to reproduce the seasonal dynamics of above ground biomass, and of CO_2 and CH_4 fluxes for a wide range of nitrogen fertilizer inputs, climate conditions, and for grazing versus cutting management. This site level evaluation study is a prerequisite for upscaling, in a companion paper, the model results to the scale of the European continent.

Appendix A

[33] Table A1 presents the variables and parameters of the equations of the PASIM model as modified in this paper.

[34] **Acknowledgment.** This work was funded by the EC under the contract EVK2-CT2001-00105 "GreenGrass."

References

- Ammann, C., C. Flechard, J. Fuhrer, and A. Nefel (2004), Greenhouse gas budget of intensively and extensively managed grassland, in *Land Use Systems in Grassland Dominated Regions*, edited by A. Lüscher, pp. 130–132, vdf Hochschulverlag AG, Zurich, Switzerland.

- Blaxter, K. L., and J. L. Clapperton (1965), Prediction of the amount of methane produced by ruminants, *Brit. J. Nutr.*, 19, 511–522.
- Boadi, D. A., K. M. Wittenberg, and A. D. Kennedy (2002), Validation of the sulphur hexafluoride (SF₆) tracer gas technique for measurement of methane and carbon dioxide production by cattle, *Can. J. Animal Sci.*, 82(2), 125–131.
- Boeckx, P., and O. VanCleemput (2001), Estimates of N₂O and CH₄ fluxes from agricultural lands in various regions in Europe, *Nutr. Cycling Agroecosyst.*, 60, 35–47.
- Bouman, B. A. M., A. H. C. M. Schapendonk, W. Stol, and D. W. G. van Kraalingen (1996), Description of LINGRA, a model approach to evaluate potential productivities of grasslands in different European climate regions, in *Quantitative Approaches in System Analysis 7*, p. 58, DLO Res. Inst. for Agrobiol. and Soil Fertil., Wageningen, Netherlands.
- Bouwman, A. F. (1996), Direct emission of nitrous oxide from agricultural soils, *Nutr. Cycling Agroecosyst.*, 46, 53–70.
- Brown, L., B. Syed, S. C. Jarvis, R. W. Sneath, V. R. Phillips, K. W. T. Goulding, and C. Li (2002), Development and application of a mechanistic model to estimate emission of nitrous oxide from UK agriculture, *Atmos. Environ.*, 36(6), 917–928.
- Corrall, A. J. (1988), Prediction of production from grassland, in *Information Bulletin of the FAO European Research Co-operative Network on Pastures and Field Crops, Herba 1*, pp. 25–28, Agron. Inst. of the Univ. of Florence, Florence, Italy.
- Crutzen, P. J., I. Aseilmann, and W. Seiler (1986), Methane production by domestic animals, wild ruminants, other herbivorous fauna, and humans, *Tellus, Ser. B*, 38, 271–284.
- de Vries, W., J. Kros, O. Oenema, and J. de Klein (2003), Uncertainties in the fate of nitrogen: II. A quantitative assessment of the uncertainties in major nitrogen fluxes in the Netherlands, *Nutr. Cycling Agroecosyst.*, 66, 71–102.
- Freibauer, A. (2003), Regionalised inventory of biogenic greenhouse gas emissions from European agriculture, *Eur. J. Agron.*, 19(2), 135–160.
- Gilmanov, T. G., W. J. Parton, and D. S. Ojima (1997), Testing the 'CENTURY' ecosystem level model on data sets from eight grassland sites in the former USSR representing a wide climatic/soil gradient, *Ecol. Modell.*, 96(1–3), 191–210.
- Gottschalk, P., M. Wattenbach, A. Neftel, J. Fuhrer, M. Jones, G. Lanigan, P. Davis, C. Campbell, J.-F. Soussana, and P. Smith (2007), Factors affecting uncertainty associated with estimates of Net Ecosystem Exchange by two ecosystem models at grassland sites in Europe, *Agric. Ecosyst. Environ.*, in press.
- Groot, J. C. J. (1999), Modelling grass digestibility on the basis of morphological and physiological plant characteristics, Ph.D. thesis, Wageningen Univ., Wageningen, Netherlands.
- Guthery, F. S., and R. L. Bingham (1996), A theoretical basis for study and management of trampling by cattle, *J. Range Manage.*, 49(3), 264–269.
- Intergovernmental Panel on Climate Change (1995), *Climate Change 1994: Radiative Forcing of Climate Change and an Evaluation of the IPCC IS92 Emission Scenarios*, 339 pp., Cambridge Univ. Press, New York.
- Janssens, I. A., et al. (2003), Europe's terrestrial biosphere absorbs 7 to 12% of European anthropogenic CO₂ emissions, *Science*, 300(5625), 1538–1542.
- Jarvis, S. C., S. Yamulki, and L. Brown (2001), Sources of nitrous oxide emissions in intensive grassland managements, *Phyton Horn Austria*, 41(3), 107–118.
- Johnson, K., M. Huyler, H. Westberg, B. Lamb, and P. Zimmerman (1994), Measurement of methane emissions from ruminant livestock using a SF₆ tracer technique, *Environ. Sci. Technol.*, 28(2), 359–362.
- Krinner, G., N. Viovy, N. de Noblet-Ducoudre, J. Ogee, J. Polcher, P. Friedlingstein, P. Ciais, S. Sitch, and I. C. Prentice (2005), A dynamic global vegetation model for studies of the coupled atmosphere-biosphere system, *Global Biogeochem. Cycles*, 19, GB1015, doi:10.1029/2003GB002199.
- Machefert, S. E., N. B. Dise, K. W. T. Goulding, and P. G. Whitehead (2002), Nitrous oxide emission from a range of land uses across Europe, *Hydrol. Earth Syst. Sci.*, 6(3), 325–337.
- Mitchell and Gauthier Associates (1993), *Advanced Continuous Simulation Language (ACSL) Reference Manual*, Concord, Mass.
- Moncrieff, J. B., Y. Malhi, and R. Leuning (1996), The propagation of errors in long-term measurements of land-atmosphere fluxes of carbon and water, *Global Change Biol.*, 2, 231–240.
- Mougin, E., D. Losen, S. Rambal, A. Gaston, and P. Hiernaux (1995), A regional Sahelian grassland model to be coupled with multispectral satellite data: 1. Model description and validation, *Remote Sens. Environ.*, 52(3), 181–193.
- Murray, P. J., E. Gill, S. L. Balsdon, and S. C. Jarvis (2001), A comparison of methane emissions from sheep grazing pastures with differing management intensities, *Nutr. Cycling Agroecosyst.*, 60, 93–97.
- Parton, W. J., J. W. B. Stewart, and C. V. Cole (1988), Dynamics of C, N, P and S in grassland soils: A model, *Biogeochemistry*, 5, 109–131.
- Pinarès-Patino, C. S., P. D'Hour, J. P. Jouany, and C. Martin (2007), Effects of stocking rate on methane and carbon dioxide emissions from grazing cattle, *Agric. Ecosyst. Environ.*, in press.
- Riedo, M., A. Grub, M. Rosset, and J. Fuhrer (1998), A pasture simulation model for dry matter production, and fluxes of carbon, nitrogen, water and energy, *Ecol. Modell.*, 105, 141–183.
- Riedo, M., D. Gyalistras, A. Fischlin, and J. Fuhrer (1999), Using an ecosystem model linked to GCM-derived local weather scenarios to analyse effects of climate change and elevated CO₂ on dry matter production and partitioning and water use in temperate managed grasslands, *Global Change Biol.*, 5, 213–223.
- Riedo, M., D. Gyalistras, and J. Fuhrer (2000), Net primary production and carbon stocks in differently managed grasslands: Simulation of site-specific sensitivity to an increase in atmospheric CO₂ and to climate change, *Ecol. Modell.*, 134, 207–227.
- Schapendonk, A. H. C. M., W. Stol, D. W. G. van Kraalingen, and B. A. M. Bouman (1998), LINGRA, a sink/source model to simulate grassland productivity in Europe, *Eur. J. Agron.*, 9(2–3), 87–100.
- Schmid, M., J. Fuhrer, and A. Neftel (2001a), Nitrous oxide concentrations in the soil of a mown grassland: Comparison of model results with soil profile measurements, *Water Air Soil Pollut. Focus*, 1, 437–446.
- Schmid, M., A. Neftel, M. Riedo, and J. Fuhrer (2001b), Process-based modelling of nitrous oxide emissions from different nitrogen sources in mown grassland, *Nutr. Cycling Agroecosyst.*, 60, 177–187.
- Schulze, E.-D. (2000), *Carbon and Nitrogen Cycling in European Forest Ecosystems*, 500 pp., Springer, New York.
- Sheehy, J. E., and I. R. Johnson (1988), Physiological models of grass growth, in *The Grass Crop: The Physiological Basis for Production*, edited by M. B. Jones and A. Lazenby, pp. 243–275, CRC Press, Boca Raton, Fla.
- Soussana, J. F., P. Loiseau, N. Vuichard, E. Ceschia, J. Balesdent, T. Chevallier, and D. Arrouays (2004), Carbon cycling and sequestration opportunities in temperate grasslands, *Soil Use Manage.*, 20, 219–230.
- Sozanska, M., U. Skiba, and S. Metcalfe (2002), Developing an inventory of N₂O emissions from British soils, *Atmos. Environ.*, 36(6), 987–998.
- Thornley, J. H. M. (1998), *Grassland Dynamics: An Ecosystem Simulation Model*, 241 pp., CABI Publ., Cambridge, Mass.
- Vermorel, M. (1995), Emissions annuelles de méthane d'origine digestive par les bovins en France, *Prod. Anim.*, 8(4), 265–272.
- Vleeshouwers, L. M., and A. Verhagen (2002), Carbon emission and sequestration by agricultural land use: A model study for Europe, *Global Change Biol.*, 8, 519–530.
- C. Ammann, P. Calanca, and J. Fuhrer, Agroscope Reckenholz-Tänikon, Research Station ART Air Pollution and Climate Group, Reckenholzstr. 191, CH-8046 Zurich, Switzerland.
- P. Ciais and N. Viovy, Laboratoire des Sciences du Climat et de l'Environnement, CEA-CNRS, Bat 701, Orme des Merisiers, F-91191 Gif-sur-Yvette, France.
- J. Clifton-Brown, Plant Breeding and Genetics, IGER, Plas Gogerddan, Aberystwyth, SY23 3EB, UK.
- M. Jones, Botany Department, Trinity College of Dublin, Dublin 2, Ireland.
- C. Martin, Unité de Recherches sur les Herbivores, INRA Theix, St Genès Champanelle F-63100, France.
- J.-F. Soussana, INRA, UR874 Grassland Ecosystem Research, 234 Av. du Brézet, F-63100 Clermont-Ferrand, France.
- N. Vuichard, Department of Forest Science and Environment, University Of Tuscia, Via de Lellis snc, I-01100 Viterbo, Italy. (vuichard@unitus.it)