Safety assessment of dairy microorganisms: The Lactococcus genus

Erick Casalta, Marie-Christine M.-C. Montel

► To cite this version:

Erick Casalta, Marie-Christine M.-C. Montel. Safety assessment of dairy microorganisms: The Lactococcus genus. International Journal of Food Microbiology, 2008, 126 (3), pp.271-273. 10.1016/j.ijfoodmicro.2007.08.013 . hal-02667687

HAL Id: hal-02667687

<https://hal.inrae.fr/hal-02667687>

Submitted on 31 May 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Safety assessment of dairy microorganisms: The *Lactococcus* genus[☆]

Erick Casalta^{a,*}, Marie-Christine Montel^b

^a INRA, UR45 Recherches sur le Développement de l'Élevage, Campus Grossetti, F-20250 Corté, France

^b INRA, UMT545 Recherches Fromagères, 36, rue de Salers, F-15000 Aurillac, France

Abstract

The *Lactococcus* genus includes 5 species. *Lactococcus lactis* subsp. *lactis* is the most common in dairy product but *L. garviae* has been also isolated. Their biotope is animal skin and plants. Owing to its biochemical characteristics, strains of *L. lactis* are widely used in dairy fermented products processing. Cases of human infections due to lactococci are very seldom reported even if *Lactococcus garviae* can be involved in fish diseases. Then *L. lactis* can be considered as safe and it is most commonly considered as Generally Recognized as Safe.

© 2008 Published by Elsevier B.V.

Keywords: *Lactococcus*; Taxonomy; Dairy use; Human safety

1. Introduction

The genus *Lactococcus* was proposed by Schleifer and colleagues in 1985 to reclassify some species of the genera *Streptococcus* (Lancefield group N lactic streptococci) and *Lactobacillus*. It has been defined on the basis of chemotaxonomic studies confirmed by 16s rRNA sequencing (Schleifer et al., 1985; Schleifer and Killper-Bälz, 1987; Collins et al., 1989). The *Lactococcus* genus includes five species, *L. garviae* (formerly *E. serolicida*), *L. piscium*, *L. plantarum*, *L. raffinolactis* (formerly *S. raffinolactis*) and *L. Lactis*, which is differentiated into subspecies *L. lactis* subsp. *cremoris*, *L. lactis* subsp. *hordniae* (formerly *Lactobacillus hordniae*) and *L. lactis* subsp. *lactis* (formerly *Lactobacillus xylosus*, *Streptococcus lactis*) (www.bacterio.cict.fr).

Lactococci are Gram positive cocci and belong to the group of Lactic Acid Bacteria. They are homofermentative and exclusively produce L(+) lactic acid. They are not β hemolytic and they are poorly α hemolytic.

2. Biotope and concerned food

Lactococci are generally found on plants and the skins of animals. *L. plantarum* is mainly isolated from plants, *L. garviae* from fish, animals and milk, and *L. piscium* from salmon (Williams et al., 1990). The presence of lactococci in raw milk is due to contamination from forage during milking. The two lactococci most commonly found in raw milk, cheese and other dairy products are *L. lactis* subsp. *lactis* and *L. lactis* subsp. *cremoris*. These two subspecies generally reach a high level ($>10^8$ CFU g⁻¹) as early as the first day of manufacturing and maintain it throughout the ripening period of many raw milk cheeses such as Camembert (Corrolier et al., 1999), Serra (Macedo et al., 1996), Venaco (Casalta, 2003) and Pecorino Sardo (Ledda et al., 1996). *L. raffinolactis* has occasionally been found in raw milk and cheeses (Perez Elortondo et al., 1999; Lopez-Diaz et al., 2002). *L. garviae* may also be isolated from raw milk (Villani et al., 2001) and raw milk cheeses: PDO Salers (Callon et al., 2004), Egyptian cheeses (El-Baradei et al., 2005), Jben cheese (Ouadghiri et al., 2005), Italian Piedmontese PDO Toma cheeses (Fortina et al., 2003). A study of 35 European artisanal dairy products indicated lactococci as the most commonly found LAB genus, accounting for 38% of the bacterial isolates identified (Cogan et al., 1997).

L. lactis subsp. *lactis* and to a lesser extent *L. lactis* subsp. *cremoris* have long been extensively used in starter cultures

[☆] Contribution to the safety assessment of technological microflora found in fermented dairy products, PART II.

* Corresponding author.

E-mail address: eca@corte.inra.fr (E. Casalta).

for dairy fermentation (i.e. in cheeses, sour cream and butter), composed of single or multiple strains with or without other lactic acid bacteria (Beresford et al., 2001). Their main role in dairy fermentation is acidification, mainly by producing L-lactic acid. They contribute to the development of texture by producing exopolysaccharides, or to flavor by producing aromatic compounds (alcohols, ketones, aldehydes) or by citrate, amino acid or fat metabolism (Smit et al., 2005). They can also be used for food preservation due to their ability to produce organic acids and bacteriocins, nisin being the best characterized and recognized (Delves-Broughton et al., 1996). Their use as probiotics has been also considered (Ouwehand et al., 1999). The annotation of the genomes of different subspecies will undoubtedly open up new prospects for identifying new useful functions in the species (Kok et al., 2005). It will also be a great help in assessing the safety of lactococci (Kok et al., 2005).

3. Taxonomy

The identification of the most common *Lactococcus* species found in dairy products can be successfully performed by rapid and accurate molecular techniques.

PCR-DGGE (denaturing gradient gel electrophoresis) analyses can be used for differentiating *Lactococcus lactis* from other lactic acid bacteria (Coppola et al., 2001). rRNA oligonucleotide probes have been designed for identifying *L. lactis* subsp. *cremoris* (Salama et al., 1991). Several PCR or multiplex PCR reactions are now available, exploiting the diversity of sequences of 16 S rRNA genes in *L. lactis* (Pu et al., 2002) and *L. garvieae* (Zlotkin et al., 1998) and the polymorphism of the 16 S-23 S rDNA spacer region (Blaiotta et al., 2002) or other functional genes, histidine biosynthesis operon (Corrolier et al., 1999), *acmA* gene (Garde et al., 1999), *sodA* gene (Fihman et al., 2006). The genetic diversity of *Lactococcus* can be analyzed by random polymorphism DNA (RAPD) (Tailliez et al., 1998) or multiple locus microsatellite analysis (Quénée et al., 2005) for *L. lactis* or by pulsed-field gel electrophoresis (PFGE) for *L. garvieae* (Vela et al., 2000).

Knowledge of the *L. lactis* genome will make it easy, in the future, to develop new genomic tools for increasingly reliable identification (Kok et al., 2005).

4. Safety assessment

Members of *Lactococcus* genus are most commonly classed as Generally Recognized as Safe (GRAS) (Salminen et al., 1998). *L. lactis* cannot be considered as an opportunist pathogen, as only two cases of endocarditis have been reported in the medical literature over a period of fifty years (Wood et al., 1955; Mannion and Rothburn, 1990).

Lactococcus garvieae has been associated with septicemic infections of fish, mainly in intensive modern aquaculture; it is the main risk factor for the Mediterranean European trout industry (Schmidtke and Carson, 2003; Vela et al., 2000; Eyngor et al., 2004). It has been also isolated from bovine mastitis (Teixeira et al., 1996). Human infections due to *L.*

garvieae are scarce, since only six cases have been described, all of which concerned elderly or immunosuppressed patients. *L. garvieae* has been incriminated in three cases of infection of prosthetic valves (Fihman et al., 2006), one case of native valve infection (Fefer et al., 1998), one case of osteomyelitis (James et al., 2000) and one case of bacteremia with a liver abscess (Mofredj et al., 2000). The number of cases may be underestimated as the bacterial agent responsible for endocarditis cases has not always been identified with certainty.

5. Conclusion

Lactococci are ubiquitous in the environment and in food. They are widely used as starters, with a long history of use for the sake of their technological properties, especially in dairy products, from small-scale manufacture to industrial-scale processes. Their identification no longer poses problems.

Lactococci from dairy products can be generally considered safe, in view of their extensive, daily consumption by humans and their low incidence in human infections.

References

- Beresford, T.P., Fitzsimons, N.A., Brennan, N.L., Cogan, T.M., 2001. Recent advances in cheese microbiology. *International Dairy Journal* 11, 259–274.
- Blaiotta, G., Pepe, O., Mauriello, G., Villani, F., Andolfi, R., Moschetti, G., 2002. 16 S-23 S rDNA intergenic spacer region polymorphism of *Lactococcus garvieae*, *Lactococcus raffinolactis* and *Lactococcus lactis* as revealed by PCR and nucleotide sequence analysis. *Systematic and Applied Microbiology* 25, 520–527.
- Callon, C., Millet, L., Montel, M.C., 2004. Diversity of lactic acid bacteria isolated from AOC Salers cheese. *Journal of Dairy Research* 71, 231–244.
- Casalta, E., 2003. Bases scientifiques de la qualité du Venaco, fromage traditionnel au lait cru. Mise au point de ferments sélectionnés spécifiques. Thèse de microbiologie, Université de Bourgogne. ENSBANA, Dijon (136 p.).
- Collins, M.D., Ash, C., Farrow, J.A.E., Wallbanks, S., Williams, A.M., 1989. 16 s ribosomal ribonucleic acid sequence analyses of lactococci and related taxa. Description of *Vagococcus fluvialis* gen. nov., sp. nov. *Journal of Applied Microbiology* 67, 453–467.
- Cogan, T.M., Barbosa, M., Beuvier, E., Bianchi-Salvadori, B., Cocconcelli, P.S., Fernandes, I., Gomez, J., Gomes, R., Kalantzopoulos, G., Ledda, A., Medina, M., Rea, M.C., Rodriguez, E., 1997. Characterisation of the lactic acid bacteria in artisanal dairy products. *Journal of Dairy Research* 64, 409–421.
- Coppola, S., Blaiotta, G., Ercolini, D., Moschetti, G., 2001. Molecular evaluation of microbial diversity occurring in different types of Mozzarella cheese. *Journal of Applied Microbiology* 90, 414–420.
- Corrolier, D., Desmasures, N., Gueguen, M., 1999. Correlation between polymerase chain reaction analysis of the histidine biosynthesis operon, randomly amplified polymorphic DNA analysis and phenotypic characterization of dairy *Lactococcus* isolates. *Applied Microbiology and Biotechnology* 51, 91–99.
- Delves-Broughton, J., Blackburn, P., Evans, R.J., Hugenoltz, J., 1996. Applications of the bacteriocin nisin. *Antonie van Leeuwenhoek* 69, 193–202.
- El-Baradei, G., Delacroix-Buchet, A., Pery, P., Ogier, J.C., 2005. Occurrence of *Lactococcus garvieae* in four types of Egyptian cheeses by specific polymerase chain reaction assay. *Egyptian Journal of Dairy Science* 33, 35–41.
- Eyngor, M., Zlotkin, A., Ghittino, C., 2004. Clonality and diversity of the fish pathogen *Lactococcus garvieae* in Mediterranean countries. *Applied and Environmental Microbiology* 70, 5132–5137.
- Fefer, J.J., Ratzan, K.R., Sharp, S.E., Saiz, E., 1998. *Lactococcus garvieae* endocarditis: report of case and review of the literature. *Diagnostic Microbiology and Infectious Disease* 32, 127–130.

- Fihman, V., Raskine, L., Barrou, Z., Kiffe, C., Riahi, J., Berçot, B., Sanson-Le Pors, M.J., 2006. *Lactococcus garvieae* endocarditis: identification by 16 S rRNA and *sodA* sequence analysis. *Journal of Infection* 52, e3–e6.
- Fortina, M.G., Ricci, G., Acquati, A., Zeppa, G., Gandini, A., Manachini, P.L., 2003. Genetic characterization of some lactic acid bacteria occurring in an artisanal protected denomination origin (PDO) Italian cheese, the Toma piemontese. *Food Microbiology* 20, 397–404.
- Garde, S., Babin, M., Gaya, P., Nuñez, M., Medina, M., 1999. PCR amplification of the gene *acmA* differentiates *Lactococcus lactis* subsp. *lactis* and *L. lactis* subsp. *cremoris*. *Applied and Environmental Microbiology* 65, 5151–5153. <http://www.bacterio.cict.fr>.
- James, P.R., Hardman, S.M., Patterson, D.L., 2000. Osteomyelitis and possible endocarditis secondary to *Lactococcus garvieae*: a first case report. *Postgraduate Medical Journal* 76, 301–303.
- Kok, J., Buist, G., Zomer, A.L., van Hijum, S.A.F.T., Kuipers, O.P., 2005. Comparative and functional genomics of lactococci. *FEMS Microbiology Reviews* 29, 411–433.
- Ledda, A., Floris, R., Mannu, L., Scintu, M.F., 1996. Studies on the microbial population and ecosystem of Pecorino Sardo cheese made from raw ewe's milk. *Flora 3rd plenary meeting, Thessaloniki*, pp. 89–95. 10–12 october.
- Lopez-Diaz, T.M., Alonso, C., Roman, C., Garcia-Lopez, M.L., Moreno, B., 2002. Lactic acid bacteria isolated from a hand made blue cheese. *Food Microbiology* 17, 23–32.
- Macedo, A.C., Costa, M.L., Malcata, F.X., 1996. Changes in the microflora of Serra cheese: evolution throughout ripening time, lactation period and axial location. *International Dairy Journal* 6, 79–94.
- Mannon, P.T., Rothburn, M.M., 1990. Diagnosis of bacterial endocarditis by *Streptococcus lactis* and assisted by immunoblotting of serum antibodies. *Journal of Infection* 21, 317–326.
- Mofredj, A., Baraka, D., Cadrane, J.F., Le Maitre, P., Kloeti, G., Dumont, J.L., 2000. *Lactococcus garvieae* septicemia with liver abscess in an immunosuppressed patient. *American Journal of Medicine* 109, 513–514.
- Ouadghiri, M., Amar, M., Vancanneyt, M., 2005. Biodiversity of lactic acid bacteria in Moroccan soft white cheese (Jben). *FEMS Microbiology Letters* 251, 267–271.
- Ouwehand, A.C., Kirjavainen, P.V., Grönlund, M., Isolauri, M., Salminen, S.J., 1999. Adhesion of probiotic micro-organisms to intestinal mucus. *International Dairy Journal* 9, 623–630.
- Perez Elortondo, F.J., Aldamiz Echobarria, P., Albisu, M., Barcina, Y., 1999. Indigenous lactic acid bacteria in Idiazabal ewes' milk cheese. *International Dairy Journal* 8, 725–732.
- Pu, Z.Y., Dobos, M., Limsowtin, G.K.Y., Powell, I.B., 2002. Integrated polymerase chain reaction-based procedures for the detection and identification of species and subspecies of the Gram-positive bacterial genus *Lactococcus*. *Journal of Applied Microbiology* 93, 353–361.
- Quénée, P., Lepage, E., Kim, W.S., Vergnaud, G., Gruss, A., 2005. Minisatellite polymorphism as a tool to distinguish closely related *Lactococcus lactis* strains. *FEMS Microbiology Letters* 248, 101–109.
- Salama, M., Sandine, W., Giovannoni, S., 1991. Development and application of oligonucleotide probes for identification of *Lactococcus lactis* subsp. *cremoris*. *Applied and Environmental Microbiology* 57, 1313–1318.
- Salminen, S., von Wright, A., Morelli, L., Marteau, P., Brassart, D., de Vos, W.M., Fondén, R., Saxelin, M., Collins, K., Mogensen, G., Birkeland, S.E., Mattila-Sandholm, T., 1998. Demonstration of safety of probiotics — a review. *International Journal of Food Microbiology* 44, 93–106.
- Schleifer, K.H., Kraus, J., Dvorak, C., Kilpper-Bälz, R., Collins, M.D., Fischer, W., 1985. Transfer of *Streptococcus lactis* and related streptococci to the genus *Lactococcus* gen. nov. *Systematic and Applied Microbiology* 6, 183–195.
- Schleifer, K.H., Kilpper-Bälz, R., 1987. Molecular and chemotaxonomic approaches to the classification of streptococci, enterococci and lactococci: a review. *Systematic and Applied Microbiology* 10, 1–19.
- Schmidtke, L.M., Carson, J., 2003. Antigen recognition by rainbow trout (*Oncorhynchus mykiss*) of whole cell proteins expressed by *Lactococcus garvieae* when obtained directly from fish and under iron limited culture conditions. *Veterinary Research* 34, 63–71.
- Smit, G., Smit, B.A., Engels, W.J.M., 2005. Flavour formation by lactic acid bacteria and biochemical flavour profiling of cheese products. *FEMS Microbiology Reviews* 29, 591–610.
- Tailliez, P., Tremblay, J., Ehrlich, S.D., Chopin, A., 1998. Molecular diversity and relationship within *Lactococcus lactis*, as revealed by randomly amplified polymorphic DNA (Rapid). *Systematic and Applied Microbiology* 21, 530–538.
- Teixeira, L.M., Merquior, V.L., Vianni, M.C.E., Carvalho, M.G.S., Fracalanza, S.E.L., Steigerwalt, A.G., Brenner, D.J., Facklam, R.R., 1996. Phenotypic and genotypic characterization of atypical *Lactococcus garvieae* strains isolated from water buffalos with subclinical mastitis and confirmation of *L. garvieae* as a senior subjective synonym of *Enterococcus seriolicida*. *International Journal of Systematic Bacteriology* 46, 664–668.
- Vela, A.I., Vazquez, J., Gibello, A., Blanco, M.M., Moreno, M.A., Liebana, P., Albendea, C., Alcalá, B., Mendez, A., Dominguez, L., Fernandez-Garayzabal, J.F., 2000. Phenotypic and genetic characterization of *Lactococcus garvieae* isolated in Spain from lactococcosis outbreaks and comparison with isolates of other countries and sources. *Journal of Clinical Microbiology* 38, 3791–3795.
- Villani, F., Aponte, M., Blaiotta, G., Mauriello, G., Pepe, O., Moschetti, G., 2001. Detection and characterization of a bacteriocin, garviecin L1-5, produced by *Lactococcus garvieae* isolated from raw cow's milk. *Journal of Applied Microbiology* 90, 430–439.
- Williams, A.M., Fryer, J.L., Collins, M.D., 1990. *Lactococcus piscium* sp. nov. a new *Lactococcus* species from salmonid fish. *FEMS Microbiology Letters* 68, 109–114.
- Wood, H.F., Jacobs, K., McCarty, M., 1955. *Streptococcus lactis* isolated from a patient with subacute bacterial endocarditis. *American Journal of Medicine* 18, 345–347.
- Zlotkin, A., Eldar, A., Ghittino, C., Bercovier, H., 1998. Identification of *Lactococcus garvieae* by PCR. *Journal of Clinical Microbiology* 36, 983–985.