



**HAL**  
open science

## Microbial dynamics in anaerobic enrichment cultures degrading di-n-butyl phthalic acid ester

Eric Trably, Damien Batstone, Nina Christensen, Dominique Patureau, Jens  
E. Schmidt

► **To cite this version:**

Eric Trably, Damien Batstone, Nina Christensen, Dominique Patureau, Jens E. Schmidt. Microbial dynamics in anaerobic enrichment cultures degrading di-n-butyl phthalic acid ester. *FEMS Microbiology Ecology*, 2008, 66 (2), pp.472-483. 10.1111/j.1574-6941.2008.00570.x . hal-02668691

**HAL Id: hal-02668691**

**<https://hal.inrae.fr/hal-02668691>**

Submitted on 9 Aug 2023

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

1

## 2 **Microbial Dynamics in Anaerobic Enrichment Cultures**

### 3 **Degrading Di-n-Butyl Phthalic Acid Ester**

4

5

6 Eric Trably<sup>1,3,\*</sup>, Damien J. Batstone<sup>1,2</sup>, Nina Christensen<sup>1</sup>, Dominique Patureau<sup>1,3</sup> and

7 Jens E. Schmidt<sup>1,4</sup>

8 <sup>1</sup> *Technical University of Denmark (DTU), Department of Environmental Engineering, Byg.113 Bygningstorvet, 2800*  
9 *Kgs. Lyngby, Denmark. [nic@env.dtu.dk](mailto:nic@env.dtu.dk),*

10 <sup>2</sup> *Present address:Advanced Wastewater Management Centre, The University of Queensland, St. Lucia, 4072*  
11 *Queensland, Australia. [damiemb@awmc.uq.edu.au](mailto:damiemb@awmc.uq.edu.au)*

12 <sup>3</sup> *Present address: INRA, UR050, Laboratoire de Biotechnologie de l'Environnement, Narbonne, F-11100, France.*  
13 *[trably@supagro.inra.fr](mailto:trably@supagro.inra.fr) ; [patureau@supagro.inra.fr](mailto:patureau@supagro.inra.fr)*

14 <sup>4</sup> *Present address NRG, Biosystems Department, Risø National Laboratory for Sustainable Energy, Technical*  
15 *University of Denmark, P.O. 49, DK-4000 Roskilde, Denmark; [jens.ejbye.schmidt@risoe.dk](mailto:jens.ejbye.schmidt@risoe.dk)*

16

#### 17 **[Correspondent footnote]**

18 \*corresponding author. Mailing Address: INRA, UR50, Laboratoire de Biotechnologie de l'Environnement, Avenue  
19 des Etangs, Narbonne, F-11100, France.

20 Phone: (+33) 468 425 172 . Fax: (+33) 468 425 160. Email: [trably@supagro.inra.fr](mailto:trably@supagro.inra.fr)

21

#### 22 **[Keywords]**

23 Anaerobic; Biodegradation; DBP; Fluorescent In Situ Hybridization; Sludge; Single Strand

24 Conformation Polymorphism

25

26 **[Abstract]**

27 Although anaerobic biodegradation of di-n-butyl phthalic acid ester (DBP) has been studied over  
28 the past decade, only a little is known about the microorganisms involved in the biological  
29 anaerobic degradation pathways. The aim of this work is to characterize the microbial community  
30 dynamics in enrichment cultures degrading phthalic acid esters under methanogenic conditions. A  
31 selection pressure was applied by adding DBP at 10 and 200 mg L<sup>-1</sup> in semi-continuous anaerobic  
32 reactors. The microbial dynamics were monitored by Single Strand Conformation Polymorphism  
33 (SSCP). While only limited abiotic losses were observed in the sterile controls (20 % - 22 %),  
34 substantial DBP biodegradation was shown in the enrichment cultures (90 % - 99 %). In addition,  
35 significant population changes were observed. The dominant bacterial species in the DBP-  
36 degrading cultures was affiliated to *Soehngenia saccharolytica*, a microbe previously described as  
37 anaerobic benzaldehyde degrader. Within the archaeal community, there was a shift between two  
38 different species of the genus *Methanosaeta* sp., indicating a highly specific impact of DBP or  
39 degradation products on archaeal species. RNA directed probes were designed from SSCP  
40 sequences, and Fluorescence *In Situ* Hybridization (FISH) observations confirmed the dominance  
41 of *Soehngenia saccharolytica*, and indicated floccular microstructures, likely providing favourable  
42 conditions for DBP degradation.

43

44 **[Introduction]**

45 Phthalic acid esters (PAEs) are specifically addressed by EU regulations because of their increasing  
46 amounts released in the environment. Over the past years, more than 900,000 tons of PAEs have  
47 been produced each year in Europe (ECPI, 2004). Most PAEs are used as plasticizers to increase  
48 the flexibility of the polyvinylchloride (PVC) resins, and also as additives in other resins such as  
49 polyvinyl acetates, celluloses, and polyurethanes (Staples *et al.*, 1997). The Di-n-Butyl Phthalic  
50 acid ester (DBP) is used as additive in epoxy resins, cellulose esters, specialized adhesive

51 formulations, and also as a solvent for dyes, insecticides and other organic compounds (Staples *et*  
52 *al.*, 1997). Widely found in urban wastewaters, DBP is surface active and highly hydrophobic, and  
53 readily adsorbs onto sludge organic particles during wastewater clarification. Therefore, DBP up  
54 concentrates to orders of magnitude above the values in the original wastewaters. By spreading  
55 sludge, DBP may not only accumulate in soils (Hu *et al.*, 2003; Mougin *et al.*, 2006; Patureau *et al.*,  
56 2007), but also readily transfer to plants and animals along the food chain (Yin *et al.*, 2003;  
57 Jarosova, 2006). Long term exposure to DBP can adversely affect human reproduction and  
58 development, as well as on plants and animals (CEHR, 2000; Kim *et al.*, 2002). Therefore,  
59 limitation of DBP contents in sludge before land disposal is strongly recommended. To date, only  
60 Denmark within the European Union has fixed a limit of  $50 \text{ mg kg}_{\text{TS}}^{-1}$  for the phthalic acid ester  
61 concentration in sludge. Future European Union legislation will probably fix a target limit value of  
62  $100 \text{ mg kg}_{\text{TS}}^{-1}$  for PAEs (Directive 2455/2001/EC).

63 Several studies have been conducted to assess PAE biodegradability under aerobic (Angelidaki *et*  
64 *al.*, 2000; CEHR, 2000; Yuan *et al.*, 2002), denitrifying (Benckriser & Ottow, 1982), sulfate-  
65 reducing (Chauret *et al.*, 1996) and methanogenic conditions (Angelidaki *et al.*, 2000; Wang *et al.*,  
66 2000; Gavala *et al.*, 2003; Chang *et al.*, 2005). DBP is generally described as one of the most  
67 readily biodegradable phthalic acid esters because of the shortness of the alkyl branching chain  
68 (CEHR, 2000; Yuan *et al.*, 2002; Gavala *et al.*, 2003). Anaerobic conditions remain however less  
69 favourable to DBP degradation than aerobic conditions where biodegradation rates are up to ten  
70 fold higher (Staples *et al.*, 1997; Yuan *et al.*, 2002). Strong inhibition of methanogenesis was  
71 reported after addition of  $200 \text{ mg L}^{-1}$  of DBP in organic waste-treating reactors (Angelidaki *et al.*,  
72 2000; Gavala *et al.*, 2003). Furthermore, Staples *et al.* (1997) proposed an anaerobic PAE  
73 biodegradation pathway based on theoretical considerations. This pathway suggests a first  
74 hydrolytic step of one ester bond to form a monoester phthalate and a corresponding alcohol. The  
75 second ester bond is hydrolysed and lead to the formation of phthalic acid. Then, the anaerobic

76 mineralization of the phthalic acid by syntrophic methanogenic consortia is possible via the  
77 benzoate degradation pathway (Kleerebezem *et al.*, 1999; Qiu *et al.*, 2004). Although several DBP-  
78 degrading bacteria have been isolated under aerobic conditions (CEHR, 2000), little is known about  
79 the microorganisms involved in DBP biodegradation under anaerobic conditions. Despite the  
80 evidence of efficient DBP biodegradation under methanogenic conditions (O'Connor *et al.*, 1989;  
81 Ejlertsson *et al.*, 1996; Angelidaki *et al.*, 2000; Gavala *et al.*, 2003), no highly enriched culture or  
82 pure cultures have been obtained and characterized in the literature, likely because of the complex  
83 syntrophic relationships occurring in anaerobic reactors.

84 The aim of this work is to characterize microbial dynamics in enrichment cultures degrading DBP  
85 under methanogenic conditions. An adapted anaerobic ecosystem was enriched in DBP degraders,  
86 and microbial dynamics were monitored by molecular methods.

87

## 88 **[Material and methods]**

89 **Source of methanogenic inoculum.** The methanogenic ecosystem used to inoculate the  
90 enrichment reactors was sampled from a full-scale anaerobic sludge digester at Lynetten wastewater  
91 treatment plant (Denmark) treating household and industrial wastewaters from the Copenhagen  
92 area. The plant treated a number of upstream sources constantly contaminated by phthalic acid  
93 esters. The anaerobic digester had been fed with a mixture of primary and secondary sewage  
94 sludge, and operated at 35°C. After sampling, residual biodegradable organic carbon was depleted  
95 by storing the inoculum for 15 days at 37°C.

96 **Chemicals and preparation of the medium.** All chemicals were of analytical grade (>98%). The  
97 Pentane and Diethyl-ether solvents as well as the Di-n-Butyl Phthalate solutions were provided by  
98 Sigma Aldrich (Saint-Louis, USA). Borosilicate glassware and experimental apparatus were treated  
99 overnight at 200°C to remove trace contaminants. Basal anaerobic (BA) medium was prepared  
100 according to Angelidaki *et al.* (2000). The BA medium was supplemented with 2 g L<sup>-1</sup> of yeast

101 extract, flushed with N<sub>2</sub>:CO<sub>2</sub> (80:20, v/v) for 10 min, and autoclaved (120°C, 30 min.). Vitamins  
102 were added under sterile conditions, according to Angelidaki *et al.* (2000). Before feeding the  
103 reactors, 200 mL of BA medium were freshly amended with 2 mL of DBP solutions at 0 g L<sup>-1</sup>, 1 g  
104 L<sup>-1</sup> or 20 g L<sup>-1</sup> in Pentane : Diethyl ether (15:85, v/v). The final concentrations of DBP in BA  
105 medium were of 0, 10 or 200 mg L<sup>-1</sup>, respectively.

106 **Experimental enrichment procedure.** Five enrichment reactors were operated: Three were  
107 performed with DBP at 0, 10 and 200 mg.L<sup>-1</sup>, called **Blank**, **R10**, and **R200**, respectively. Two  
108 sterile control reactors were performed with 1.4% (w/v) sodium azide, 2 % (w/v) formaldehyde and  
109 DBP at 10 and 200 mg.L<sup>-1</sup>, called **CTRL10** and **CTRL200**, respectively. The enrichments were  
110 carried out in 250 mL serum bottles sealed with Teflon coated rubber stoppers and aluminium caps.  
111 Inoculation corresponded to 200 mL of anaerobic digested sludge flushed with a mixture of N<sub>2</sub>:CO<sub>2</sub>  
112 (80:20, v/v) for 10 min. Enrichments were carried out for 100 days under semi-continuous  
113 conditions with daily manual feeding corresponding to an average hydraulic retention time of 20  
114 days. A two step feeding procedure was performed: It involved sampling of 10 mL reactor content  
115 followed, by the addition of 10 mL BA medium supplemented with DBP at 0, 10 or 200 mg.L<sup>-1</sup>,  
116 respectively. No DBP was initially added to the inoculum. The sampling procedure was carried  
117 out under vigorous agitation to obtain homogeneous outlet sample from the reactor. Reactors were  
118 operated in a temperature controlled room (37°C), under continuous magnetic stirring. Biogas  
119 production was daily measured, and biogas composition of the headspace was weekly analysed.

120 **Analytical procedure.** Gas components (methane, carbon-dioxide and nitrogen) analysis and  
121 volatile fatty acid (VFA) analysis were performed by GC-TCD and GC-FID respectively (Sorensen  
122 *et al.*, 1991). Total Solids (TS) and Volatile Solids (VS) were analysed in triplicates according to  
123 the standard methods for examination of wastewater (APHA, 1995). Phthalic acid isomers and  
124 benzoic acid concentrations were quantified by high pressure liquid chromatography coupled to UV  
125 detection, as described elsewhere (Kleerebezem *et al.*, 1999).

126 **DBP analysis.** 1 mL of sample was diluted in 9 mL of ultrapure water pH 12, and added to 2 mL  
127 of extraction solvent {Pentane : Diethyl ether (15:85 v/v) with 6.6 mg L<sup>-1</sup> Fluoranthene-d10  
128 (Cambridge Isotope Laboratories, Andover, MA) as internal extraction standard}. Extraction was  
129 performed in 15 mL Pyrex tubes capped with a Teflon lined stopper. The tubes were shaken at  
130 room temperature in a tube rotator for 24 hours at 170 rpm (Struers, Gerhardt, Germany). The  
131 extract was then centrifuged (1500 g, 15 min.) and 0.5 mL of supernatant was added to 0.5 mL of  
132 GC injection standard {1 mg L<sup>-1</sup> Phenanthrene-d10 (Cambridge Isotope Laboratories, Andover,  
133 MA) in Pentane : Diethyl ether (15:85 v/v)}. DBP concentrations in the extract were quantified by  
134 gas chromatography (Agilent Technologies 6890N) coupled to mass spectrometry (Agilent  
135 Technologies 5973N) (Christensen *et al.*, 2004).

136 **Single Strand Conformation Polymorphism (SSCP) procedure.** The procedure of DNA  
137 fingerprinting of environmental communities by single strand conformation polymorphism was  
138 performed according to Delbes *et al.* (2000), except the following: An aliquot of 2 mL of sludge  
139 sample was first centrifuged (6000 g, 10 min.), and the pellet was resuspended in 2 mL of 4 M  
140 guanidine thiocyanate-tris HCl pH 7.5 0.1 M and 600 µL of 10 % (w/v) N-Lauroyl-Sarcosine.  
141 Extraction and purification of bacterial genomic DNA was performed with a QIAAmp DNA stool  
142 Mini Kit (Quiagen, Hilden, Germany). The V3 region of the bacterial Small SubUnit rDNA was  
143 amplified by PCR with the primers EF330-FUR500 (Table 1). Because of the low amount of  
144 *Archaea* in the enrichment cultures, the whole Small SubUnit rDNA of this group was first  
145 amplified by PCR using the primers AF333-UR1492 (Table 1). The V3 region of the Small  
146 SubUnit rDNA of the *Archaea* was then amplified with the primers AF333-FUR500. The PCR  
147 products were analysed by SSCP by addition of a size standard (Genescan-400 Rox; Applied  
148 Biosystems), electrophoresis, and computing correction (Genescan software, Applied Biosystems),  
149 according to Delbes *et al* (2000).

150 [TABLE 1]

151

152 **Construction of 16S rDNA clone library and phylogenetic analysis.** The V3 region of the total  
153 16S rDNA was amplified with the primers AF333-UR500 for *Archaea* and EF330-UR500 for  
154 *Bacteria* (Table1). The PCR products were then cloned according to the TOPO TA cloning kit  
155 recommendations (Invitrogen). The clones were then selected to identify individual peaks of SSCP  
156 profiles and sequenced (Delbes *et al.*, 2000). An equal portion of rRNA gene (*E.coli* position 326  
157 to 450) was used for the sequence analysis. Sequences were submitted to Genbank for preliminary  
158 analysis. The NCBI Blast Software was used to identify putative close phylogenetic relatives.  
159 Sequences were aligned to their nearest neighbour with the automated alignment tool of the ARB  
160 software package, and manually checked. The sequences have been submitted to the Genbank  
161 database under the accession numbers EF380210 to EF380215.

162 **Fluorescent *In Situ* Hybridization (FISH) procedure and microscopy observations.** The  
163 method of Hugenholtz *et al.* (2001) was used for fixation and *in situ* hybridization of the samples.  
164 In this study, several 16S rDNA probes were designed and tested for their specificity in targeting  
165 the microorganisms identified as potentially involved in DBP degradation (Table 2). The probes  
166 were optimised with a hybridisation temperature of 46°C, and wash temperature of 48°C, using  
167 blanks as negative controls (Hugenholtz *et al.*, 2001). There was no response to non-target  
168 microbes at any formamide concentration, and strongest emission was found at 0 % and 20 %  
169 formamide (v/v). The slides were examined using a Zeiss LSM 510 confocal laser scanning  
170 microscope (CLSM) with an upright Axioplan 2 microscope and ApoChromat 63/1.4 aperture.  
171 Appropriate excitation lasers and emission filters were used for indocarbocyanine (CY3) and  
172 fluorescein (FITC) labels. In general, the target bacterial cells were labelled using CY3, all bacteria  
173 in FITC, and *Archaea* in CY3.

174 [TABLE 2]

175



176 **[Results]**

177 **Methanogenic activity of the enrichment cultures.** All active cultures showed significant  
178 methanogenic activity during the 110 days experiments (Table 3). The sterile control reactors  
179 produced no gas. The biologically active reactors contained approximately 60 % of methane in the  
180 biogas. No significant inhibition of methanogenesis by DBP at 10 mg L<sup>-1</sup> was observed compared  
181 to the blank containing no DBP (t value of a t-test = 1.34 < 4.3 at 95 % confidence). In contrast,  
182 total biogas production was significantly lower in reactor R200 than in the blank, indicating an  
183 inhibitory effect of DBP at 200 mg L<sup>-1</sup> (value of a t-test = 11.1 > 4.3 at 95 % confidence).  
184 Furthermore, no VFA accumulation was detected in R200 (< 5 mM), suggesting that degradation of  
185 VFA was not specifically inhibited, and that inhibition was rather affecting initial biodegradation  
186 steps. No phthalic acid or benzoic acid accumulation was observed in the biological reactors.  
187 In contrast, because of dilution in the reactors, the concentration of total biomass (VS) constantly  
188 decreased in the sterile control reactors over enrichment time from 7.2 ± 0.6 g L<sup>-1</sup> to 1.51 ± 0.04 g  
189 L<sup>-1</sup> (CTRL10) and 1.34 ± 0.03 g L<sup>-1</sup> (CTRL200) at steady state. The VS content of these reactors  
190 corresponded to the remaining yeast extract and DBP in reactor outlet. The higher VS amounts in  
191 the controls were likely due to a lack of biological hydrolytic activity on the remaining solids. The  
192 lowest VS value was found in the blank (0.80 ± 0.02 g L<sup>-1</sup>). The methanogenic activity in the blank  
193 resulted from degradation of residual particulate substrate, yeast extract, or from autotrophic decay.  
194 Final VS contents in the DBP-degrading biological reactors, i.e. R10 and R200, were similar with  
195 an average value of 0.89 ± 0.02 g L<sup>-1</sup> (F value of ANOVA-test = 2.4 < 7.71 at 95% confidence).  
196 Since no DBP degradation was observed in previous enrichment attempts in absence of yeast  
197 extract, the addition of yeast extract could not be avoided (data not shown).

198 **[TABLE 3]**

199

200 **DBP biodegradation.** DBP concentrations in biological reactors, as well as DBP removal  
201 efficiencies are presented in Figure 1. Both R10 and R200 reached similar final effluent DBP  
202 concentrations of  $1.01 \pm 0.07 \text{ mg L}^{-1}$  and  $1.3 \pm 0.65 \text{ mg L}^{-1}$ , respectively. The theoretical DBP  
203 concentration was based on a mass balance model, indicating DBP accumulation without  
204 degradation (Fig.1). In the sterile reactors, DBP losses, as compared to the accumulation model,  
205 occurred during the first 40 days to reach a maximum of 65 % at  $10 \text{ mg L}^{-1}$  DBP. Since residual  
206 biogas production was also observed (approx.  $3 \text{ mL week}^{-1}$ ), DBP removal was attributed to  
207 incomplete sterility of the reactor. Thenceforth, the sodium azide concentration was increased from  
208 7 to  $14 \text{ g L}^{-1}$  at day 41 (still 2 % (w/v) formaldehyde). The biogas production then stopped and the  
209 DBP level increased towards the theoretical concentration. At the final points, both sterile control  
210 reactors had approximately 20 % DBP losses ( $21.8 \pm 7.5 \%$  at  $10 \text{ mg L}^{-1}$ , and  $21.4 \pm 7.8 \%$  at  $200 \text{ mg}$   
211  $\text{L}^{-1}$ ), attributed to abiotic removal. Considering that DBP is a volatile compound, the highest abiotic  
212 loss was probably due to volatilization rather than sorption to the glass parts of the reactor but this  
213 needs further investigations. No DBP was detected in the blank, indicating no external  
214 contamination. High and constant DBP removal rates were observed in the biological reactors fed  
215 with DBP at  $10 \text{ mg L}^{-1}$  ( $89.7 \pm 0.8 \%$ ) and at  $200 \text{ mg L}^{-1}$  ( $99.3 \pm 0.3 \%$ ) (Fig. 1). In addition,  
216 reactor steady-state was defined as the stable period of time where DBP removal variations were  
217 lower than 5 % around the final average value. It was observed that sterile reactors were not as  
218 stable as the biological reactors, likely because of the higher DBP concentrations causing spatial  
219 heterogeneity. The reactor at  $10 \text{ mg L}^{-1}$  reached a DBP-removal steady state after 70 days of  
220 enrichment, while DBP removal stability was earlier achieved at  $200 \text{ mg L}^{-1}$  (16 days). Based on a  
221 mass balance kinetic model, estimated removal rates at steady state were assessed in R10 and R200  
222 at  $0.46 \pm 0.01 \text{ mg}_{\text{DBPdeg}} (\text{L day})^{-1}$  and  $9.97 \pm 0.1 \text{ mg}_{\text{DBPdeg}} (\text{L day})^{-1}$ , respectively. According to VS  
223 contents in R10 and R200, specific DBP degradation rates were assessed to be  $0.52 \pm 0.02 \text{ mg}_{\text{DBPdeg}}$   
224  $(\text{g}_{\text{VS}} \text{ day})^{-1}$  and  $11.1 \pm 0.35 \text{ mg}_{\text{DBPdeg}} (\text{g}_{\text{VS}} \text{ day})^{-1}$ , respectively.

225 [FIGURE 1]

226

227 **Dynamics of SSCP microbial profiles in the enrichment cultures.** Both bacterial and archaeal  
228 communities were characterized in the biological reactors. Sterile control reactors did not contain  
229 enough DNA material to perform suitable SSCP analysis without introducing unspecific PCR  
230 amplification. Figure 2 compares SSCP profiles of the inoculum and after 100 days of enrichment  
231 for blank, R10, and R200. Area of the SSCP peaks was representative of the abundance of the  
232 associated 16S rDNA sequence. For *Archaea*, the blank maintained its dominant peak (Arc3, 76 %  
233 of peak area) with several sub-dominant species (Arc1, Arc2 and Arc4, 5 %, 8 %, and 7 % of peak  
234 area respectively) (Fig.2). At 10 mg L<sup>-1</sup> of DBP, the population shifted to a bipolar dominance of  
235 Arc3 and Arc4 with approximately equal relative abundance (43 % and 37 % respectively). At 200  
236 mg L<sup>-1</sup> of DBP, species Arc4 was highly dominant (83 %) whereas Arc3 was only found at trace  
237 levels (approx. 6%). Comparatively, *Bacteria* presented more complex profiles (Fig. 2).  
238 Nonetheless, a lower number of SSCP peaks (< 20) were observed in the final *Bacteria* profiles  
239 compared to the inoculum. Owing to the use of synthetic medium in the inlet, the enrichment  
240 procedure led to simplify the total bacterial community in the reactors. In the blank reactor, the  
241 species Bac3 was dominant with approx. 29 % of relative abundance. The abundance of the other  
242 sub-dominant peaks was lower than 5 %. At 10 mg L<sup>-1</sup> DBP, additional dominant peaks appeared,  
243 and especially, Bac1 was found at the same level as Bac3 in the final enrichment culture (17 % and  
244 19 % respectively of relative abundance). At 200 mg L<sup>-1</sup> DBP, the SSCP profile was highly  
245 simplified with only four main peaks: Bac1, Bac2, Bac3, and Bac4 with respectively 23 %, 5 %, 15  
246 %, and 11 % of relative abundance.

247 [FIGURE 2]

248 Microbial SSCP profiles dynamics were evaluated over the entire enrichment procedure and are  
249 presented in Figure 3. Although the archaeal community was phylogenetically stable in the blank  
250 reactor, a progressive shift from Arc3 to Arc4 was observed slowly (after 10 weeks) at 10 mg L<sup>-1</sup>,

251 and more rapidly (within 2 weeks) at 200 mg L<sup>-1</sup> of DBP. Concerning the bacterial community, the  
252 increasing dominance of species Bac1 to Bac4, as well as the disappearance of a group of  
253 intermediary peaks located between Bac1 and Bac2 occurred (Fig.3). In the blank, Bac3 species  
254 preferentially developed and was mainly dominant after 100 days of experimentation. At 10 mg L<sup>-1</sup>  
255 of DBP, Bac3 also developed but the main dominance in the final enrichment culture was supported  
256 by Bac1. The Bac1 dominance occurred after approximately 8 to 9 weeks of enrichment (Fig. 3).  
257 At 200 mg L<sup>-1</sup> of DBP, Bac1 developed more rapidly to dominance. In both reactors, the emergence  
258 of Bac1 occurred simultaneously with the achievement of a stationary phase in term of DBP  
259 removal. As well, the system was considered as phylogenetically stable with regard to no major  
260 changes of the microbial relative abundance, after 8 to 9 weeks at 10 mg L<sup>-1</sup>, and only after 2 weeks  
261 at 200 mg L<sup>-1</sup>.

262 [FIGURE 3]

263

264 **Phylogenetic identification.** The phylogenetic trees representing the affiliation of individual SSCP  
265 clones are shown in Figure 4. Arc1 and Arc2 were not identified because of their low abundance in  
266 the final enrichment cultures. Although a significant shift of the archaeal population was observed  
267 by addition of 10 and 200 mg L<sup>-1</sup> of DBP, both Arc3 and Arc4 belong to the *Methanosaeta* genus.  
268 The divergence between 16S rDNA fragments of the two microorganisms was of 3.9 %. Although  
269 these two clones were phylogenetically close, Arc4 was even closer to *Methanosaeta concilii* strain  
270 GP6 than Arc3 (2.5 % and 4.7 % divergence, respectively).

271 The dominant clones within *Bacteria* phylum in the final DBP enrichments were also  
272 phylogenetically similar (Fig.4). The most dominant species -Bac1- was closely related to  
273 *Soehngenia saccharolytica*. In contrast, the main dominant species Bac3 found in the blank and in  
274 the enrichment cultures was related to the genus *Bacteroides*. Bac2 and Bac4 peaks corresponded  
275 to an identical species belonging to the phylum *Bacteroidetes*, and probably to the order  
276 *Bacteroidales*. Additionally, Bac2 and Bac4 SSCP peaks corresponded to two isomers of the same

277 16S rDNA fragment belonging to one species. A similar artefact of double peaks corresponding to  
278 one 16S rDNA sequence was previously observed under the same analytical conditions (Delbes *et*  
279 *al.*, 2000).

280 [FIGURE 4]

281

282 ***In situ* characterization of the enrichment cultures by FISH.** The probes SOE01-432 and  
283 BCT01-409 were designed to specifically target *Soehngenia saccharolytica* (Bac1), and  
284 *Bacteroides* sp. (Bac2-Bac4). The FISH images are shown in Figure 5. In all samples, microbial  
285 material was flocculant (10-100 µm). No microbes in the blank or inoculum were detected  
286 responding to the probes SOE01-432 or BCT01-409, but organisms responding to both SOE01-432  
287 and BCT01-409 were observed in both enrichment cultures. Organisms responding to SOE01-432  
288 (Bac1) were abundantly present throughout flocs, as well as *Archaea* responding to ARC-915  
289 (approximately 10% of total microorganisms in the enrichment cultures). Based on visual  
290 estimation, Bac1 represented 70 % of the total *Bacteria* community in the flocs, which is  
291 significantly higher than in the SSCP profiles representing the distribution of flocculating and non-  
292 flocculating bacteria (approximately 25 %). The remainder of bacteria (EUB338/338+), as  
293 observed by FISH appeared to be Bac2-Bac4 (BCT01-409) with even, but non structural  
294 distribution throughout the flocs.

295 [FIGURE 5]

296

297 [Discussion]

298 **DBP biodegradation under strict anaerobic conditions.** Although DBP biodegradation has  
299 previously been observed in anaerobic environments (Angelidaki *et al.*, 2000; Wang *et al.*, 2000;  
300 Gavala *et al.*, 2003; Chang *et al.*, 2005), the present study reports for the first time, the possibility to  
301 enrich microbial cultures with highly efficient DBP biodegradation ability under strict anaerobic  
302 conditions. Working with volatile and hydrophobic organic compounds requires the consideration

303 of the potential abiotic losses caused by the experimental setup, such as volatilization or adsorption  
304 on experimental glassware. In the present study, abiotic disappearance of DBP was estimated at 20  
305 % in both sterile controls. The significant differences between controls and biological reactors  
306 indicated effective biological degradation of DBP. Mass balance did not provide evidence of full  
307 mineralization of DBP, since yeast extract was necessary to be added and mineralization could not  
308 be separately evaluated.

309 Nevertheless, primary attack of DBP was clearly shown in both biological reactors. At both 10 mg  
310 L<sup>-1</sup> and 200 mg L<sup>-1</sup> inlet concentrations, the effluent DBP concentrations were of approximately 1.1  
311 mg L<sup>-1</sup>. This value likely corresponds to a threshold concentration where DBP biodegradation was  
312 limited by bioavailability.

313 Because the DBP solvents (pentane and diethyl ether) were added at the same concentration in all  
314 experiments, inhibition due to these specifically would have been the same. Considering this,  
315 specific and significant impact of DBP was observed at 200 mg L<sup>-1</sup> compared to lower  
316 concentrations. Previous study reported methanogenesis inhibition at a similar level (Angelidaki *et*  
317 *al.*, 2000). In contrast, O'Connor *et al.* (1989) reported no toxic effect on methanogenic activity for  
318 concentration above 300 mg L<sup>-1</sup>. This suggests that the nature and the composition of the microbial  
319 community impact on its own sensitivity to phthalic acid ester inhibition. In our enrichment  
320 cultures, since the DBP concentrations were relatively low in the enrichment reactors, direct  
321 inhibition is unlikely. It is more likely that selection of DBP degrading microorganisms and the  
322 resulting specialization of the bacterial population caused a decrease in overall methanogenic  
323 activity. Since no by-product accumulation was observed (VFA or possible aromatic  
324 intermediates), this decrease of activity was due to a decrease in hydrolytic activity rather than  
325 methanogenic activity.

326 **Microbial dynamics and phylogenetic affiliation of the microorganisms involved in DBP**  
327 **biodegradation.** In all biological cultures, a decrease in diversity was observed by SSCP

328 throughout the enrichment procedure. This was mainly due to the wash out of non-growing  
329 microorganisms, such as in the blank. The simplicity of the SSCP profiles showed that only few  
330 microbial species from the inoculum were able to grow on yeast extract under the applied operating  
331 conditions. Moreover, the application of DBP selection pressure in inlet favoured the specialization  
332 of the DBP-degrading consortium by improving the growth of DBP-degrading microorganisms.  
333 Because of the low in-reactor DBP concentration in the enrichment cultures, DBP growth inhibition  
334 did probably not occur. This is consistent with the VS contents in the biological reactors, which  
335 were significantly higher than in the blank without DBP. Phylogenetic stability of the consortium  
336 was observed at steady state, i.e. no major changes in the DBP-degrading microbial population.  
337 Furthermore, the emergence of the final bacterial profile was very slow at 10 mg L<sup>-1</sup> and faster at  
338 200 mg L<sup>-1</sup>. The early profile of microbial dynamics at 200 mg L<sup>-1</sup> corresponded to the final  
339 profiles at lower concentration (10 mg L<sup>-1</sup>), suggesting that selection of microorganisms followed  
340 similar steps over time, and that higher DBP concentrations speed up selection of the specific  
341 degrading microbial consortium. Additionally, the emergence of the final microbial profiles  
342 correlated well with the time to reach a stationary phase for DBP removal. All these results are  
343 consistent with direct involvement of the microbial consortium in DBP biodegradation pathway.  
344 Interestingly, the experimental setup of the enrichment procedure only influenced the dynamics of  
345 the microbial communities that finally tend to a similar consortium whatever the DBP selection  
346 pressure.

347 In this study, a population shift within *Archaea* kingdom occurred. This suggested a rapid  
348 adaptation of the methanogens to DBP (2 weeks at 200 mg L<sup>-1</sup> DBP). The selected methanogen  
349 (Arc4) likely had lower sensitivity to DBP compared to the original species (Arc3) found in the  
350 blank. Co-dominance of both species at 10 mg L<sup>-1</sup> indicated that low DBP concentrations slightly  
351 favour Arc4 emergence, which outcompete Arc3. These results support the involvement of Arc4 as  
352 a partner of DBP degraders at high DBP concentration. Surprisingly, although the abundance shift

353 was clear according to the increasing DBP concentrations, the phylogenetic shift was only very  
354 limited, with Arc3 and Arc4 both being members of the genus *Methanosaeta*. It was therefore  
355 concluded that the final DBP-degrading methanogenic consortium was highly specific to their local  
356 environment, either linked to acetate affinity since both *Methanosaeta* species carried the same  
357 function, or due to physicochemical properties (e.g., surface properties) suitable for DBP  
358 degradation by other microbes necessarily involved in the process.

359 Nevertheless, the presence of *Methanosaeta* sp. in anaerobic enrichment culture was unsurprising.  
360 Leclerc *et al.* (2004) previously reported that *Methanosaeta* sp. represented more than 75 % in  
361 abundance of the archaeal species amongst 44 different anaerobic digesters, and was found in 84 %  
362 of the anaerobic reactors. Although physiological properties would rather favour the  
363 implementation of fast growing hydrogenotrophs (Leclerc *et al.*, 2004), the hydrophobic properties  
364 and the high affinity for acetate as substrate favour mainly the implementation of *Methanosaeta* sp.  
365 in flocs and granules in anaerobic reactors (Grotenhuis *et al.*, 1991; Schmidt & Ahring, 1996;  
366 Sekiguchi *et al.*, 1999; Leclerc *et al.*, 2004). Moreover, *Methanosaeta* sp. outcompete other fast-  
367 growing acetate users, such as *Methanosarcina* sp., at low acetate concentrations (Conklin *et al.*,  
368 2006). Considering that phthalic acid esters are hydrophobic compounds and most of the potential  
369 DBP-degraders concentrated within flocs in the enrichment cultures as observed by FISH, the  
370 presence of *Methanosaeta* sp. moreover likely favoured local hydrophobic environment within  
371 flocs. Although the two identified archaeal species found in blank and enrichments were closely  
372 related, Arc4 was even closer to several microbes (AF229777- AF229778- AF229774) previously  
373 found in methanogenic consortium degrading terephthalate, an isomer of the probable intermediate  
374 orthophthalate in DBP degradation pathway (Wu *et al.*, 2001).

375 In addition, among the two bacteria –Bac1 and Bac2/Bac4 - identified as DBP-degrading  
376 candidates, Bac1 was phylogenetically affiliated to *Soehngenia saccharolytica*, an anaerobic  
377 benzaldehyde degrader (Parshina *et al.*, 2003). The emergence of *Soehngenia saccharolytica* as



378 dominant bacteria in the enrichment cultures correlated well with the stationary phase for DBP  
379 removal in both biological reactors. This is consistent with direct involvement of Bac1 in DBP  
380 biodegradation pathway. One strain of *Soehngenia saccharolytica* was previously reported to  
381 require yeast extract for growth coupled to detoxification by dismutation of benzaldehyde to  
382 benzoate and benzylalcohol (strain BOR), but was not reported to perform aromatic ring fission  
383 (Parshina *et al.*, 2000). No intermediate was detected in our enrichment cultures suggesting that  
384 complete mineralization occurred. This implied the occurrence of primary attack likely performed  
385 by Bac1 affiliated to *Soehngenia saccharolytica*, followed by a ring fission and further oxidation  
386 steps carried by other emerging bacteria (e.g. Bac2/Bac4). According to Staples *et al.* (1997),  
387 phthalic acid, a probable intermediate, is a central intermediate in the biological degradation under  
388 methanogenic conditions of phthalate esters and is then converted to CH<sub>4</sub> (Kleerebezem *et al.*,  
389 1999).

390 The second emerging group (Bac2/Bac4) belonged to the genus *Bacteroides*, commonly found in  
391 anaerobic environments. Because of this sub-dominant group appeared later over the enrichment  
392 procedure, it was concluded that Bac2/Bac4 probably corresponded to bacteria growing on by-  
393 products, especially aromatic rings. Chen *et al.* (2004) reported that clones related to *Bacteroides*  
394 sp. may be involved at a lower extent in terephthalate degradation under thermophilic conditions.  
395 Additionally, molecular tools may present biases, especially with regards to the retrieved sequences  
396 that are only representative of dominant species, as well as the limited specificity of PCR primers  
397 (Delbes *et al.*, 2000; Leclerc *et al.*, 2004). Nevertheless, the FISH observations presented here,  
398 using probes developed from SSCP sequence information, confirmed the SSCP results. At last, the  
399 expression of the degradation function by sub-dominant species, as previously discussed by Delbes  
400 *et al.* (2000), was unlikely here because of the low complexity of the SSCP profiles after enrichment  
401 and the high specialization of the degrading consortium.

402

403 **Enrichment cultures of anaerobic phthalate ester degraders.** Development of enrichment  
404 cultures under strict anaerobic conditions is subject to scientific and technical constraints that were  
405 addressed in this study. First, the PAEs are only found at trace levels in the environment, and the  
406 selection and adaptation of an efficient PAE-degrading ecosystem is very time-consuming  
407 (Kleerebezem *et al.*, 1999; Hayes *et al.*, 1999; Qiu *et al.*, 2004). Edwards & Grbic-Galic (1994)  
408 showed that *ex-situ* adaptation of anaerobic ecosystems to single aromatic compounds, such as  
409 toluene and o-xylene, required more than 100 and 200 days of adaptation in lab systems,  
410 respectively. The levels of exposure (Yuan *et al.*, 2002) as well as the period of contamination  
411 (Hayes *et al.*, 1999) affect the capability of the anaerobic microbial consortium to degrade aromatic  
412 compounds. This issue was addressed in the current study by a preliminary screening of several  
413 potential inocula. In particular, the ability of the anaerobic ecosystem to degrade DBP was not  
414 widely distributed and only long-term naturally contaminated sludge exhibited a substantial  
415 potential for DBP degradation.

416 Second, it is commonly assumed that biodegradation of phthalic acids require syntrophic microbial  
417 populations to occur under methanogenic conditions (Kleerebezem *et al.*, 1999; Qiu *et al.*, 2004).  
418 Obtaining highly enriched cultures depends on the ability to maintain an active syntrophic  
419 consortium of oxidising bacteria, and hydrogen utilising methanogenic *Archaea* throughout the  
420 enrichment procedure. In our study, less than 4 months were necessary in semi-continuous reactors  
421 to select highly enriched cultures by applying strong selection pressure – both dilution rates, and  
422 high loading rates. In contrast, Kleerebezem *et al.* (1999) reported that stable enrichment cultures  
423 on phthalates were obtained after a period of more than 1 year and through numerous transfers into  
424 fresh medium. Qiu *et al.* (2004) reported that over 2 years of enrichment were required to establish  
425 phthalate-degrading enrichment cultures. Such usual enrichment method consisting in successive  
426 transfers into fresh medium is therefore time consuming since microbial growth rates of anaerobic  
427 cultures are low, within a range from 0.08 to 0.25 day<sup>-1</sup> with phthalic acids (Kleerebezem *et al.*,

428 1999; Qiu *et al.*, 2004), and 0.1 day<sup>-1</sup> with other aromatic compounds (Edwards & Grbic-Galic,  
429 1994). The use of a semi-continuous, rather than transfer system therefore favoured more rapid  
430 selection. Nevertheless, the application of higher dilution rates under methanogenic conditions was  
431 restricted by the presence of slow-growing methanogens and DBP-degrading bacteria in the  
432 degradative consortium.

433 At last, stability of the enrichment culture was reached with stable DBP degradation rates near the  
434 end of the test. Therefore, at this dilution rate, cell growth matched washout and decay of non  
435 degrading microorganisms, and 20 days hydraulic retention time was highly suitable to maintain the  
436 ability to degrade DBP. In contrast, Kleerebezem *et al.* (1999) reported that phthalate-enriched  
437 cultures were unstable at low-rates or when less than 20% of cultures were transferred. Qiu *et al.*  
438 (2004) reported similar observations with the possibility of losing the ability to grow on pure  
439 phthalate.

440

#### 441 **[Acknowledgments]**

442 The work was supported by the 6<sup>th</sup> Framework Program – Intra European Fellowship – MEIF-CT-  
443 2003-500956-Xenomic project and the 5<sup>th</sup> EU framework program project (QLK5-CT-2002-01138-  
444 BIOWASTE “Bioprocessing of sewage sludge for safe disposal on agricultural land”). Pr. Jean-  
445 Jacques Godon, Olivier Zemb and Valérie Bru from the INRA-Narbonne (FR), as well as Hector  
446 Garcia from the DTU (DK) are especially thanked for their collaboration and their technical  
447 support.

448

#### 449 **[References]**

450 Angelidaki I, Mogensen AS & Ahring BK (2000) Degradation of organic contaminants found in organic waste.  
451 *Biodegradation* **11**: 377-383.  
452 APHA (1995) *Standard methods for the examination of water and wastewater*. American Public Health Association,  
453 Washington DC.

454 Benckriser G & Ottow JCG (1982) Metabolism of the plasticizer di-n-butylphthalate by *Pseudomonas*  
455 *pseudoalcaligenes* under anaerobic conditions, with nitrate as the only electron acceptor. *Appl Environ Microbiol* **44**:  
456 576-578.

457 CEHR, Center for the Evaluation of risks to Human Reproduction (2000) *National Toxicology Program – CEHR*  
458 *monograph on the potential human reproductive and developmental effects of di-n-butyl phthalate (DBP)*. [Online]  
459 <http://cerhr.niehs.nih.gov>

460 Chang BV, Liao CS & Yuan SY (2005) Anaerobic degradation of diethyl phthalate, di-n-butyl phthalate, and di-(2-  
461 ethylhexyl) phthalate from river sediment in Taiwan. *Chemosphere* **58**: 1601-1607.

462 Chauret C, Iniss WE & Mayfield CI (1996) Biotransformation at 10°C of di-n-butyl phthalate in subsurface  
463 microcosms. *Ground water* **34**: 791-794.

464 Chen CL, Macarie H, Ramirez I, Olmos A, Ong SL, Monroy O & Liu WT (2004) Microbial community structure in a  
465 thermophilic anaerobic hybrid reactor degrading terephthalate. *Microbiol* **150**: 3429–3440.

466 Christensen N, Batstone DJ, He Z, Angelidaki I & Schmidt JE (2004) Removal of polycyclic aromatic hydrocarbons  
467 (PAHs) from sewage sludge by anaerobic degradation. *Water Sci Technol* **50** (9): 237-244.

468 Conklin A, Stensel HD & J Ferguson (2006) Growth kinetics and competition between *Methanosarcina* and  
469 *Methanosaeta* in mesophilic anaerobic digestion. *Water Environ Res* **78**:486-496.

470 Daims H, Bruhl A, Amann R, Schleifer KH & Wagner M (1999) The domain-specific probe EUB338 is insufficient for  
471 the detection of all bacteria: development and evaluation of a more comprehensive probe set. *Syst Appl Microbiol* **22**:  
472 434-444.

473 Delbes C, Moletta R & Godon JJ (2000) Monitoring of activity dynamics of an anaerobic digester bacterial community  
474 using 16S rRNA polymerase chain reaction-single-strand conformation polymorphism analysis. *Environ Microbiol* **2**:  
475 506-515.

476 ECPI, European Council for Plasticizers and Intermediates (2004) [Online.] <http://www.ecpi.org/>

477 Edwards EA & Grbic-Galic D (1994) Anaerobic degradation of toluene and o-xylene by a methanogenic consortium.  
478 *Appl Environ Microbiol* **60**: 313-322.

479 Ejlertsson J, Meyerson U & Svensson BH (1996) Anaerobic degradation of phthalic acid esters during digestion of  
480 municipal solid waste under landfilling conditions. *Biodegradation* **7**: 345-352.

481 Gavala HN, Alatrisme-Mondragon F, Iranpour R & Ahring BK (2003) Biodegradation of phthalate esters during the  
482 mesophilic anaerobic digestion of sludge. *Chemosphere* **52**: 673-682.

483 Grotenhuis JT, Smit M, Plugge CM, Xu YS, Van Lammeren AA, Stams AJ & Zehnder AJ (1991) Bacteriological  
484 composition and structure of granular sludge adapted to different substrates. *Appl Environ Microbiol* **57**: 1942-1949.

485 Hayes LA, Nevin KP & Lovley DR (1999) Role of prior exposure on anaerobic degradation of naphthalene and  
486 phenanthrene in marine harbor sediments. *Org. Geochem.* **30**: 937-945.

487 Hu XY, Wen B & Shan XQ (2003) Survey of phthalate pollution in arable soils in China. *J Environ Monit* **5**: 649-653.

488 Hugenholtz P, Tyson GW & Blackall L (2001) Design and evaluation of 16S rRNA-targeted oligonucleotide probes for  
489 Fluorescence *In Situ* Hybridization. *Methods in Molecular Biology*, Vol. 176 (Lieberman BA, ed.), pp. 29-41. Humana  
490 Press Inc, Totowa, NJ.

491 Jarosova (2006) Phthalic Acid Esters (PAEs) in the food chain. *Czech J Food Sci.* **5**: 223-231.

492 Kim SH, Kim SS, Kwon O, Sohn KH, Kwack SJ, Choi YW, Han SY, Lee MK & Park KL (2002) Effects of dibutyl  
493 phthalate and monobutyl phthalate on cytotoxicity and differentiation in cultured rat embryonic limb bud cells. *J*  
494 *Toxicol Environ Health* **5-6**: 461-472.

495 Kleerebezem R, Hulshoff Pol LW & Lettinga G (1999) Anaerobic degradation of phthalate isomers by methanogenic  
496 consortia. *Appl Environ Microbiol* **65**: 1152-1160.

497 Leclerc M, Delgenes JP & Godon JJ (2004) Diversity of the archaeal community in 44 anaerobic digesters as  
498 determined by Single Strand Conformation Polymorphism analysis and 16S rDNA sequencing. *Environ Microbiol* **6**:  
499 809-819.

500 Mougin C, Dappozze F, Brault A, Malosse C, Schmidt JE, Amellal-Nassr N & Patureau D (2006) Phthalic acid and  
501 benzo[a]pyrene in soil-plant-water systems amended with contaminated sewage sludge. *Environ Chem Lett* **4**: 201-206.

502 O'Connor OA, Rivera MD & Young LY (1989) Toxicity and biodegradation of phthalic acid esters under  
503 methanogenic conditions. *Environ Toxicol Chem* **8**: 569-576.

504 Parshina SN, Kleerebezem R, Van Kempen E, Nozhevnikova AN, Lettinga G & Stams AJM (2000) Benzaldehyde  
505 conversion by two anaerobic bacteria isolated from an upflow anaerobic sludge bed reactor. *Proc Biochem* **36**: 423-429.

506 Parshina SN, Kleerebezem R, Sanz JL, Lettinga G, Nozhevnikova AN, Kostrikina NA, Lysenko AM & Stams AJM  
507 (2003) *Soehngenia saccharolytica* gen. nov., sp. nov. and *Clostridium amygdalinum* sp. nov., two novel anaerobic,  
508 benzaldehyde-converting bacteria. *Int J Syst Evol Microbiol* **53**: 1791-1799.

509 Patureau D, Laforie M, Lichtfouse E, Caria G, Denaix L & Schmidt JE (2007) Fate of organic pollutants after sewage  
510 sludge spreading on agricultural soils: a 30-years field-scale recording. *Water Practice & Technol* **2** (1) / doi  
511 10.2166/wpt.2007.008.

512 Qiu YL, Sekiguchi Y, Imachi H, Kamagata Y, Tseng IC, Cheng SS, Ohashi A & Harada H (2004) Identification and  
513 isolation of anaerobic, syntrophic phthalate isomer-degrading microbes from methanogenic sludges treating wastewater  
514 from terephthalate manufacturing. *Appl Environ Microbiol* **70**: 1617-1626.

515 Schmidt JE & Ahring BK (1996) Granular sludge formation in upflow anaerobic sludge blanket (UASB) reactors.  
516 *Biotech Bioeng* **49**: 229-246.

517 Sekiguchi Y, Kamagata Y, Nakamura K, Ohashi A & Harada H (1999) Fluorescence *in situ* hybridization using 16S  
518 rRNA-targeted oligonucleotides reveals localization of methanogens and selected uncultured bacteria in mesophilic and  
519 thermophilic sludge granules. *Appl Environ Microbiol* **65**: 1280-1288.

520 Sorensen AH, Winther-Nielsen M & Ahring BK (1991) Kinetics of lactate, acetate and propionate in unadapted and  
521 lactate-adapted thermophilic, anaerobic sewage sludge: the influence of sludge adaptation for start-up of thermophilic  
522 UASB-reactors. *Appl Microbiol Biotechnol* **34**: 823-827.

523 Stahl DA & Amann R (1991) Development and application of nucleic acid probes. *Nucleic acid techniques in bacterial*  
524 *systematics*. (Stackebrandt E & Goodfellow M, eds), pp. 205-248. Academic Press, Chichester, UK.

525 Staples CA, Peterson DR, Parkerton TF & Adams WJ (1997) The environmental fate of phthalate esters: a literature  
526 review. *Chemosphere* **35**: 667-749.

527 Wang J, Chen L, Shi H & Qian Y (2000) Microbial degradation of phthalic acid esters under anaerobic digestion of  
528 sludge. *Chemosphere* **41**: 1245-1248.

529 Wu JH, Liu WT, Tseng IC & Cheng SS (2001) Characterization of microbial consortia in a terephthalate-degrading  
530 anaerobic granular sludge system. *Microbiol* **147**: 373-382.

531 Yin R, Lin XG, Wang SG & Zhang HY (2003) Effect of DBP/DEHP in vegetable planted soil on the quality of  
532 capsicum fruit. *Chemosphere* **50**: 801-805.

533 Yuan SY, Liu C, Liao CS & Chang BV (2002) Occurrence and microbial degradation of phthalate esters in Taiwan  
534 river sediments. *Chemosphere* **49**: 1295-1299.

535

537 **Table 1** Primer sequences used for PCR amplification of the total or partial 16S Small SubUnit rRNA genes  
 538

Name	Specificity	<i>E.coli</i> position	Sequence (5' to 3')
UR500	Universal <sup>a</sup>	500	TTA CCG CGG CTG CTG GCA G
FUR500	Universal <sup>a,b</sup>	500	6-FAM- TTA CCG CGG CTG CTG GCA G
UR1492	Universal <sup>a</sup>	1492	GNT ACC TTG TTA CGA CTT
AF3	<i>Archaea</i>	3	ATT CYG GTT GAT CCY GSC RG
AF333	<i>Archaea</i>	333	TCC AGG CCC TAC GGG G
EF330	<i>Bacteria</i>	330	ACG GTC CAG ACT CCT ACG GG

539 <sup>a</sup> all prokaryotes including *Bacteria* and *Archaea*

540 <sup>b</sup> 6-FAM= 6-carboxyfluorescein, terminal DNA fluorescent label

541

542

543  
544  
545

**Table 2** Fluorescent labelled oligonucleotides used for Fluorescent *In Situ* Hybridization probing

Name	Target group	Formamide (%)	Non-group hits NCBI database	Probe sequence (5' to 3')	<i>E.coli</i> position	Reference
SOE01-432	<i>Soehngenia saccharolytica</i>	20	1 <sup>a</sup>	GTCATTATCTCCCCTAGGACAGAGC	432	This study
BCT01-409	Uncultured <i>Bacteroides</i> sp.	20	0	CAACCCCTTAGGGCCGCCTTC	409	This study
EUB-338 <sup>b</sup>	<i>Bacteria</i> – most	20	0	GCTGCCTCCCGTAGGAGT	338	(Stahl & Ammann, 1991)
EUB-338 <sup>+b</sup>	<i>Bacteria</i> -remaining	20	0	GCWGCCACCCGTAGGTGT	338	(Daims <i>et al.</i> , 1999)
ARC-915	<i>Archaea</i>	20	0	GTGCTCCCCCGCCAATTCCT	915	(Stahl & Ammann, 1991)

546  
547  
548  
549

<sup>a</sup> *Catonia barnesae* (AB38361)

<sup>b</sup> EUB-338/EUB-338+ were used simultaneously to target all *Bacteria* (EUB-Mix).



550  
551

**Table 3** Methanogenic performances in the biological and control cultures, after 100 days of enrichment

Bioreactor Name	DBP concentration in inlet (mg L <sup>-1</sup> )	Methane content in the biogas (%) <sup>a</sup>	Average biogas production rate (mL week <sup>-1</sup> ) <sup>a</sup>	VS content (g L <sup>-1</sup> ) <sup>a</sup>
Blank	0	61.3 ± 1.6	41.7 ± 1.5	0.80 ± 0.02
R10	10	62.3 ± 0.6	38.0 ± 3.5	0.88 ± 0.01
CTRL10	10 (sterile control)	n.d	∅	1.51 ± 0.04
R200	200	60.7 ± 1.1	13.7 ± 3.2	0.90 ± 0.02
CTRL200	200 (sterile control)	n.d	∅	1.34 ± 0.03

552  
553  
554  
555  
556

<sup>a</sup> Indicated errors corresponded to 95% confidence intervals of triplicate analyses at steady state  
∅ not measurable production (< 0.5 mL.week<sup>-1</sup>)  
n.d. not determined

557

558 **[Figure Legends]**

559 **Fig. 1.** Theoretical and measured DBP concentrations in the sterile controls (CTRL10 and CTRL200) and  
560 biological reactors (R10 and R200). DBP removal corresponds to the ratio between the theoretical curve and  
561 the measured values. Plain lines indicate biological reactors, while stripped lines correspond to sterile  
562 controls. Error bars of the concentration values represent the standard deviation of triplicate analyses. Error  
563 bars of the DBP removal curves represent 95 % confidence interval of the calculated value.




564 **Fig. 2.** SSCP profiles of the archaeal (a) and bacterial (b) communities in the sludge inoculum and after  
565 100 days of enrichment at 0, 10 and 200 mg L<sup>-1</sup> of DBP. Species of interest are marked by an arrow and  
566 numbered.

567 **Fig. 3.** Three dimensional representations of the archaeal (a) and bacterial (b) SSCP profiles over  
568 enrichment time, and according to DBP concentrations. The SSCP peaks of interest are marked by an arrow  
569 and numbered. (A.U.=Arbitrary Unit)


570 **Fig. 4.** Phylogenetic trees of 16S rDNA fragment of the dominant archaeal (A) and bacterial (B) species  
571 in the blank and enrichment cultures. The trees were generated by using neighbour joining distance method  
572 in ARB software, with distant microorganisms as roots. Numbers at the nodes indicate the bootstrap values  
573 above 50 %, for 1000 bootstrap calculations. The scale bar represents the number of substitutions per  
574 nucleotide. The phylogenetic divergence correspond to the comparison of partial sequences from *E.coli*  
575 nucleotide 330 to nt 500 (*Bacteria*) and nt 333 to nt 500 (*Archaea*). The sequence roots correspond to  
576 *Methanosarcina mazei* (**AF028691**) and *Escherichia coli* (**AJ567617**) for *Archaea* and *Bacteria* trees,  
577 respectively. The sequences obtained from the present study are indicated in bold.

578 **Fig. 5.** Fluorescent In Situ Hybridization microscopic observations of identified DBP degraders in the  
579 200 mg L<sup>-1</sup> DBP enrichment culture. A: sample hybridized with EUB338-FITC, ARC915-CY3, and the  
580 specific probe SOE01-432-CY3 to give target coloured cells yellow, other *Bacteria*, green, and *Archaea* red.  
581 B: sample hybridized with EUB338-FITC, ARC915-CY3, and the specific probe BCT01-409-CY3 to give  
582 target coloured cells yellow/red, other *Bacteria*, green, and *Archaea* red. In Figure 5B, most of the target  
583 cells appear red instead of yellow, due to the very strong response by other cells (presumptively Bac1) to the  
584 EUB338 probe. Bar indicates 10 µm.

585





586  
587 Fig. 1.  
588





589  
 590  
 591

**Fig. 2.**



592  
59  
59  
59  
59  
59  
59  
60  
60  
602  
603  
604  
605  
606  
607  
608  
609  
610  
611  
612  
613  
614  
615  
616  
617  
618  
619  
620  
621  
622  
623  
624



0 mg/L



10 mg/L




200 mg/L

(a)


**Fig. 3.**

625



626  
627  
628  
629


A.



630  
631  
632  
633  
634  
635  
636  
637

B.

Fig. 4.



638

639

640

641 **Fig. 5.**

642

643