



HAL
open science

Central domain of a potyvirus VPg is involved in the interaction with the host translation initiation factor eIF4E and the viral protein HcPro

Genevieve Roudet, Thierry Michon, Jocelyne J. Walter, Thierry Delaunay, Elise Redondo, Olivier O. Le Gall

► To cite this version:

Genevieve Roudet, Thierry Michon, Jocelyne J. Walter, Thierry Delaunay, Elise Redondo, et al.. Central domain of a potyvirus VPg is involved in the interaction with the host translation initiation factor eIF4E and the viral protein HcPro. *Journal of General Virology*, 2007, 88 (3), pp.1029-1033. 10.1099/vir.0.82501-0 . hal-02669156

HAL Id: hal-02669156

<https://hal.inrae.fr/hal-02669156>

Submitted on 31 May 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - ShareAlike 4.0 International License

Short Communication

Central domain of a potyvirus VPg is involved in the interaction with the host translation initiation factor eIF4E and the viral protein HcPro

G. Roudet-Tavert, T. Michon, J. Walter, T. Delaunay, E. Redondot and O. Le Gall

Correspondence

Olivier Le Gall
olegall@bordeaux.inra.fr

IPV, UMR GDPP INRA-Bordeaux 2, IBVM, BP 81, F-33883 Villenave d'Ornon Cedex, France

Received 26 August 2006
Accepted 27 November 2006

Using recombinant proteins produced in bacteria or in infected plants, interactions between the VPg and HcPro of *Lettuce mosaic potyvirus* (LMV) and between LMV VPg and the lettuce translation initiation factor 4E, the cap-binding protein (eIF4E), were demonstrated *in vitro*. Interaction with eIF4E and HcPro both involved the same VPg central domain. The structure of this domain in the VPg context was predicted to include an amphiphilic α -helix, with the amino acids related to biological functions in various potyviruses exposed at the hydrophilic side.

Viral infection relies on the availability of host factors to interact with virus proteins. Limited information is available on the proteins involved in the cycle of plant viruses, but interactions between host and virus proteins have been documented in recent years. In the genus *Potyvirus*, the interaction between VPg (a small virus-encoded protein bound to the 5' end of the virus RNA genome) and the host cap-binding protein, eIF4E (Wittmann *et al.*, 1997), has received more attention since eIF4E defects were associated with an inability of several hosts to support potyvirus infection (Gao *et al.*, 2004; Lellis *et al.*, 2002; Nicaise *et al.*, 2003; Ruffel *et al.*, 2002). This suggested a biological significance for the eIF4E–VPg interaction, which was confirmed recently (Kang *et al.*, 2005). In lettuce, eIF4E is encoded by *mol1*, a recessive resistance gene against *Lettuce mosaic potyvirus* (LMV) (Nicaise *et al.*, 2003). VPg also interacts with other proteins encoded by the genome of the host (Dunoyer *et al.*, 2004; Schaad *et al.*, 1997; Yambao *et al.*, 2003) or of the virus (Guo *et al.*, 2001; Hong *et al.*, 1995), including the multifunctional protein HcPro (Yambao *et al.*, 2003). In several potyviruses, variations in the VPg central domain are associated with resistance-breaking (Borgström & Johansen, 2001; Keller *et al.*, 1998; Masuta *et al.*, 1999; Moury *et al.*, 2004; Nicolas *et al.*, 1997; Rajamäki & Valkonen, 1999; Schaad *et al.*, 1997). In this work, we investigated the interactions of LMV VPg, in particular its central domain, with lettuce eIF4E and LMV HcPro.

Recombinant lettuce (*Lactuca sativa*) eIF4E was expressed in bacteria from pDest17 (Invitrogen). Recombinant LMV VPg and HcPro were expressed in bacteria from pTrcHis (Invitrogen). These three proteins were purified from

bacterial extracts as poly(His) fusions. In some experiments, His-tagged LMV HcPro purified from infected plants (Plisson *et al.*, 2003) was used, yielding essentially similar results to those shown below (data not shown). Protein purity was controlled by SDS-PAGE (not shown). Rabbit polyclonal antibodies were obtained against VPg and eIF4E. Mouse monoclonal antibodies 1H5 and 21M were raised against VPg and VPg-CD (a synthetic peptide corresponding to the VPg central domain with a non-viral C-terminal extension, VFSDIGLVQDAFGKERLHAAAHAY), respectively. These mAbs were assessed by Western blotting and ELISA using infected lettuce or recombinant VPg (not shown); in addition, the reactivity of 21M for the VPg central domain was confirmed, as increasing concentrations of VPg-CD resulted in loss of ELISA signal (not shown). A monoclonal antibody against HcPro, 3C5, has been described previously (Roudet-Tavert *et al.*, 2002). An ELISA-derived (Clark & Adams, 1977; Léonard *et al.*, 2000) interaction assay was used. The surface was coated with eIF4E or HcPro [$4 \mu\text{g (ml carbonate buffer)}^{-1}$ overnight at 4°C] and saturated with fetal bovine serum (FBS; 30 min at room temperature) before incubation with VPg [$8 \mu\text{g (ml PBS/0.2 \% Tween)}^{-1}$, 0.2 % FBS for 1 h at 4°C]. Interactions were revealed with 1H5 [$5 \mu\text{g (ml PBS/0.2 \% Tween)}^{-1}$ for 2 h at 37°C], followed by antibodies conjugated to alkaline phosphatase. In reverse experiments, the wells were coated with VPg ($4 \mu\text{g ml}^{-1}$) and the interaction with eIF4E or HcPro ($8 \mu\text{g ml}^{-1}$) was revealed similarly using anti-eIF4E or 3C5.

A specific interaction was observed between VPg and eIF4E, as well as between VPg and HcPro (Fig. 1a). An N-terminally truncated version of HcPro (Plisson *et al.*, 2003) also interacted with VPg (not shown), indicating that the first 99 N-terminal residues of HcPro are dispensable for the interaction with VPg. The interaction

†Present address: BIOGEMMA, 24 avenue des Landais, F-63170 Aubière, France.




Fig. 1. Immunochemical characterization of the interaction between VPg and eIF4E or HcPro. (a) VPg interacts with eIF4E and HcPro. The bars represent the A_{405} values observed in ELISA with the coating and ligand proteins and the antibody indicated below each bar. BSA was used as a control protein. mAbs against LMV VPg (1H5) and HcPro (3C5) and polyclonal antibodies raised against recombinant lettuce eIF4E purified from *Escherichia coli* after overexpression were used. The means \pm SD of three replicates obtained in a typical experiment is shown. (b) Anti-VPg mAb inhibits binding of VPg to eIF4E and HcPro. Wells were coated with an anti-VPg polyclonal antibody (pAb), eIF4E or HcPro, and the interaction with VPg was assayed, with (open bars) or without (filled bars) a preliminary incubation with the anti-VPg-CD mAb 21M. Interactions were revealed using the anti-VPg mAb 1H5. Means \pm SD of A_{405} values (mean of four replicates) of a typical experiment are shown. (c) eIF4E and HcPro compete for interaction with VPg. Wells were coated with HcPro ($5 \mu\text{g ml}^{-1}$), and interaction with VPg ($1 \mu\text{g ml}^{-1}$) was assayed after a preliminary incubation with serial concentrations of eIF4E (open bars) or HcPro (filled bars) at the concentrations indicated ($\mu\text{g ml}^{-1}$). Interactions were revealed using the anti-VPg Mab 1H5. Means \pm SD of A_{405} values (mean of three replicates) of a typical experiment are shown.




Fig. 2. Spectroscopic characterization of the interaction between eIF4E and VPg. eIF4E tryptophan fluorescence was measured following the addition of VPg (○) or VPg-CD (●) (0.05–5 μM) was added to a 2.5 μM eIF4E solution in 1 ml of 20 mM HEPES/KOH (pH 7.6), 25 mM KCl, 1 mM DTT, 10 % glycerol. The mixture was incubated (5 min at 25 °C) until a steady-state fluorescence at 342 nm was reached upon excitation at 280 nm. AU, arbitrary units. Inset: variation of eIF4E fluorescence as a function of increasing concentrations of VPg (○) or VPg-CD (●).

between eIF4E and VPg was confirmed by a spectrofluorometric approach, whereby binding of a ligand results in quenching of the fluorescence of eIF4E tryptophans. A decrease in eIF4E tryptophan fluorescence was observed in the presence of increasing concentrations of VPg (Fig. 2) but not of BSA used as a control (not shown). The data suggested a stoichiometry of one molecule bound per eIF4E and a dissociation constant in the same range as that of the cap-eIF4E interaction (Khan *et al.*, 2006; Michon *et al.*, 2006).

In *Turnip mosaic virus* (TuMV) (Léonard *et al.*, 2000), although not in *Tobacco etch virus* (TEV) (Schaad *et al.*, 2000), amino acid changes in the VPg central region that impair eIF4E interaction are lethal, suggesting a direct interaction with eIF4E. The VPg central domain is also involved in HcPro binding (Yambao *et al.*, 2003). Therefore, the involvement of the LMV VPg central domain in both HC-Pro and eIF4E interactions was assayed *in vitro* as described above. The anti-VPg-CD mAb 21M did not reveal an interaction between VPg and eIF4E (not shown), suggesting that the VPg central domain was not accessible to 21M once the VPg-eIF4E interaction had taken place

and hence that this domain of VPg was involved directly in eIF4E interaction.

To test the hypothesis that eIF4E, HcPro and 21M bound in the same area on VPg, VPg was pre-incubated with 21M (1 : 3, w/w, for 2 h at 37 °C) and assayed for eIF4E binding. Such pre-treatment of VPg significantly diminished binding to eIF4E, as well as to HcPro (Fig. 1b). When an anti-VPg polyclonal antibody was used for immunocapture of VPg in a control experiment, no reduction in signal was observed after pre-treatment, indicating that 21M did not interfere with recognition by 1H5, the mAb used to reveal the interaction. Taken together, this confirmed that 21M, when bound to the VPg central domain, interferes with further interaction with eIF4E or HcPro, and thus demonstrated that the central domain must remain available for specific interactions with eIF4E and HcPro to occur.

In order to confirm the localization of the interaction domain on VPg, the wells were coated with the synthetic peptide VPg-CD (5 µg ml⁻¹) and incubated with eIF4E. The results were revealed using a mAb (2B12) directed against the N-terminal tag of the recombinant eIF4E. In a typical experiment, A_{405} values averaged 0.264 ± 0.040 for VPg-CD compared with 0.028 ± 0.006 for BSA, confirming a specific interaction. The interaction between eIF4E and VPg-CD was confirmed by spectrofluorometry, as above (Fig. 2), with the same dissociation constant as for VPg. Similarly, in a binding assay revealed using the anti-HcPro mAb 3C5, VPg-CD interacted with HcPro ($A_{405} = 0.420 \pm 0.190$ vs 0.028 ± 0.025).

VPg-CD overlaps the region involved in the interaction between eIF4E and TuMV VPg (Léonard *et al.*, 2000) and between HcPro and VPg of *Clover yellow vein virus* (Yambao *et al.*, 2003). The present work therefore provides the first evidence that, for a single given potyvirus, HcPro and eIF4E bind to the same central domain of VPg (residues 89–105). This clearly suggests the possibility that they could compete for binding to VPg. In order to assess this possibility, we conducted an ELISA-based competition assay in which the effect of lettuce eIF4E on LMV HcPro–VPg binding was studied. In such experiments, the wells were coated with HcPro (5 µg ml⁻¹) followed by VPg (1 µg ml⁻¹) pre-incubated with either eIF4E or HcPro (1 h at 4 °C). In both cases, inhibition of HcPro–VPg interaction could be observed (Fig. 1c), confirming that HcPro and eIF4E compete for VPg binding.

A three-dimensional model of VPg from *Potato virus Y* has been predicted based on that of an unrelated protein showing a similar distribution of hydrophobic/hydrophilic residues (Płochocka *et al.*, 1996). VPg appeared as an egg-shaped protein, with the domain involved in the interaction exposed at one pole. Although this model was obtained through an unusual approach, it is consistent with an involvement in protein interactions.

To understand better the possible relationships between VPg function and structure, attempts were made to predict

the folding of LMV VPg. Homology-based modelling was unsuccessful due to the lack of homologous proteins of known structure in databases (not shown). To predict the secondary structure of the interaction domain (Fig. 3), ten VPg sequences were aligned and the secondary structure was predicted for each of them using GOR-IV (Garnier *et al.*, 1996). Other algorithms operated online (<http://www.expasy.org> and <http://npsa-pbil.ibcp.fr>) provided essentially similar results (not shown). A hydrophobic (Kyte & Doolittle, 1982) β -sheet was returned for all sequences except those of bymoviruses, between the RNA-bound Tyr-64 (Murphy *et al.*, 1991) and Asp-78, a residue associated with eIF4E-binding in TuMV (Léonard *et al.*, 2000). For all viruses analysed, the downstream portion (residues 90–120) was predicted to be folded in a long amphiphilic α -helix (Fig. 3). The amino acid variations associated with different biological properties (Borgström & Johansen, 2001; Masuta *et al.*, 1999; Moury *et al.*, 2004; Nicolas *et al.*, 1997; Rajamäki & Valkonen, 1999; Schaad *et al.*, 1997), where identified unambiguously, mapped to the hydrophilic – presumably exposed – side of this helix, which is consistent with a role in protein interactions.

Despite obvious discrepancies between this two-dimensional prediction and the available three-dimensional model (Płochocka *et al.*, 1996), both are compatible with a surface exposition of the VPg central domain. Further structural and biochemical studies are needed to clarify the VPg modelling and the roles of the VPg central domain in the success of virus infection through interactions with eIF4E, HcPro and possibly other proteins. The biological implications of the duality of the interaction in which the VPg central domain is engaged with the host protein eIF4E and the virus protein HcPro will also require investigation. For instance, it has been established that changes in the VPg central domain that affect long-distance movement of TEV in tobacco are not necessarily associated with a loss of eIF4E interaction (Schaad *et al.*, 2000). It is possible that this type of phenotype could be related to changes in affinity with HcPro, a viral protein involved in long-distance movement (Cronin *et al.*, 1995).

Acknowledgements

This work was partially supported by EPR Aquitaine (ref. 20.000307004), by the Genoplante programme ('Newvir', Gen56) and by the French National Agency for Research ('Poty4E', ANR-05-Blan-0302-01).

References

- Borgström, B. & Johansen, I. E. (2001). Mutations in *Pea seedborne mosaic virus* genome-linked protein VPg alter pathotype-specific virulence in *Pisum sativum*. *Mol Plant Microbe Interact* **14**, 707–714.
- Clark, M. F. & Adams, A. N. (1977). Characteristics of the microplate method of enzyme-linked-immunosorbent-assay for the detection of plant viruses. *J Gen Virol* **34**, 475–483.
- Cronin, S., Verchot, J., Haldeman-Cahill, R., Schaad, M. C. & Carrington, J. C. (1995). Long-distance movement factor: a transport

- function of the potyvirus helper component proteinase. *Plant Cell* **7**, 549–559.
- Dunoyer, P., Thomas, C., Harrison, S., Revers, F. & Maule, A. (2004). A cysteine-rich plant protein potentiates potyvirus movement through an interaction with the virus genome-linked protein VPg. *J Virol* **78**, 2301–2309.
- Gao, Z., Johansen, E., Eyers, S., Thomas, C. L., Noel Ellis, T. H. & Maule, A. J. (2004). The potyvirus recessive resistance gene, *sbm1*, identifies a novel role for translation initiation factor eIF4E in cell-to-cell trafficking. *Plant J* **40**, 376–385.
- Garnier, J., Gibrat, J. F. & Robson, B. (1996). GOR method for predicting protein secondary structure from amino acid sequence. *Methods Enzymol* **266**, 540–553.
- Guo, D., Rajamäki, M. L., Saarma, M. & Valkonen, J. P. (2001). Towards a protein interaction map of potyviruses: protein interaction matrixes of two potyviruses based on the yeast two-hybrid system. *J Gen Virol* **82**, 935–939.
- Hong, Y., Levay, K., Murphy, J. F., Klein, P. G., Shaw, J. G. & Hunt, A. G. (1995). A potyvirus polymerase interacts with the viral coat protein and VPg in yeast cells. *Virology* **214**, 159–166.
- Kang, B. C., Yeam, I., Frantz, J. D., Murphy, J. F. & Jahn, M. M. (2005). The *pvr1* locus in *Capsicum* encodes a translation initiation factor eIF4E that interacts with *Tobacco etch virus* VPg. *Plant J* **42**, 392–405.
- Keller, K. E., Johansen, I. E., Martin, R. R. & Hampton, R. O. (1998). Potyvirus genome-linked protein (VPg) determines *Pea seed-borne mosaic virus* pathotype-specific virulence in *Pisum sativum*. *Mol Plant Microbe Interact* **11**, 124–130.
- Khan, M. A., Miyoshi, H., Ray, S., Natsuaki, T., Suehiro, N. & Goss, D. J. (2006). Interaction of genome-linked protein (VPg) of turnip mosaic virus with wheat germ translation initiation factors eIFiso4E and eIFiso4F. *J Biol Chem* **281**, 28002–28010.
- Kyte, J. & Doolittle, R. F. (1982). A simple method for displaying the hydropathic character of a protein. *J Mol Biol* **157**, 105–132.
- Lellis, A. D., Kasschau, K. D., Whitham, S. A. & Carrington, J. C. (2002). Loss-of-susceptibility mutants of *Arabidopsis thaliana* reveal an essential role for eIF(iso)4E during potyvirus infection. *Curr Biol* **12**, 1046–1051.
- Léonard, S., Plante, D., Wittmann, S., Daigneault, N., Fortin, M. G. & Laliberté, J. F. (2000). Complex formation between potyvirus VPg and translation eukaryotic initiation factor 4E correlates with virus infectivity. *J Virol* **74**, 7730–7737.
- Masuta, C., Nishimura, M., Morishita, H. & Hataya, T. (1999). A single amino acid change in viral genome-associated protein of *Potato virus Y* correlates with resistance breaking in ‘Virgin A mutant’ tobacco. *Phytopathology* **89**, 118–123.
- Michon, T., Estevez, Y., Walter, J., German-Retana, S. & Le Gall, O. (2006). The potyviral virus genome-linked protein VPg forms a ternary complex with the eukaryotic initiation factors eIF4E and eIF4G and reduces eIF4E affinity for a mRNA cap analogue. *FEBS J* **273**, 1312–1322.
- Moury, B., Morel, C., Johansen, E., Guilbaud, L., Souche, S., Ayme, V., Caranta, C., Palloix, A. & Jacquemond, M. (2004). Mutations in *Potato virus Y* genome-linked protein determine virulence toward recessive resistances in *Capsicum annuum* and *Lycopersicon hirsutum*. *Mol Plant Microbe Interact* **17**, 322–329.
- Murphy, J. F., Rychlik, W., Rhoads, R. E., Hunt, A. G. & Shaw, J. G. (1991). A tyrosine residue in the small nuclear inclusion protein of tobacco vein mottling virus links the VPg to the viral RNA. *J Virol* **65**, 511–513.
- Nicaise, V., German-Retana, S., Sanjuan, R., Dubrana, M. P., Mazier, M., Maisonneuve, B., Candresse, T., Caranta, C. & Le Gall, O. (2003). The eukaryotic translation initiation factor 4E controls lettuce susceptibility to the potyvirus *Lettuce mosaic virus*. *Plant Physiol* **132**, 1272–1282.
- Nicolas, O., Dunnington, S. W., Gotow, L. F., Pirone, T. P. & Hellmann, G. M. (1997). Variations in the VPg protein allow a potyvirus to overcome *va* gene resistance in tobacco. *Virology* **237**, 452–459.
- Plisson, C., Drücker, M., Blanc, S., German-Retana, S., Le Gall, O., Thomas, D. & Bron, P. (2003). Structural characterization of HC-Pro, a plant virus multifunctional protein. *J Biol Chem* **278**, 23753–23761.
- Plochocka, D., Wejnicky, M., Zielenkiewicz, P. & Ostojka-Zagorski, W. (1996). Three-dimensional model of the potyviral genome-linked protein. *Proc Natl Acad Sci U S A* **93**, 12150–12154.
- Rajamäki, M. L. & Valkonen, J. P. (1999). The 6K2 protein and the VPg of *Potato virus A* are determinants of systemic infection in *Nicotiana glauca*. *Mol Plant Microbe Interact* **12**, 1074–1081.
- Roudet-Tavert, G., German-Retana, S., Delaunay, T., Delecote, B., Candresse, T. & Le Gall, O. (2002). Interaction between potyvirus helper component-proteinase and capsid protein in infected plants. *J Gen Virol* **83**, 1765–1770.
- Ruffel, S., Dussault, M. H., Palloix, A., Moury, B., Bendahmane, A., Robaglia, C. & Caranta, C. (2002). A natural recessive resistance gene against *Potato virus Y* in pepper corresponds to the eukaryotic initiation factor 4E (eIF4E). *Plant J* **32**, 1067–1075.
- Schaad, M. C., Anderberg, R. J. & Carrington, J. C. (2000). Strain-specific interaction of the *Tobacco etch virus* NIa protein with the translation initiation factor eIF4E in the yeast two-hybrid system. *Virology* **273**, 300–306.
- Schaad, M. C., Lellis, A. D. & Carrington, J. C. (1997). VPg of tobacco etch potyvirus is a host genotype-specific determinant for long-distance movement. *J Virol* **71**, 8624–8631.
- Thompson, J. D., Gibson, T. J., Plewniak, F., Jeanmougin, F. & Higgins, D. G. (1997). The CLUSTAL_X windows interface: flexible strategies for multiple sequence alignment aided by quality analysis tools. *Nucleic Acids Res* **25**, 4876–4882.
- Wittmann, S., Chatel, H., Fortin, M. G. & Laliberté, J. F. (1997). Interaction of the viral protein genome linked of *Turnip mosaic potyvirus* with the translational eukaryotic initiation factor (iso) 4E of *Arabidopsis thaliana* using the yeast two-hybrid system. *Virology* **234**, 84–92.
- Yambao, M. L., Masuta, C., Nakahara, K. & Uyeda, I. (2003). The central and C-terminal domains of VPg of *Clover yellow vein virus* are important for VPg–HCPro and VPg–VPg interactions. *J Gen Virol* **84**, 2861–2869.