



HAL
open science

A gene-based radiation hybrid map of chicken microchromosome 14: comparison to human and alignment to the assembled chicken sequence

Mireille Morisson, Sophie S. Leroux, Carine Jiguet-Jiglaire, Sirine Assaf, Frederique Pitel, Sandrine Lagarrigue, Suzanne Bardes, Katia K. Feve, Thomas Faraut, Denis Milan, et al.

► To cite this version:

Mireille Morisson, Sophie S. Leroux, Carine Jiguet-Jiglaire, Sirine Assaf, Frederique Pitel, et al.. A gene-based radiation hybrid map of chicken microchromosome 14: comparison to human and alignment to the assembled chicken sequence. *Genetics Selection Evolution*, 2005, 37, pp.229-251. 10.1051/gse:2004046 . hal-02670446

HAL Id: hal-02670446

<https://hal.inrae.fr/hal-02670446>

Submitted on 31 May 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Copyright

A gene-based radiation hybrid map of chicken microchromosome 14: Comparison to human and alignment to the assembled chicken sequence

Mireille MORISSON^{a*}, Sophie LEROUX^a, Carine JIGUET-JIGLAIRE^a,
Sirine ASSAF^b, Frédérique PITEL^a, Sandrine LAGARRIGUE^b,
Suzanne BARDES^a, Katia FEVE^a, Thomas FARAUT^a, Denis MILAN^a,
Alain VIGNAL^a

^a Laboratoire de génétique cellulaire, Institut national de la recherche agronomique,
31326 Castanet-Tolosan, France

^b Laboratoire de génétique animale, Institut national de la recherche agronomique,
École nationale supérieure d'agronomie de Rennes, 35042 Rennes Cedex, France

(Received 3 May 2004; accepted 11 October 2004)

Abstract – We present a gene-based RH map of the chicken microchromosome GGA14, known to have synteny conservations with human chromosomal regions HSA16p13.3 and HSA17p11.2. Microsatellite markers from the genetic map were used to check the validity of the RH map and additional markers were developed from chicken EST data to yield comparative mapping data. A high rate of intra-chromosomal rearrangements was detected by comparison to the assembled human sequence. Finally, the alignment of the RH map to the assembled chicken sequence showed a small number of discordances, most of which involved the same region of the chromosome spanning between 40.5 and 75.9 cR₆₀₀₀ on the RH map.

chicken / comparative mapping / radiation hybrids / microchromosome 14 / intrachromosomal rearrangement

1. INTRODUCTION

The numerous efforts made these last years in the field of chicken genomics [2, 12] come from the importance of this species in agriculture and its great value for research in virology, developmental biology, oncology and immunology [6]. Thus, a large genomic toolset was developed including

* Corresponding author: mmorriso@toulouse.inra.fr

a detailed consensus linkage map of the genome including over two thousand markers [15, 16]. Large collections of chicken expressed sequence tags (EST) were released [1, 4, 37] and bacterial artificial chromosome (BAC) libraries [9, 21] were constituted and used to assemble both local and genome-wide chicken BAC contig maps [11, 29]. BAC contigs are usually used as a platform on which full genome sequences are assembled. They also serve as a bridge between the genome sequence and the linkage map, the essential tool for QTL analysis [30]. In March 2004, a first draft assembly of the chicken genome sequence was released by the Washington University Genome Sequencing Center (WUGSC) and the National Human Genome Research Institute (<http://www.genome.gov/11510730>). Due to a significantly lower rate of interspersed repetitive elements, this draft chicken genome sequence is probably more accurate than the first draft human genome sequences originally published three years ago [18, 38]. Nevertheless, the integration of all chicken genomic resources such as the BAC contigs, the genetic and the radiation hybrid (RH) maps, will be essential for the assembly of the whole genome sequence data to a reliable and more informative resource. Thus, in addition to the BAC contig map, an RH map will provide an independent platform to assist the chicken genome sequence assembly process towards a finished quality sequence. RH mapping is a powerful tool for locating genes since it uses the simple polymerase chain reaction (PCR): contrariwise to genetic markers, RH markers do not require the development of polymorphism. The whole genome radiation hybrid (WGRH) panel we produced [24] has already been used to build radiation maps for chicken microchromosome 15 [20], macrochromosome 4 and 7 [25, 28], and maps of other chromosomes are under construction.

The first comparisons of chicken gene maps with those of humans have revealed an unexpectedly high level of conserved syntenies [8, 14, 27, 32, 33]. However, subsequent and more detailed mapping studies have revealed high levels of intra-chromosomal rearrangements within them [7, 10, 19, 34, 36]. The expectations are therefore, that the number of segments of the conserved gene order will increase with the number of genes mapped in the chicken.

To develop a dense RH map of chicken microchromosome GGA14, we adopted a strategy based on the use of markers from the genetic map to check the validity of the RH map and from EST or genes whose location on GGA14 could be predicted from known data on conserved syntenies. The human/chicken comparative data published in 2000 by Schmid *et al.* [31] showed that two genes (*HBA* and *NTN2*) localised on GGA14 were both located on HSA16p13.3, suggesting that this human chromosome should be used as a base for developing gene-based markers. More recently, the gene *SREBP1*,

orthologous to *SREBF1* localised on HSA17p11.2 in human, was also shown to be located on GGA14 by segregation analysis in the East Lansing reference family [3], providing another source for the development of markers.

The GGA14 RH map obtained was compared to the assembled human sequence to detect the chromosome rearrangements that have occurred in the lineages leading to humans and chickens. Finally, the RH map was compared to the newly available assembled chicken genome sequence in order to detect discordances pointing to potential assembly problems of the sequence.

2. MATERIALS AND METHODS

2.1. Radiation hybrid panel

The production of the RH panel has already been described [24]. Briefly, normal diploid female chicken fibroblasts irradiated at 6000 rads by gamma rays from a Cesium-137 source were fused to the hypoxanthine guanine phosphoribosyl transferase (HPRT)-deficient hamster cell line, Wg3hC12 [13]. The hybrid cells were selected on HAT (hypoxanthine-aminopterin-thymidine) media, tested for marker retention and subjected to large-scale culture for DNA extraction. The final panel was composed of 90 clones with a mean marker retention frequency of 22%.

2.2. Markers from the genetic map

Nine microsatellites (*ADL0118*, *ADL0263*, *LEI0066*, *LEI0098*, *MCW0123*, *MCW0136*, *MCW0225*, *MCW0296* and *ROS0005*) and 2 SSCP (*GCT0903* and *GCT0908*) markers from the two known genetic linkage groups attributed to GGA14 were used. In the course of the whole genome RH mapping work going on in the laboratory, the microsatellite marker *ADL0205* was found linked to GGA14. More information for these 12 markers is available at <http://www.thearkdb.org/browser> or <https://acedb.asg.wur.nl/>.

2.3. EST (expressed sequence tag) and gene markers

All publicly available chicken EST (>420 000) from Genbank and other sources [1, 4, 37] were collected in a local database, after which EST selection and primer design were performed after comparison to the human genome sequence using the Iccare web server [26]. EST markers corresponding to the *COQ7* and *DREVI* genes had been previously developed in our laboratory and

were found to be linked to the GGA14 RH linkage group. Finally, an additional gene marker was developed to map the chicken *SREBP1* gene, orthologous to human *SREBF1* [3] (Tab. I).

2.4. Markers from the sequence assembly

Eight markers (*SEQ0168*, *SEQ0170*, *SEQ0171*, *SEQ0172*, *SEQ0173*, *SEQ0174*, *SEQ0175* and *SEQ0177*) were designed directly from a portion of the GGA14 sequence assembly for which no markers existed on the GGA14 RH map. Primers are given in Table II.

2.5. PCR conditions

PCR were performed in 15 μ L, containing 25 ng hybrid DNA, concentrations of $MgCl_2$ ranging from 1 to 3 mM as determined by test experiments, 0.3 U Taq DNA polymerase (Life technologies: Carlsblad, CA, USA), 1 X buffer (Life technologies), 200 μ M of each dNTP, 0.2 μ M of each primer and 1X loading buffer composed of 350 mM sucrose and 0.2 mM cresol. After denaturation for 10 min at 94 °C, 33 PCR cycles of 30 s at 94 °C, 30 s at the specific annealing temperature, 30 s at 72 °C, and a final elongation step of 10 min at 72 °C were performed. PCR products were analysed using 2% agarose gels and visualised using ethidium bromide staining. Chicken DNA was used as the positive control and Wg3hCl2 DNA and TE (Tris-EDTA) buffer as negative controls. Each marker was genotyped twice and a third genotyping experiment was performed when discrepancies between the first two experiments were found.

2.6. Map construction

All markers were scored as present or absent in each of the hybrids. Ambiguous results were also reported. Pairwise and multipoint data analysis was completed using RH2PT and RHMAXLIK programmes in the RHMAP3.0 software package [5, 22]. We assumed random breakage along the chromosomes and equiprobable retention of fragments. The RH map was constructed in three steps: (1) a two point analysis identified markers linked together with an LOD score greater than 6, thus defining linkage groups; (2) multipoint analyses were done with RHMAXLIK to define a framework map with the markers from the largest linkage group, using a stepwise marker addition strategy and

a LOD threshold of 3. The resulting framework map was further tested by removing one marker at a time and calculating the LOD score for all of its possible positions; (3) a comprehensive map was built by calculating the location of additional markers relative to the framework markers. Finally the map figure was created using MapChart, a software for the graphical presentation of linkage maps and QTL developed by Voorrips and colleagues [39].

2.7. Comparative mapping data – Alignment to the chicken sequence assembly

Data on human gene order were obtained from the Ensembl (<http://www.ensembl.org/>) or NCBI Mapview (<http://www.ncbi.nlm.nih.gov/mapview/>) browsers. Data on the location of chicken genes were obtained by BLASTN or BLAT searches using the Ensembl (<http://www.ensembl.org/>) or Golden Path (<http://genome.ucsc.edu/>) browsers.

3. RESULTS

3.1. Development of EST (expressed sequence tag) markers

Genes located in the regions of HSA16 and HSA17 identified as having a conservation of synteny with GGA14 were used to search for chicken orthologous EST with the Iccare software. Primers were then selected according to the constraints of RH mapping: species-specific amplification of an exonic region. Eighty such chicken EST could thus be used to design PCR primers suitable for RH mapping and 63 (80%) led to successful amplifications.

Markers corresponding to the *COQ7*, *DREVI* and *SREBF1* genes were added to our study (see Materials and Methods). Altogether, 66 EST or gene markers, as shown in Table I, were used to build the GGA14 RH map. However, mapping data concerning a few human genes have evolved since we designed the primers. Hence, *GCT1270* is similar to the open reading frame *Res4-22C* now localised on HSA4 and *UBC* (*GCT1264*) is now on HSA12. Therefore, only 64 of our EST markers were orthologous to genes located on HSA16p13.3 or HSA17p11.2.

3.2. Construction of the GGA14 RH map, marker retention

Genotyping data on the ChickRH6 panel for the 12 genetic and 66 gene markers were used to generate an RH map. After two point analysis at a LOD

Table I. Primer sequences for chicken markers corresponding to human genes and EST. Position on the genome sequence and chromosome information are from the Ensembl (<http://www.ensembl.org/>) or NCBI Mapview (<http://www.ncbi.nlm.nih.gov/mapview/>) genome browsers. Unknown: the genomic sequence exists but could not be assigned to a chromosome in the genome sequence assembly. No hits: the sequence was not found by BLASTN in the genome sequence.

Human				Chicken		
Gene	Genbank ID	Sequence data		Genbank ID	Sequence data	
		Chromosome	Localisation (Mb)		Chromosome	Localisation (Mb)
<i>A2BP1</i>	NM_145893	HSA16	7,383	AL588477	GGA14	13,241
<i>ABAT</i>	NM_000663	HSA16	8,736	BU384127	GGA14	12,243
<i>ABCA3</i>	NM_001089	HSA16	2,267	BI067240	GGA14	1,554
<i>ACCN1</i>	NM_001094	HSA17	31,487	BU281335	GGA27	0,277
<i>AKAP10</i>	NM_007202	HSA17	19,971	BU248455	GGA19	6,355
<i>BLMH</i>	NM_000386	HSA17	28,721	AJ445455	GGA19	5,818
<i>BM045</i>	NM_018459	HSA16	4,880	BI394462	unknown	
<i>CACNA1H</i>	AF223563	HSA16	1,204	BU313977	GGA14	7,320
<i>CGTHBA</i>	NM_012075	HSA16	0,074	AI979784	GGA14	13,944
<i>CLCN7</i>	NM_001287	HSA16	1,435	BI064676	GGA14	14,126
<i>COQ7</i>	NM_016138	HSA16	19,046	BU127240	GGA14	10,660
<i>COX10</i>	NM_001303	HSA17	14,173	BU297164	GGA18	2,459
<i>CREBBP</i>	NM_004380	HSA16	3,777	BU133172	GGA14	14,342
<i>CRYBA1</i>	NM_005208	HSA17	27,719	M15658	GGA19	5,568
<i>DECR2</i>	NM_020664	HSA16	0,392	BG713681	GGA14	2,063
<i>DKFZp434F054</i>	NM_032259	HSA16	0,675	BI392137	GGA14	1,942
<i>DREV1</i>	NM_016025	HSA16	21,577	AJ444698	GGA14	10,917
<i>E4F1</i>	NM_004424	HSA16	2,214	BU134214	GGA14	8,538
<i>ELAC2</i>	NM_018127	HSA17	13,096	BU481814	GGA18	0,988
<i>FLJ14154</i>	NM_024845	HSA16	3,494	BU446205	GGA14	14,176
<i>FLJ20040</i>	NM_018992	HSA16	2,732	AL587387	GGA14	9,409
<i>GIT1</i>	NM_014030	HSA17	28,046	BU365311	No Hits	
<i>GRAP</i>	NM_006613	HSA17	19,125	BU374370	GGA14	7,218
<i>HAGH</i>	NM_005326	HSA16	1,799	AJ396406	GGA14	8,085
<i>HBZ</i>	NG_000006	HSA16	0,220	AL586775	GGA14	13,962
<i>HMOX2</i>	NM_002134	HSA16	4,527	BI394804	GGA14	14,271
<i>HSCARG</i>	NM_020677	HSA16	4,512	BI390022	GGA14	14,573
<i>HSPC055</i>	NM_014153	HSA16	11,811	BU467982	GGA14	3,092
<i>JJAZ1</i>	NM_015355	HSA17	30,410	AJ443766	GGA18	6,405
<i>KIAA0643</i>	AB014543	HSA16	3,551	BG711879	unknown	
<i>LLGL1</i>	NM_004140	HSA17	18,329	BU222034	unknown	

Table I. Continued.

Human				Chicken		
Gene	Genbank ID	Sequence data		Genbank ID	Sequence data	
		Chromosome	Localisation (Mb)		Chromosome	Localisation (Mb)
<i>LOC51061</i>	NM_015914	HSA16	11,739	BU117620	GGA14	3,096
<i>LUC7L</i>	NM_018032	HSA16	0,179	AJ397480	GGA14	14,119
<i>Magmas</i>	AL833954	HSA16	4,371	BI392448	GGA14	14,524
<i>MAP2K3</i>	NM_002756	HSA17	21,364	BU215684	GGA14	6,604
<i>MGC15416</i>	BC001912	HSA16	0,638	BG712373	unknown	
<i>MGC2605</i>	NM_032304	HSA16	0,717	BI394017	GGA14	8,081
<i>MRPS34</i>	NM_023936	HSA16	1,762	AL585124	unknown	
<i>MYH11</i>	NM_002474	HSA16	15,763	X06546	GGA14	9,602
<i>NCOR1</i>	NM_006311	HSA17	16,136	BU143568	GGA19	6,068
<i>NF1</i>	NM_000267	HSA17	29,568	S62087	GGA19	8,557
<i>NLK</i>	NM_016231	HSA17	26,515	AJ443268	GGA19	8,826
<i>NT5M</i>	NM_020201	HSA17	17,407	BI394839	GGA14	6,862
<i>NTHL1</i>	NM_002528	HSA16	2,030	BU249652	GGA14	8,303
<i>NUBP2</i>	NM_012225	HSA16	1,777	BG711301	GGA14	1,889
<i>PDPK1</i>	NM_002613	HSA16	2,528	AL587372	GGA14	9,385
<i>PM5</i>	NM_014287	HSA16	16,293	BU315883	GGA14	9,790
<i>PMP22</i>	NM_000304	HSA17	15,334	BU408850	GGA18	2,175
<i>PPL</i>	NM_002705	HSA16	4,933	BU284851	GGA14	1,838
<i>PRPSAP2</i>	NM_002767	HSA17	18,962	BU389510	GGA14	7,179
<i>Rab11</i>	NM_014700	HSA16	0,416	AL585583	GGA18	6,198
<i>RAB26</i>	NM_014353	HSA16	2,139	BU205196	GGA14	8,454
<i>RES4-22C</i>	AB000461	HSA4	2,659	BU140047	GGA4	82,790
<i>RGS11</i>	NM_003834	HSA16	0,258	BU460973	GGA14	14,083
<i>SDOS</i>	NM_032349	HSA16	4,744	AF095446	GGA14	14,180
<i>SOX8</i>	AK024491	HSA16	0,972	BU451000	GGA14	7,800
<i>SREBF1</i>	NM_004176	HSA17	17,917	AY029224	GGA14	6,985
<i>STUB1</i>	NM_005861	HSA16	0,670	BU449476	unknown	
<i>TRAF4</i>	NM_004295	HSA17	27,217	BU258287	GGA19	5,446
<i>TRAP1</i>	NM_016292	HSA16	3,708	BE139917	GGA14	14,326
<i>UBC</i>	NM_021009	HSA12	123,749	BU409506	GGA15	4,481
<i>UBE2I</i>	AK024172	HSA16	1,289	AJ396786	GGA14	9,438
<i>UBN1</i>	NM_016936	HSA16	4,898	BU390306	GGA14	1,865
<i>ULK2</i>	BC034988	HSA17	19,862	BU204959	GGA19	6,388
<i>USP7</i>	NM_003470	HSA16	8,810	BQ038106	GGA14	12,143
<i>WSB1</i>	NM_015626	HSA17	25,767	BU318529	GGA19	8,635

Table I. Continued.

Human gene	Marker			Product size
	Name	Forward primer (5'-3')	Reverse primer (5'-3')	
<i>A2BP1</i>	<i>GCT1062</i>	GGAAGCTACAGCACCTCC	TGCTCTGGAACAGTGGTTTG	147
<i>ABAT</i>	<i>GCT1348</i>	CAGGAAAATTCTGGGCACAC	GGCCTGAACTCCTCTTTGTG	112
<i>ABCA3</i>	<i>GCT1054</i>	CATTGTCTGCACCTTGAAC	AAGATGATGGATTGCCACAC	160
<i>ACCN1</i>	<i>GCT1344</i>	TCGTCAGACATGGGCTTAGAC	GCATGTGAACCATTTTGACAG	112
<i>AKAP10</i>	<i>GCT1265</i>	TCAAGCCACACATCCTCTTG	TGCCGTAAGGGAGTAGTGAAAC	111
<i>BLMH</i>	<i>GCT1341</i>	AATGCCTTTGTGGAACAGC	ATATCCCACTGGCCTCCATC	101
<i>BM045</i>	<i>GCT1067</i>	GAGCAGAGCAGAGGTCGTTC	TAAAAGGACCCGACCGAGAC	245
<i>CACNA1H</i>	<i>GCT1243</i>	GCTTTTTCTCCAGGCGTC	TCATCGTCAGCTTCTTCGTG	135
<i>CGTHBA</i>	<i>GCT1047</i>	AACAATTTTGCTACAAAGTCAG	AACGTAGGGTGACCAACAAAG	87
<i>CLCN7</i>	<i>GCT1052</i>	GTGGAGGCAGGGAGAGGAC	ACTGGAGAAGGCAGCAACAC	107
<i>COQ7</i>	<i>GCT1469</i>	GCACAGTGGCGGTGGAAG	GCATCAGTCTTTGTACTTTTC	100
<i>COX10</i>	<i>GCT1259</i>	CCCTGCTTGCTGTATGCTTC	GTGTTGGCGATGCTCATTC	157
<i>CREBBP</i>	<i>GCT1271</i>	CACTAAGAGCCATGACCACAAG	GCACAACCCGTTTTCATCTTC	206
<i>CRYBA1</i>	<i>GCT1339</i>	CACAAGGAATCCAAGATCACTG	ATCTTCATGGAGCCCACTTC	128
<i>DECR2</i>	<i>GCT1227</i>	ACATCCGAGTGAACAGCTTG	GACCAGTGTGGTGCCAGTC	200
<i>DKFZp434F054</i>	<i>GCT1050</i>	CGAGCTTTGGCCTCTCTG	CTGCATCCAAGCCCAAC	122
<i>DREV1</i>	<i>GCT1470</i>	GGGCTCAATGTTTGTGTTTTTC	GACCTCTCCATCTCCAGCTC	109
<i>E4F1</i>	<i>GCT1248</i>	AGTCATTGTGGCAGGAGACC	TCATGTGAGCTTCGAGGATG	214
<i>ELAC2</i>	<i>GCT1255</i>	TGGATGAAGATGAATGCTG	AATCCAACCTTCTCAGGAAGTC	103
<i>FLJ14154</i>	<i>GCT1346</i>	CCATCTACCTGCACGTCTCTC	GGTGGGTGTCCCACTATTG	160
<i>FLJ20040</i>	<i>GCT1055</i>	TGCAGTGAGTTGTACAGTTG	GCCAGTTGACACCAGCTTATAG	117
<i>GIT1</i>	<i>GCT1340</i>	CTCAGGCCAACTTCTTCCAC	CGTACACCACCAGCAGCTC	99
<i>GRAP</i>	<i>GCT1260</i>	CAGCACGTCCAGCACTTC	CCTGTAGAAGTCCACCAGCTC	102
<i>HAGH</i>	<i>GCT1066</i>	TCAAAGACAGTCGCACAAGC	CCACTATTGCAGCTTCTCTTC	109
<i>HBZ</i>	<i>GCT1239</i>	CCGAGGACAAGAAGCTCATC	GGTCGAAGTGGGGGAAGTAG	132
<i>HMOX2</i>	<i>GCT1060</i>	CTGGAAGAGGAAATGGATCG	GCTTCTTTCCGGTGAAGCTC	83
<i>HSCARG</i>	<i>GCT1232</i>	AATGTGCCACCACAACC	TGGCCTATGTAAGTCTCTGG	200
<i>HSPC055</i>	<i>GCT1355</i>	AGAAACAGTGGCAGGGTCAC	AAACATCCGGTTGGAAGC	106
<i>JJAZ1</i>	<i>GCT1342</i>	CCAAAGCGTACGAAAGCAAG	ATTTCTTGTGGACGGAGAGG	140
<i>KIAA0643</i>	<i>GCT1057</i>	AGATGGCTATGCGGTGAGAG	TGAATTTGCTGCTGTCTTCC	113
<i>LLGL1</i>	<i>GCT1257</i>	GCTGGGGAAGGAGATCCAG	AGGAGATGAGCATGGAGTGG	155
<i>LOC51061</i>	<i>GCT1354</i>	TGAATACAAAGGCCCAATGAG	TTTTGATCGAGACGAGATGTAGAG	91
<i>LUC7L</i>	<i>GCT1048</i>	ATCCTTCATTGCGGAGTGTG	CAGCACTGATCTTCTCTGTG	80
<i>Magmas</i>	<i>GCT1059</i>	AAAGCCGAGACGTGACTCC	GAGGACACGGCGTTTATTG	83
<i>MAP2K3</i>	<i>GCT1262</i>	CGGGGATGACTCTTAACTGG	AAGGTGTTGGCTTCCAAGG	192
<i>MGC15416</i>	<i>GCT1063</i>	GGCCAAAATTGAAAAGTCTC	TGGGGTGTGTTGTTGTTG	150
<i>MGC2605</i>	<i>GCT1064</i>	GGAAGGAGGATGTGTTCTC	CTCATCCGACCCATACAC	203
<i>MRPS34</i>	<i>GCT1065</i>	GAAGACCCCTTGCCTACTG	CGACCAGGAGTCCAAAGTTC	185
<i>MYH11</i>	<i>GCT1353</i>	TGAAGAGAAGAACCTCCTCCAG	TCAATCTGGCCTCCATCTC	138
<i>NCOR1</i>	<i>GCT1263</i>	GAAAATAATCCACGCAGGAAAG	GAAATCGCTTGTGTTCTC	103

Table I. Continued.

		Marker		Product
Human gene	Name	Forward primer (5'-3')	Reverse primer (5'-3')	size
<i>NF1</i>	<i>GCT1343</i>	GTCATAGCATTCCGCTCCAG	TCCATGATCTCCAGCAAAGC	119
<i>NLK</i>	<i>GCT1337</i>	GGCCAGGGTGAAGAATTAG	TGCAAAGATGCAGCCTACAG	142
<i>NT5M</i>	<i>GCT1261</i>	GCTGGTGGACATGGATGG	AGGGCAATGTAGGGCTTGTC	93
<i>NTHL1</i>	<i>GCT1246</i>	TCTCTGGAGGGAGATAAACTGG	TGCTCAAAGAAGCCAGAATG	146
<i>NUBP2</i>	<i>GCT1053</i>	GGAAGAGCACCATCAGCAC	CATTGTCTGCACCTTGAAC	221
<i>PDPK1</i>	<i>GCT1056</i>	TGCTACCAAAAAGATTAGGCTGTG	TGCTACCAAAAAGATTAGGCTGTG	151
<i>PM5</i>	<i>GCT1352</i>	TGGGAAAGCAGGCTTATTTATG	ACTGTGATGAGAGGGGAAGG	102
<i>PMP22</i>	<i>GCT1258</i>	CAGTCTGTCCAAGCAATGATG	CTTGGTGAGCGTGAAGAGC	90
<i>PPL</i>	<i>GCT1251</i>	CTCCCTCTACAGCCCTACCC	AACATGGAGCTGACGTACCC	212
<i>PRPSAP2</i>	<i>GCT1256</i>	TCAGAAGGAAATACAGGGCTTC	TTCTTGGATGTACTGTAGTAAAAATGG	82
<i>Rab11</i>	<i>GCT1049</i>	GCAGCGGAAGTTGATCTC	CAGCCTCCTTCTGTAATCC	226
<i>RAB26</i>	<i>GCT1240</i>	CGGCCTCAATGTGGAATTAG	GCAGCTACAAAACGGTTGC	177
<i>RES4-22C</i>	<i>GCT1270</i>	TTGTAGCTGTATGCCGTTGG	AGGTCCATTCCTTTGTCAGG	195
<i>RGS11</i>	<i>GCT1247</i>	CAGGTGAAATCATCCATCTGC	TAGCCCAGTAGGTGGTGTGCG	130
<i>SDOS</i>	<i>GCT1061</i>	ACGCAGATAGCGGAGAGAAC	ATTACAAAACCGGGGCATC	218
<i>SOX8</i>	<i>GCT1268</i>	GACCCCAAAACTGACCTG	CGAAGGTCTCCATGTTGTTG	152
<i>SREBF1</i>	<i>EXT002</i>	CAGCCCCACTTTCATCAAGGC	GGCCAGGATGGTCCCTCCAC	249
<i>STUB1</i>	<i>GCT1242</i>	CGCTGTGCTACCTGAAGATG	GGCCTCGTCGTAGTTCTCCA	140
<i>TRAF4</i>	<i>GCT1338</i>	ACATACCGGAGACCTTCCAC	CCTCGTGGGAGATGAACCTTG	123
<i>TRAP1</i>	<i>GCT1058</i>	CCAAGGACATCGGTGAGTG	GCTCAGGCACATAGAAGATGC	135
<i>UBC</i>	<i>GCT1264</i>	AAGACCCTCACTGGCAAGAC	GGACCAGATGGAGAGTGAC	196
<i>UBE2I</i>	<i>GCT1051</i>	GCACAAGCCAAGAAGTTTGC	GTGGCGAATATGGAAAATGG	178
<i>UBN1</i>	<i>GCT1252</i>	GCTGTCGCTCACTCTCTTTG	AGAGCCTCCGCTTTATGTCTC	168
<i>ULK2</i>	<i>GCT1266</i>	AGTGTTCCTCCAGCAGATCG	GGAACCACACAGAGTTGCAG	205
<i>USP7</i>	<i>GCT1349</i>	GATAACAGTGAGCTGCCAACAG	TCGTTGGGGATGGTTTTATC	92
<i>WSB1</i>	<i>GCT1336</i>	CATGGCACAAAAGAATGTTGC	GTCCTGCCCAAATTTGAAAC	201

threshold of 6, a large linkage group containing 60 markers (49 genes or EST, plus 11 genetic markers) was defined. By lowering the LOD threshold down to 4, the microsatellite marker *ADL0118* could be added. A framework map 342 cR₆₀₀₀ long, containing 23 markers and covering the entire chicken chromosome 14 was constructed (Fig. 1). Thirty-eight additional markers linked to GGA14, but whose positions were not supported by a LOD score greater than 3, are indicated on the side of the map to avoid size inflation and to keep track of different possible local orders.

The average retention rate for the markers was 23.7%, in accordance with the first estimates for the panel [24]. The retention frequency along the chromosome, as estimated by plotting the retention of the markers from the framework map against their positions, demonstrates that the higher retention rates were




Figure 1. Alignment of the chicken chromosome 14 genetic and radiation hybrid maps. The genetic maps on the left of the figure are from Groenen *et al.* [15]. For clarity, only the framework markers and additional markers present both on the genetic and radiation hybrid (RH) maps are shown. The RH map is in the middle. Markers positioned on the vertical double bars of the RH map are the framework markers and additional markers are indicated to the right of the map. Microsatellite markers are underlined. The microsatellite marker *ADL0118* in brackets was added to the map by lowering the twopoint lod threshold from 6 to 4. *GCT903* and *GCT908* (boxed) are located on the same microchromosome by FISH [23]. The chart to the right indicates the retention rate of the framework markers. The marker *MCW0123* which was used for selecting the clones for the panel is indicated by an arrow on the retention chart. The suggested centromeric region is indicated by a vertical bar to the right of the RH map.



located around the microsatellite marker *LEI0066*, suggesting that the centromere could be located in this area (Fig. 1). The marker *MCW0123*, which was used for selecting the clones while constructing the panel [24] did not show a particularly high level of retention, when compared to the surrounding markers.

3.3. Colinearity between genetic and RH maps, resolution of the panel

Eleven loci (*ADL0118*, *ADL0263*, *GCT0903*, *GCT0908*, *LEI0066*, *LEI0098*, *MCW0123*, *MCW0136*, *MCW0225*, *MCW0296* and *ROS0005*) were shared between the genetic and the RH maps (the microsatellite marker *ADL0205* is not on the genetic map; *MCW0225* corresponds to *NTN2* on the genetic map).

Comparing both maps indicates a good overall agreement with several improvements in marker ordering. One major change comes from the localisation of the centromere around *LEI0066*, suggesting a reverse orientation for the GGA14 genetic map. Markers *MCW0136*, *ADL0118* and *GCT0903* are mapped with a higher precision on the RH map. Finally, the localisation of *GCT0908* close to *MCW0296* indicates that the genetic linkage group C37 is a part of GGA14.

The part of the RH map between *LEI0066* and *MCW0296* was 259.7 cR₆₀₀₀ long while the genetic distance between these two markers was 72 cM. The ratio between the two maps was thus 3.6 cR₆₀₀₀ to 1 cM.

3.4. Comparison to the assembled GGA14 sequence

When compared, the GGA14 genome sequence assembly and the RH map presented an overall good colinearity (Fig. 2), although two major discrepancies could be detected.

The first concerned a region from positions 1.554 to 2.063 Mb, terminal on the sequence assembly, but found in position 40.5 to 75.9 cR₆₀₀₀ of the RH map, shown in red in Figure 2. This region includes six genes (*DECR2*, *DKFZp434F054*, *NUBP2*, *UBN1*, *PPL* and *ABCA3*) on the sequence map, as well as the microsatellite marker *ADL0205*. On the RH map, it contains additionally *Rab11*, located on the sequence contig of GGA18, and the three markers *STUB1*, *KIAA0643* and *MGC15416*, matching all three with sequence data of unknown location in the chicken genome sequence assembly. The second main discrepancy concerns the region between 14.573 Mb (*HSCARG*) and 20.310 Mb (*MCW0225*) of the sequence assembly, for which no marker could be found on the RH map, despite a high density. To test this region, we developed new markers: *SEQ0168*, *SEQ0170*, *SEQ0171*, *SEQ0172*, *SEQ0173*, *SEQ0174*, *SEQ0175* and *SEQ0177*, directly from the genomic sequence. As a result, marker *SEQ0177* was the only one to be linked to the GGA14 RH map, close to *LEI0066*. Markers *SEQ0172*, *SEQ0173*, *SEQ0174* and *SEQ0175* were linked by RH mapping to markers from GGA3 and markers *SEQ0168*, *SEQ0170* and *SEQ0171* were linked to markers not yet positioned on our RH maps.

Finally, a few improvements over the sequence assembly concerned three markers corresponding to an existing genomic sequence of unknown location. One (*LLGL1*) was at position 239.4 cR₆₀₀₀. The two others (*MRPS34* and *BM045*) at position 0 cR₆₀₀₀, extended the RH map further than the sequence assembly.

4. DISCUSSION

4.1. Development of the EST markers

The first constraint on the choice of primers for RH mapping was to avoid the presence of introns, whose positions in the chicken were predicted on the basis of the human genomic sequence. The second was to design primers in the most divergent regions of the human-chicken sequence alignments so as to avoid cross-amplification with the hamster DNA present in the hybrids. Using Iccare proved to be very efficient, with 80% of the primer pairs designed yielding usable RH mapping data, enabling the mapping of a high number of

Table II. Primer sequences for chicken markers designed from the sequence assembly. Positions on the genome sequence are from the Ensembl (<http://www.ensembl.org/>) genome browser.

Name	Seq. loc. (Mb)	Forward primer (5'-3')	Reverse primer (5'-3')
<i>SEQ0168</i>	14.6	TGGAGAAGAGAAGGCTGTGG	TCCCTCTCACATCCAACCTC
<i>SEQ0170</i>	15.0	TTCAGGCAGCACAAAGTTCAG	AGCATTTCCACCAGATCGAC
<i>SEQ0171</i>	16.0	GTGATTTACCCCGATCTTC	GTAGCTGAGGGGCAGCTATG
<i>SEQ0172</i>	17.0	AAAAGAAGGCAAGCAACACG	AACAGCTCAAACCTGCATGG
<i>SEQ0173</i>	18.0	GTAGAGACGTGGCGCTGAG	AGCAATTTGCAACTGCTCTG
<i>SEQ0174</i>	19.0	ACTGTGCTCACCATTTGCAG	TTTGGGTTCCCTAAAAGTGC
<i>SEQ0175</i>	19.5	ATCCCCACACCTTGTGTTC	TCGAGGAAGATAGGCAGGTG
<i>SEQ0177</i>	20.2	TCATTCCTGGCCCTACTCTG	AGCGTGCTAAAGACCACCTC

EST and genes on the GGA14 RH map. Moreover, the development of EST through the use of Iccare minimises the chances of choosing wrong ortholog genes, since it performs a BLASTN comparison of all chicken EST against the complete set of human Unigene clusters.

4.2. Marker retention and position of the centromere

Preferential retention in RH clones of chromosome fragments from pericentromeric regions of donor cells has been shown in various species, including humans [17,35] and chickens [25]. The retention frequency of markers shown in Figure 1 indicates a drop from 35% down to 15% over the first 200 cR₆₀₀₀ from one end of the map, after which the retention of the markers varies only slightly, with values comprised between 15 and 20%. This data suggests a position for the centromere towards one end of the RH map and is therefore compatible with an acrocentric microchromosome. As a result of our observations and so as to position the centromere conventionally towards the top of the figures, we suggest to reverse the orientation of the genetic map and of the Mb counting on the sequence assembly.

A similar trend, with a drop of 45% down to 15% over a similar distance of 200 cR₆₀₀₀ from the centromere was observed for GGA7, although this chromosome is twice the size of GGA14 [25]. This can explain the higher retention rate observed for microchromosome markers when constructing the panel [24], since they have a higher chance than macrochromosome markers of being close to the centromere.

GGA14 sequence (Mb)

Radiation map (cR₆₀₀₀)




Figure 2. Alignment of the RH map to the assembled sequence of GGA14. The numbering in Mb of the genomic sequence assembly (left) from <http://www.ensembl.org/>, is in inverse order to that of the cR₆₀₀₀ on the RH map (right), as the latter takes into account the suggested position of the centromere. Markers localised on the genetic maps are in bold and loci not yet found on the sequence assembly are underlined on the RH map. Markers placed in a different position on the GGA14 RH map and GGA14 genome sequence assembly are in red. *Rab11* in purple is on the GGA18 in the genome sequence assembly. Markers in the GGA14 genome sequence assembly mapped to other chromosomes by RH are in green (linked to GGA3 markers) or blue (linked to markers not yet assigned to a chromosome on the current RH maps). The dashed lines indicate the relative position of fragments on the two maps.



4.3. Comparison to the genetic map

Fluorescent *in situ* hybridisation (FISH) experiments with the BAC clones P1-8 and P6-V11 from which the genetic SSCP markers *GCT0903* and *GCT0908* are derived, suggested that the small linkage group C37 could be linked to GGA14 [23]. Here we confirm this result by the inclusion of both *GCT0903* and *GCT0908* on the RH map. The reason why these two linkage groups are independent on the genetic map is that *GCT0908* and *COM0079* were only mapped in the Compton population and the nearby marker *MCW0296* only on the Wageningen population. Due to the history of its development using three independent populations, the chicken genetic map still contains a number of small linkage groups, whose chromosome assignment has to be determined. The RH map also enables a greater precision for the mapping of the two markers *MCW0136* and *ADL0118*. The ratio of 3.6 cR₆₀₀₀ to 1 cM was close, although slightly lower, to the previous observation (4cR₆₀₀₀ to 1 cM) for GGA7 [25]. The recombination rate of microchromosomes being higher than that of macrochromosomes, a lower cR₆₀₀₀ to cM ratio was expected for GGA14 than for GGA7.

4.4. Comparison to the genome sequence assembly

Although the genomic sequence assembly for GGA14 covers a total of 20.4 Mb (Fig. 2), a few discrepancies were found with the RH map, one of them suggesting that a portion of 5 Mb from the sequence assembly (between *HSCARG* and *MCW0225*) belongs in fact for a large part to GGA3 and possibly to other chromosomes. As a confirmation, the microsatellite marker

MCW0083, from the GGA3 genetic linkage group was also found at position 19.2 Mb of the GGA14 sequence assembly. This brings the length of the GGA14 sequence down to 15 Mb instead of 20 Mb. On the contrary, the RH map extends 19 cR further than the available sequence, with the addition of *MRPS34* and *BM045*. The length of the RH map to be compared to the sequence was thus 324 cR₆₀₀₀ and not 343 cR₆₀₀₀. By using these figures, the ratio between the two maps was 46 kb/cR₆₀₀₀. The previously published figure of 61 kb/cR₆₀₀₀ for GGA7 used size estimations for this chromosome based on cytogenetic data [25]. The updated value for GGA7 using the genomic sequence assembly was thus 56 kb/cR₆₀₀₀. Similarly, a value of 37 kb/cR₆₀₀₀ can be calculated for GGA15 [20]. The breaking of chromosomes by radiation is a physical process, suggesting similar ratios to be expected for the different chromosomes. Future studies will indicate if the differences between chromosomes observed here are due to structural differences, or to errors in the RH maps and/or the sequence assembly. Given the mean retention frequency of 23.7% of GGA14 in the ChickRH6 panel containing 90 clones, we expect an average of 21.3 observations of such breaks per marker and thus a mean expected resolution power of 215 kb. However, since the retention frequency varies along the chromosome from 35% close to the centromere down to an average of 15% elsewhere, the expected resolution of the panel will vary accordingly from 146 to 340 kb.

It is noteworthy, that the two main discrepancies between the RH map and the genome sequence assembly involves the region close to the centromere: a portion of GGA14 sequence from this region was moved to the telomere region and was replaced by sequence fragments from other chromosomes, mainly GGA3. It is noticeable, that the proportion of markers corresponding to existing sequence fragments of unknown location in the sequence assembly was higher in this region of the RH map than in the other regions and that the only marker of the GGA14 RH map found on another chromosome in the genomic sequence assembly (*Rab1* on GGA18), was also mapped there. One explanation for the difficulties to assemble the sequence in this region could be a similarity between subtelomeric and pericentromeric repeats creating false joining of sequences.

4.5. Comparative mapping

By increasing the number of genes assigned to GGA14 to 48, we greatly improved the comparative mapping data available for this chromosome. If we consider a group of conserved gene segments as containing at least

two genes, 10 groups of global conservation were found with the human region HSA16p13.3 and one with the human region HSA17p11.2, indicated by coloured boxes in Figure 3. This comparison also showed eight genes, indicated in black in Figure 3, which could not be assigned to the conserved gene segments we defined. Apart for *MAP2K3* located on HSA17, all other seven genes (*ABCA3*, *BM045*, *CLCN7*, *MGC15416*, *MRPS34*, *NUBP2* and *UBE21*) are located in the region between 1.2 and 4.9 Mb on HSA16, in which the highest density of genes and EST were developed. The higher level of resolution thus obtained may partly account for the detection of such small segments of conserved gene order, but it is also possible that this region has undergone a higher number of rearrangements.

The development of EST markers based solely on the prior information of synteny conservation with HSA16p13.3 and HSA17p11.2 does not enable us to rule out the possibility that some small regions from GGA14 correspond to other HSA regions. However, as previously noted in other detailed comparative mapping studies, despite the high level of synteny conservation, a high number of intra-chromosomal rearrangements can be observed between the human and chicken genome.

Due to the lack of precision on the length of the regions of conserved synteny with humans available at the beginning of our work, we extended out of them when choosing EST markers and thus developed markers for other chicken chromosomes. This was particularly true in the case of markers from HSA17, for which the region of conserved synteny appeared to be quite small. As a result, in addition to the conservation between HSA17 and GGA14 demonstrated by *NT5M*, *SREBF1*, *LLGL1*, *PRPSAP2* and *GRAP*, small blocks homologous to regions located on GGA18 (in pink in Fig. 3) or GGA19 (in blue in Fig. 3) could be defined. Finally, *ACCN1* is located on GGA27 (in green in Fig. 3) and no hits for *GIT1* were found in the chicken genome sequence.

5. CONCLUSION

The first purpose of our work on the GGA14 RH map was to develop a dense map including a high number of genes, in order to validate the use of the ChickRH6 panel for a microchromosome and to provide detailed comparative mapping information. At the end of our project, the first draft chicken genome sequence was released and we used our data to test the sequence contig of GGA14. Although the sequence assembly is globally in good agreement with our data, we show that RH mapping can detect some errors, demonstrating its usefulness as a contribution towards a high quality assembly of the sequence.

GGA14 (cR₆₀₀₀)





Figure 3. Comparative mapping between the GGA14 RH map and the sequence maps of human regions HSA16p13.3 and HSA17p11.2. Markers localised on the genetic map are underlined on the RH map. Groups of global conserved gene order between chickens and humans are identified by coloured boxes. Genes or EST markers that could not be assigned to any such group are in black.

HSA16p13.3 (Mb)



HSA17p11.2 (Mb)




Figure 3. Continued.

Future developments towards a complete chicken RH framework map will now be based on the genomic sequence, using it for choosing STS markers regularly spaced along the chromosomes.

ACKNOWLEDGEMENTS

This work was supported by funds from the Génopôle de Toulouse Midi-Pyrénées and INRA-AGENAE.

REFERENCES

- [1] Abdrakhmanov I., Lodygin D., Geroth P., Arakawa H., Law A., Plachy J., Korn B., Buerstedde J.M., A large database of chicken bursal ESTs as a resource for the analysis of vertebrate gene function, *Genome Res.* 10 (2000) 2062–2069.
- [2] Aerts J., Crooijmans R., Cornelissen S., Hemmatian K., Veenendaal T., Jaadar A., van der Poel J., Fillon V., Vignal A., Groenen M., Integration of chicken genomic resources to enable whole-genome sequencing, *Cytogenet. Genome Res.* 102 (2003) 297–303.
- [3] Assaf S., Hazard D., Pitel F., Morisson M., Alizadeh M., Gondret F., Diot C., Vignal A., Douaire M., Lagarrigue S., Cloning of cDNA encoding the nuclear form of chicken sterol response element binding protein-2 (SREBP-2), chromosomal localization, and tissue expression of chicken SREBP-1 and -2 genes, *Poult. Sci.* 82 (2003) 54–61.
- [4] Boardman P.E., Sanz-Ezquerro J., Overton I.M., Burt D.W., Bosch E., Fong W.T., Tickle C., Brown W.R., Wilson S.A., Hubbard S.J., A Comprehensive Collection of Chicken cDNAs, *Curr. Biol.* 12 (2002) 1965–1969.
- [5] Boehnke M., Lange K., Cox D.R., Statistical methods for multipoint radiation hybrid mapping, *Am. J. Hum. Genet.* 49 (1991) 1174–1188.
- [6] Brown W.R., Hubbard S.J., Tickle C., Wilson S.A., The chicken as a model for large-scale analysis of vertebrate gene function, *Nat. Rev. Genet.* 4 (2003) 87–98.
- [7] Buitenhuis A.J., Crooijmans R.P., Bruijnesteijn van Coppenraet E.S., Veenendaal A., Groenen M.A., van der Poel J.J., Improvement of the comparative map of chicken chromosome 13, *Anim. Genet.* 33 (2002) 249–254.
- [8] Burt D.W., Bruley C., Dunn I.C., Jones C.T., Ramage A., Law A.S., Morrice D.R., Paton I.R., Smith J., Windsor D., Sazanov A., Fries R., Waddington D., The dynamics of chromosome evolution in birds and mammals, *Nature* 402 (1999) 411–413.
- [9] Crooijmans R.P., Vrebalov J., Dijkhof R.J., van der Poel J.J., Groenen M.A., Two-dimensional screening of the Wageningen chicken BAC library, *Mamm. Genome* 11 (2000) 360–363.

- [10] Crooijmans R.P., Dijkhof R.J., Veenendaal T., van der Poel J.J., Nicholls R.D., Bovenhuis H., Groenen M.A., The gene orders on human chromosome 15 and chicken chromosome 10 reveal multiple inter- and intrachromosomal rearrangements, *Mol. Biol. Evol.* 18 (2001) 2102–2109.
- [11] Crooijmans R., Dijkhof R., Aerts J., Hemmatien K., Cornelissen S., Veenendaal T., van der Poel J., Groenen M., A sequence ready BAC map of chicken chromosome 10, in: *Proceeding of the Plant and Animal Genomes XI Conference*, San Diego, California (2003) p. 225.
- [12] Dodgson J.B., Chicken genome sequence: a centennial gift to poultry genetics, *Cytogenet. Genome Res.* 102 (2003) 291–296.
- [13] Echard G., Gellin J., Gillois M., Localisation des gènes MP1, PKM2, NP sur le chromosome 3 du porc (*Sus scrofa* L.) et analyse cytogénétique d’une lignée de hamster chinois issue de la DON (wg3h), *Génét. Sél. Évol.* 16 (1984) 261–270.
- [14] Groenen M.A., Crooijmans R.P., Dijkhof R.J., Acar R., van der Poel J.J., Extending the chicken-human comparative map by placing 15 genes on the chicken linkage map, *Anim. Genet.* 30 (1999) 418–422.
- [15] Groenen M.A., Cheng H.H., Bumstead N., Benkel B.F., Briles W.E., Burke T., Burt D.W., Crittenden L.B., Dodgson J., Hillel J., Lamont S., de Leon A.P., Soller M., Takahashi H., Vignal A., A consensus linkage map of the chicken genome, *Genome Res.* 10 (2000) 137–147.
- [16] Groenen M.A.M., Crooijmans R.P.M.A., Structural genomics: integrating linkage, physical and sequence maps, in: Muir W.M., Aggrey S.E. (Ed.), *Poultry Genetics, Breeding and Technology*, CAB International, Wallingford, 2003, pp. 497–536.
- [17] Gyapay G., Schmitt K., Fizames C., Jones H., Vega-Czarny N., Spillett D., Muselet D., Prud’homme J.F., Dib C., Auffray C., Morissette J., Weissenbach J., Goodfellow P.N., A radiation hybrid map of the human genome, *Hum. Mol. Genet.* 5 (1996) 339–346.
- [18] International Human Genome Mapping Consortium, A physical map of the human genome, *Nature* 409 (2001) 934–941.
- [19] Jennen D.G., Crooijmans R.P., Kamps B., Acar R., Veenendaal A., van der Poel J.J., Groenen M.A., A comparative map of chicken chromosome 24 and human chromosome 11, *Anim. Genet.* 33 (2002) 205–210.
- [20] Jennen D.G., Crooijmans R.P., Morisson M., Grootemaat A.E., Van Der Poel J.J., Vignal A., Groenen M.A., A radiation hybrid map of chicken chromosome 15, *Anim. Genet.* 35 (2004) 63–65.
- [21] Lee M.K., Ren C.W., Yan B., Cox B., Zhang H.B., Romanov M.N., Sizemore F.G., Suchyta S.P., Peters E., Dodgson J.B., Construction and characterization of three BAC libraries for analysis of the chicken genome, *Anim. Genet.* 34 (2003) 151–152.
- [22] Lunetta K.L., Boehnke M., Multipoint radiation hybrid mapping: comparison of methods, sample size requirements, and optimal study characteristics, *Genomics* 21 (1994) 92–103.
- [23] Morisson M., Pitel F., Fillon V., Pouzadoux A., Berge R., Vit J.P., Zoorob R., Auffray C., Gellin J., Vignal A., Integration of chicken cytogenetic and genetic

- maps: 18 new polymorphic markers isolated from BAC and PAC clones, *Anim. Genet.* 29 (1998) 348–355.
- [24] Morisson M., Lemiere A., Bosc S., Galan M., Plisson-Petit F., Pinton P., Delcros C., Feve K., Pitel F., Fillon V., Yerle M., Vignal A., ChickRH6: a chicken whole-genome radiation hybrid panel, *Genet. Sel. Evol.* 34 (2002) 521–533.
- [25] Morisson M., Jiguet-Jiglaire C., Leroux S., Faraut T., Bardes S., Feve K., Genet C., Pitel F., Milan D., Vignal A., Development of a gene-based radiation hybrid map of chicken chromosome 7 and comparison to human and mouse, *Mamm. Genome* 15 (2004) 732–739.
- [26] Muller C., Denis M., Gentzbittel L., Faraut T., The Iccare web server: an attempt to merge sequence and mapping information for plant and animal species, *Nucleic Acids Res.* 32 (2004) W429–434.
- [27] Nanda I., Shan Z., Schartl M., Burt D.W., Koehler M., Nothwang H., Grutzner F., Paton I.R., Windsor D., Dunn I., Engel W., Staeheli P., Mizuno S., Haaf T., Schmid M., 300 million years of conserved synteny between chicken Z and human chromosome 9, *Nat. Genet.* 21 (1999) 258–259.
- [28] Rabie T.S., Crooijmans R.P., Morisson M., Andryszkiewicz J., van der Poel J.J., Vignal A., Groenen M.A., A radiation hybrid map of chicken Chromosome 4, *Mamm. Genome* 15 (2004) 560–569.
- [29] Ren C., Lee M.-K., Yan B., Ding K., Cox B., Romanov M.N., Price J.A., Dodgson J.B., Zhang H.-B., A BAC-Based physical map of the chicken genome, *Genome Res.* 13 (2003) 2754–2758.
- [30] Romanov M.N., Price J.A., Dodgson J.B., Integration of animal linkage and BAC contig maps using overgo hybridization, *Cytogenet. Genome Res.* 102 (2003) 277–281.
- [31] Schmid M., Nanda I., Guttenbach M., Steinlein C., Hoehn M., Schartl M., Haaf T., Weigend S., Fries R., Buerstedde J.M., Wimmers K., Burt D.W., Smith J., A'Hara S., Law A., Griffin D.K., Bumstead N., Kaufman J., Thomson P.A., Burke T., Groenen M.A., Crooijmans R.P., Vignal A., Fillon V., Morisson M., Pitel F., Tixier-Boichard M., Ladjali-Mohammedi K., Hillel J., Maki-Tanila A., Cheng H.H., Delany M.E., Burnside J., Mizuno S., First report on chicken genes and chromosomes 2000, *Cytogenet. Cell. Genet.* 90 (2000) 169–218.
- [32] Smith E.J., Lyons L.A., Cheng H.H., Suchyta S.P., Comparative mapping of the chicken genome using the East Lansing reference population, *Poult. Sci.* 76 (1997) 743–747.
- [33] Smith J., Burt D.W., Parameters of the chicken genome (*Gallus gallus*), *Anim. Genet.* 29 (1998) 290–294.
- [34] Smith J., Paton I.R., Murray F., Crooijmans R.P., Groenen M.A., Burt D.W., Comparative mapping of human Chromosome 19 with the chicken shows conserved synteny and gives an insight into chromosomal evolution, *Mamm. Genome* 13 (2002) 310–315.
- [35] Stewart E., McKusick K., Aggarwal A., Bajorek E., Brady S., Chu A., Fang N., Hadley D., Harris M., Hussain S., Lee R., Maratukulam A., O'Connor K., Perkins S., Piercy M., Qin F., Reif T., Sanders C., She X., Sun W.L., Tabar P.,

- Voyticky S., Cowles S., Fan J.B., Cox D.R., *et al.*, An STS-based radiation hybrid map of the human genome, *Genome Res.* 7 (1997) 422–433.
- [36] Suchyta S.P., Cheng H.H., Burnside J., Dodgson J.B., Comparative mapping of chicken anchor loci orthologous to genes on human chromosomes 1, 4 and 9, *Anim. Genet.* 32 (2001) 12–18.
- [37] Tirunagaru V.G., Sofer L., Cui J., Burnside J., An expressed sequence tag database of T-cell-enriched activated chicken splenocytes: sequence analysis of 5251 clones, *Genomics* 66 (2000) 144–151.
- [38] Venter J.C., Adams M.D., Myers E.W., Li P.W., Mural R.J., Sutton G.G., Smith H.O., *et al.*, The sequence of the human genome, *Science* 291 (2001) 1304–1351.
- [39] Voorrips R.E., MapChart: software for the graphical presentation of linkage maps and QTLs, *J. Hered.* 93 (2002) 77–78.