



HAL
open science

Influence of the degree of dietary fatty acid unsaturation on rumen fermentation parameters

Sylvie Giger-Reverdin, J.L. Le Pierres, Christine Duvaux-Ponter, P.
Morand-Fehr, Joseph-Alexandre J.-A. Tessier, G. Dupas, Olivier Martin, A.
Rouzeau, Daniel Sauvart

► **To cite this version:**

Sylvie Giger-Reverdin, J.L. Le Pierres, Christine Duvaux-Ponter, P. Morand-Fehr, Joseph-Alexandre J.-A. Tessier, et al.. Influence of the degree of dietary fatty acid unsaturation on rumen fermentation parameters. *Options Méditerranéennes. Série A: Séminaires Méditerranéens*, 2004, 59, pp.67-71. hal-02670693

HAL Id: hal-02670693

<https://hal.inrae.fr/hal-02670693v1>

Submitted on 31 May 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Copyright

Influence of the degree of dietary fatty acid unsaturation on rumen fermentation parameters

S. Giger-Reverdin, J.L. Le Pierres, C. Duvaux-Ponter, P. Morand-Fehr, J. Tessier, G. Dupas, O. Martin, A. Rouzeau and D. Sauvant

Physiologie de la Nutrition et Alimentation, Unité Mixte de Recherche, INRA-INA Paris-Grignon, 16 rue Claude Bernard, 75231 Paris Cedex 05, France

SUMMARY – The influence of the degree of dietary fatty acid unsaturation on rumen fermentative activity was tested by comparing two extruded oilseeds (rapeseed or R and linseed or L) incorporated at a level of 8% dry matter (DM) in ruminant diets. With the R diet (rich in C_{18:1}), the ruminal pH of fistulated goats was statistically lower and the number of protozoa was higher than with the L diet (rich in C_{18:3}). However, the molar composition of the volatile fatty acid (VFA) did not differ between diets at the different sampling times. The *in sacco* dry matter degradability of the diets was measured on fistulated cows adapted to the experimental diets. It was higher with the R diet. In conclusion, the increase in the degree of dietary fatty acid unsaturation seems to decrease the ruminal fermentative activity.

Key words: Dietary fatty acids, unsaturation, oilseeds, rumen, ruminants.

RESUME – "Influence du degré d'insaturation des acides gras sur les paramètres de fermentation ruminale". Afin de mesurer l'effet du degré d'insaturation des acides gras alimentaires sur l'activité ruminale, nous avons comparé deux graines extrudées (colza ou C et lin ou L) incorporées à un niveau de 8% de matière sèche (MS) dans des régimes destinés à des ruminants. Le régime C (riche en C_{18:1}) a induit un pH ruminal des chèvres fistulées statistiquement plus faible que le régime L (riche en C_{18:3}). Le nombre des protozoaires a été plus élevé avec le régime C qu'avec le régime L, mais la composition molaire en acides gras volatils (AGV) n'a pas été influencée par le changement de régime. La dégradabilité *in sacco* de la matière sèche des régimes a été mesurée sur des vaches recevant les deux régimes expérimentaux. Elle a été plus élevée pour le régime C. En conclusion, l'augmentation du degré d'insaturation des matières grasses alimentaires semble décroître l'activité fermentaire du rumen.

Mots-clés : Matière grasse alimentaire, insaturation, graines oléagineuses, rumen, ruminants.

Introduction

Fat is often given in early lactation to ruminants because its high energy level is of particular importance at a physiological stage when energy intake is a limiting factor for milk production. Several authors pointed out that fat might alter the ruminal microbial ecosystem, because it often has a negative effect on feed intake and fibre digestibility (Doreau and Chilliard, 1997). This effect is more noticeable with polyunsaturated fatty acids (Czerkawski, 1973; Palmquist and Jenkins, 1980; Sutton *et al.*, 1983).

The objective of this study was to test the influence of the degree of dietary fatty acid unsaturation of extruded oilseeds on rumen fermentative activity. The extrusion process is of interest because it partially protects protein from ruminal degradation (Clinquart *et al.*, 1993). Therefore, in the literature, an extruded oilseed has been usually compared to a control seed (McGuffey and Schingoethe, 1982; Ferlay *et al.*, 1992) rather than to another extruded oilseed. In this paper, we compared two extruded oilseeds [rapeseed (R) and linseed (L)] differing in the degree of their dietary fatty acids unsaturation: they were respectively rich in C_{18:1} and C_{18:3}. This kind of oilseeds can modify significantly milk fatty acids composition, and especially the C₁₈ fractions (Duvaux-Ponter *et al.*, 2001; Giger-Reverdin *et al.*, 2001). We used two complementary approaches: the *in vivo* kinetics of rumen fermentation parameters and the *in sacco* dry matter degradability of the experimental diets with fistulated animals receiving these diets.

Material and methods

Diets

The oilseeds were incorporated at a level of 8% DM in the diet which also included corn silage (50% DM) and dehydrated alfalfa (10% DM). The rest of the diet was made up of concentrate (32% DM) in order to meet the requirements of dairy goats fed in early lactation. Diets were offered *ad libitum* after morning milking (around 08:00) which is considered as T0.

In vivo kinetics of rumen parameters

Four fistulated goats in mid-lactation received the 2 diets (2 goats for each diet). After 4 weeks of adaptation, ruminal fluid was collected before the morning feed (time = 0) and 2, 4, 6 and 8 hours after feeding. Kinetics were performed in duplicate on each goat on 2 separate days. The pH was immediately recorded. Samples were quickly deep-frozen at -20°C with 5% orthophosphoric acid until their analysis for volatile fatty acid (VFA) content by capillary chromatography (Broudiscou and Lassalas, 2000). The number of protozoa was also estimated 6 hours after the morning meal by the method described by Broudiscou *et al.* (1997).

In sacco dry matter degradability

A modification of the nylon bag technique proposed by Michalet-Doreau *et al.* (1987) was used to measure the disappearance of dry matter (DM) of the two diets. Bags were removed after 2, 4, 8, 16, 24 and 48 hours of incubation. Each of the three fistulated cows received 8 kg DM of the two experimental diets at two different periods instead of the standard diet used for determining the protein dispersibility index (PDI) value of feedstuffs (Michalet-Doreau *et al.*, 1987). These daily allowances met the requirements of the animals. Within each period, 2 kinetics of measurements were performed.

Statistical analysis

In vivo kinetics of rumen parameters

The statistical design was a double-repeated one (duplicate day and several samplings on the same animal at a given day). We used a split-plot analysis with the "AR(1) proc mixed" procedure of SAS/STAT (1987). We took into account the following effects: diet, day, hour of sampling and their interactions. This model was used since the same goats were sampled at different times (Littell *et al.*, 1998).

In sacco dry matter degradability

These results were also analysed with a proc mixed procedure. Following effects were tested: diet received by the cow, feedstuff, duration of incubation and their interactions.

In all the statistical procedures, we removed from the model non-significant interactions or effects.

Results and discussion

The rapeseed meal induced more fermentation than the linseed one, especially during the first 2 hours of fermentation. The pH decreased first very quickly and significantly. It reached a plateau value of 6.03 after 4 h and then increased significantly after 6 h, just before the second meal of the day (Fig. 1). Linseed diet had a pH significantly higher than the rapeseed one, but there was no interaction diet-hour of sampling.

The quantity and the different percentages of VFA were studied at T0, T4 and T8. The dietary effect was only significant for total VFA in agreement with pH variations (Table 1).




Fig. 1. Evolution of pH after feeding in four fistulated goats receiving either a linseed diet or a rapeseed one.

Table 1. Dietary effect on the ruminal kinetics for the concentration of volatile fatty acids (VFA) and percentages of the different VFA

Diet	Sampling time					
	0 h		4 h		8 h	
	Rapeseed	Linseed	Rapeseed	Linseed	Rapeseed	Linseed
VFA (mmoles/l)	95 ^a	61 ^b	165 ^c	152 ^{c,d}	146 ^d	118 ^e
% C2	65.5 ^a	63.6 ^a	62.0 ^{b,c}	60.2 ^b	61.8 ^c	62.4 ^c
% C3	16.6 ^a	16.5 ^a	22.0 ^b	22.4 ^b	22.3 ^b	20.6 ^b
% iC4	1.6 ^a	2.5 ^b	1.0 ^c	0.7 ^c	1.0 ^c	0.9 ^c
% C4	12.6	12.2	12.3	13.2	12.5	13.2
% iC5	2.2 ^a	3.2 ^b	0.9 ^c	0.8 ^c	0.9 ^c	0.9 ^c
% C5	1.1 ^a	1.0 ^a	1.4 ^b	1.2 ^{a,b}	1.3 ^{b,c}	1.1 ^{a,c}
% C6	0.1	0.1	0.3	0.4	0.2	0.4

^{a,b,c,d,e} Within a row, values with the same letter do not differ significantly ($P > 0.05$).

The VFA composition was modified by dietary intake since the acetic acid percentage decreased very quickly while the percentage of propionic acid increased in the first 2 hours after the morning feeding. This is in agreement with the specific effect of corn silage or the addition of fat, which enhance propionate production (Broudiscou *et al.*, 1990). The non-significant dietary effect is in agreement with the results of Sutter *et al.* (2000) who compared rapeseeds and linseeds and with the review of Sauvart and Bas (2001) who pointed out that the degree of unsaturation of fatty acids did not significantly affect the C_2/C_3 ratio.

After 6 hours of fermentation, the number of total protozoa was significantly higher for the rapeseed meal compared to the linseed one ($1.36/10^6$ ml vs $1.05/10^6$ ml). This was due only to the holotrich protozoa. This agrees with Sauvart and Bas (2001) who stressed that the increase in dietary fatty unsaturation decreased the ruminal density of protozoa. Sutter *et al.* (2000) found a non-

significant decrease when replacing rapeseed by linseed. These high values point out that extruded seeds do not have a defaunating effect on the contrary to oils (Sutton *et al.*, 1983; Broudiscou *et al.*, 1990).

Whatever the duration of the incubation, DM degradability of the samples (diets R or L offered to goats) was higher when incubated in cows receiving the rapeseed diet (dotted lines, see Fig. 2). This could be related with a linearly decrease in apparent ruminal neutral detergent fibre (NDF) digestibility when esterification of fat decreased (Elliott *et al.*, 1997).




Fig. 2. Effect of the diets given to fistulated cows on *in sacco* dry matter degradability of diets samples (R: rapeseed, L: linseed).

The samples of the rapeseed diet (Sample R) were significantly less degraded than those of the linseed one (Sample L), whatever the kind of diets received by the cows. The differences were statistically significant between 4 and 24 hours of incubation. There was no significant interaction between samples and diets. This means that the rapeseed diet which was less degraded than the linseed one, seemed to induce less fermentation in cows, on the contrary to goats.

Conclusion

These two experiments bring new information about extruded oilseeds that had been seldom studied on the contrary to oils. They indicated that the degree of dietary fatty acid unsaturation had an influence on rumen fermentation. When it was increased, fermentation seemed to be less efficient, but the proportion of the different VFA in the rumen remained the same.

References

- Broudiscou, L., van Nevel, C.J. and Demeyer, D.L. (1990). Effect of soya oil hydrolysate on rumen digestion in defaunated and refaunated sheep. *Anim. Feed Sci. Tech.*, 30: 51-67.
- Broudiscou, L.P. and Lassalas, B. (2000). Effects of *Lavandula officinalis* and *Equisetum arvense* dry extracts and isoquercitrin on the fermentation of diets varying in forage contents by rumen microorganisms in batch culture. *Reprod. Nutr. Dev.*, 40: 431-440.

- Broudiscou, L.P., Papon, Y., Fabre, M. and Broudiscou, A.F. (1997). Maintenance of rumen protozoa populations in a dual outflow continuous fermenter. *J. Sci. Food Agr.*, 75: 273-280.
- Clinquart, A., Istasse, L., van Eenaeme, C., Diez, M., Dufrasne, I. and Bienfait, J.M. (1993). Influence de l'extrusion de mélanges de graines de lin et d'orge, de graines de pois et de colza, et de fèves de soja, sur la dégradabilité dans le rumen de leurs fractions azotée et lipidique et sur leur composition en acides gras. *Ann. Zootech.*, 42: 130-131.
- Czerkawski, J.W. (1973). Effect of linseed oil fatty acids and linseed oil on rumen fermentation in sheep. *J. Agr. Sci.*, 81: 517-531.
- Doreau, M. and Chilliard, Y. (1997). Digestion and metabolism of dietary fat in farm animals. *Br. J. Nutr.*, 78: 515-535.
- Duvaux-Ponter, C., Roussel, S., Giger-Reverdin, S., Morand-Fehr, P. and Sauvant, D. (2001). Influence du degré d'insaturation de matières grasses alimentaires (lin vs colza) sur la composition en acides gras de la matière grasse du colostrum caprin et le transfert passif de l'immunité chez le chevreau nouveau-né. In: *8èmes Rencontres Recherches Ruminants*, 5-6 December 2001. Institut de l'Elevage, Paris, France, p. 102.
- Elliott, J.P., Drackley, J.K., Aldrich, C.G. and Merchen, N.R. (1997). Effects of saturation and esterification of fat sources on site and extent of digestion in steers: Ruminal fermentation and digestion of organic matter, fiber, and nitrogen. *J. Anim. Sci.*, 75: 2803-2812.
- Ferlay, A., Legay, F., Bauchart, D., Poncet, C. and Doreau, M. (1992). Effect of a supply of raw or extruded rapeseeds on digestion in dairy cows. *J. Anim. Sci.*, 70: 915-923.
- Giger-Reverdin, S., Weill, P., Duvaux-Ponter, C., Morand-Fehr, P., Rouzeau, A. and Sauvant, D. (2001). Influence du degré d'insaturation de matières grasses alimentaires (lin vs colza) sur la composition en acides gras de la matière grasse du lait de chèvre en début de lactation. In: *8èmes Rencontres Recherches Ruminants*, 5-6 December 2001. Institut de l'Elevage, Paris, France, p. 101.
- Littell, R.C., Henry, P.R. and Ammerman, C.B. (1998). Statistical analysis of repeated measures data using SAS procedures. *J. Anim. Sci.*, 76: 1216-1231.
- McGuffey, R.K. and Schingoethe, D.J. (1982). Whole sunflower seeds for high producing dairy cows. *J. Dairy Sci.*, 65: 1479-1483.
- Michalet-Doreau, B., Vérité, R. and Chapoutot, P. (1987). Methodology of estimating degradability *in sacco* of nitrogen in feed in the rumen. *Bull. Tech. CRZV Theix*, 69: 5-7.
- Palmquist, D.L. and Jenkins, T.C. (1980). Fat in lactation rations: Review. *J. Dairy Sci.*, 63: 1-14.
- SAS/STAT (1987). *Guide for Personal Computers*. SAS Institute, Inc., Cary, NC, USA.
- Sauvant, D. and Bas, P. (2001). La digestion des lipides chez le ruminant. *INRA Prod. Anim.*, 5(14): 303-310.
- Sutter, F., Casutt, M.M., Ossowsky, D.A., Scheeder, M.R.L. and Kreuzer, M. (2000). Comparative evaluation of rumen-protected fat, coconut oil and various oilseeds supplemented to fattening bulls. 1. Effects on growth, carcass and meat quality. *Arch. Anim. Nutr.*, 53: 1-23.
- Sutton, J.D., Knight, R., McAllan, A.B. and Smith, R.H. (1983). Digestion and synthesis in the rumen of sheep given diets supplemented with free and protected oils. *Br. J. Nutr.*, 49: 419-432.