



HAL
open science

Isomerization increases the postprandial oxidation of linoleic acid but not alpha-linolenic acid in men

Lionel Brétilon, Jean-Michel Chardigny, Jean-Louis Sébédio, Jean-Pierre Noël, C.M. Scrimgeour, C.E. Fernie, Olivier Loreau, P. Gachon, B. Beaufrère

► **To cite this version:**

Lionel Brétilon, Jean-Michel Chardigny, Jean-Louis Sébédio, Jean-Pierre Noël, C.M. Scrimgeour, et al.. Isomerization increases the postprandial oxidation of linoleic acid but not alpha-linolenic acid in men. *Journal of Lipid Research*, 2001, 42 (6), pp.995-997. hal-02674145

HAL Id: hal-02674145

<https://hal.inrae.fr/hal-02674145v1>

Submitted on 31 May 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Isomerization increases the postprandial oxidation of linoleic acid but not α -linolenic acid in men

L. Bretillon,* J. M. Chardigny,^{1,*} J. L. Sébédio,* J. P. Noël,[†] C. M. Scrimgeour,[§] C. E. Fernie,[§] O. Loreau,[†] P. Gachon,** and B. Beaufrère**

Unité de Nutrition Lipidique,* Institut National de la Recherche Agronomique, 21065 Dijon, France; Service des Molécules Marquées,[†] CEA-Saclay, 91191 Gif-sur-Yvette, France; Scottish Crop Research Institute,[§] Dundee DD2 5DA, Scotland; and Unité du Métabolisme Protéino Énergétique,** Université d'Auvergne, INRA, CRNH, 63001 Clermont-Ferrand, France

Abstract Human lipid intake contains various amounts of *trans* fatty acids. Refined vegetable and frying oils, rich in linoleic acid and/or α -linolenic acid, are the main dietary sources of *trans*-18:2 and *trans*-18:3 fatty acids. The aim of the present study was to compare the oxidation of linoleic acid, α -linolenic acid, and their major *trans* isomers in human volunteers. For that purpose, TG, each containing two molecules of [1-¹³C]linoleic acid, α -[1-¹³C]linolenic acid, [1-¹³C]-9*cis*,12*trans*-18:2, or [1-¹³C]-9*cis*,12*cis*,15*trans*-18:3, were synthesized. Eight healthy young men ingested labeled TG mixed with 30 g of olive oil. Total CO₂ production and ¹³CO₂ excretion were determined over 48 h. The pattern of oxidation was similar for the four fatty acids, with a peak at 8 h and a return to baseline at 24 h. Cumulative oxidation over 8 h of linoleic acid, 9*cis*,12*trans*-18:2, α -linolenic acid, and 9*cis*,12*cis*,15*trans*-18:3 were, respectively, 14.0 \pm 4.1%, 24.7 \pm 6.7%, 23.6 \pm 3.3%, and 23.4 \pm 3.7% of the oral load, showing that isomerization increases the postprandial oxidation of linoleic acid but not α -linolenic acid in men.—Bretillon, L., J. M. Chardigny, J. L. Sébédio, J. P. Noël, C. M. Scrimgeour, C. E. Fernie, O. Loreau, P. Gachon, and B. Beaufrère. Isomerization increases the postprandial oxidation of linoleic acid but not α -linolenic acid in men. *J. Lipid Res.* 2001. 42: 995–997.

Supplementary key words human • oxidation • *trans* fatty acid

Heat treatment of vegetable oils induces the isomerization of essential PUFA, that is, linoleic and α -linolenic acids (1–3). These geometric isomers of essential fatty acids are accompanied in the diet by *trans* isomers of monounsaturated fatty acids, leading to consumption of *trans* fatty acids from 3.0 to 10.6 g per capita per day in Western countries (4). 9*cis*,12*trans*-18:2 and 9*cis*,12*cis*,15*trans*-18:3 are the major *trans* PUFA found in vegetable and frying oils (1–3). In rats, these compounds are desaturated and elongated into *trans* isomers of arachidonic acid and eicosapentaenoic acid (5–7). From previous studies of rats (8–12), it is concluded that *i)* dietary α -linolenic acid is oxidized more than

linoleic acid and *ii)* *trans* isomerization increases 18:2 oxidation but does not affect 18:3 oxidation. To test the hypothesis of a similar pattern in humans, we determined postprandial oxidation of ¹³C-labeled *cis* or *trans* PUFA given as TG to healthy humans.

MATERIALS AND METHODS

Materials

[1-¹³C]linoleic acid, [1-¹³C]linolenic acid, [1-¹³C]-9*cis*,12*cis*,15*trans*-18:3, and [1-¹³C]-9*cis*,12*trans*-18:2 were first synthesized as described elsewhere (13) and used for synthesizing TG following the method of Redden et al. (14). The products were >98% TG as estimated by HPLC and had a labeled fatty acid:oleic acid ratio of between 1.83 and 2.00 as estimated by gas chromatography.

Human subjects

Ten healthy men (mean age of 25 \pm 3 years, body mass index 21.3 \pm 1.8 kg/m², mean \pm SD) were recruited. Blood pressure, cholesterolemia, and triacylglycerolemia were within normal ranges. None were taking any medication, their dietary habits were stable, and they were nonsmokers.

Protocol

The study had been approved by the local ethics committee (Hôpitaux Civils de Clermont-Ferrand, France) according to the French Hurriet law. Informed written consent of the participants had been obtained. The study was designed as four periods of 2 days, separated one from another by at least 1 week. Six subjects performed the whole study and received the four fatty acids. Two additional subjects were involved in the 18:2 part of the study whereas two others performed the 18:3 study. In each study (18:2 and 18:3), the *cis* or *trans* isomers were administered in a randomized, double-blind fashion. The four study periods were strictly identical except for the nature of the ingested labeled TG.

Abbreviations: V'CO₂, rate of production of carbon dioxide.

¹ To whom correspondence should be addressed.

e-mail: chardign@dijon.inra.fr

Subjects entered the experimental unit at 7:00 AM after an overnight fast. Two samples of expired breath were collected in Vacutainer® tubes (Becton Dickinson, Grenoble, France) in order to determine the basal ^{13}C abundance in the expired CO_2 . One hour after arrival, the subjects ingested the labeled TG (1.40 g for the 18:2, and 0.87 g for the 18:3) mixed with 30 g of olive oil (Lesieur Alimentaire, Neuilly, France). Expired breath collection was performed as described above 1, 2, 3, 5, and 8 h after TG ingestion. Continuous indirect calorimetry (Deltatrac metabolic monitor; Datex, Helsinki, Finland) was performed 2, 5, and 8 h after TG ingestion for periods of 30 min, in order to measure the CO_2 production. The subjects were fasted during the first 8 h. A meal containing food items poor in ^{13}C was served afterward. Expired breath samples were again collected at 12, 24, and 48 h after ingestion but no CO_2 production measurements were performed at these time points.

Analytical procedures

^{13}C -enrichment analysis of the expired breath was done by gas chromatography-combustion-isotope ratio mass spectrometry. Briefly, 40 μl samples were injected in duplicate into a gas chromatograph (5890; Hewlett-Packard, Palo Alto, CA) equipped with a HAYSEP Q column (Chrompack, Les Ulis, France). The oven temperature was 110°C for 5 min. The CO_2 peak was then injected on-line into an isotope ratio mass spectrometer (μG as system; Fisons Instruments, VG Isotech, Middlewich, England). The ^{13}C enrichment of expired CO_2 was determined by comparison with a standard of known ^{13}C abundance (Pee Dee Belemnite, PDB). The rate of $^{13}\text{CO}_2$ production was obtained by multiplying $^{13}\text{CO}_2$ enrichment by the CO_2 production.

Calculations and statistical analyses

All results are expressed as mean \pm standard deviation. The cumulative $^{13}\text{CO}_2$ production corresponds to the area under the curve and was calculated by the trapezoidal method. Within a pair of studies (18:2 or 18:3), comparisons were made by paired, bilateral *t*-tests. Comparison between the 4 periods was made on the 6 subjects who performed the entire study, by ANOVA followed by a Fisher PLSD test.

RESULTS

The profile of ^{13}C enrichment of the expired breath air was the same, regardless of which fatty acid was ingested (Fig. 1A and B). It increased 1 h after TG ingestion, peaked between 5 and 8 h, and decreased thereafter at 12 h to reach the basal ^{13}C level at 24 and 48 h. Significantly higher ^{13}C levels were observed after ingestion of *trans*-18:2. $V'\text{CO}_2$ values were similar between the two 18:2 fatty acids ($207 \pm 10 \text{ ml}\cdot\text{min}^{-1}$ for linoleic acid and $202 \pm 10 \text{ ml}\cdot\text{min}^{-1}$ for the *trans*-18:2). Therefore the cumulative amount of tracer excreted into breath was much higher for *trans*-18:2 than for linoleic acid. The fraction of *trans*-18:2 oxidized over 8 h was $24.7 \pm 6.7\%$ of the oral load, higher than the corresponding fraction of linoleic acid, which was $14.0 \pm 4.1\%$ ($P < 0.001$; Fig. 2). In contrast to the 18:2 fatty acids, isomerization of 18:3 fatty acid did not affect its oxidation (Fig. 1B), which was similar for both isomers (23.6 ± 3.3 and $23.4 \pm 3.7\%$ for α -linolenic acid and *trans*-18:3, respectively; Fig. 2). $^{13}\text{CO}_2$ production over the 8 h after ingestion of the α -linolenic acid ($23.6 \pm 3.3\%$) was almost twice as high as after ingestion of the lin-




Fig. 1. $^{13}\text{CO}_2$ enrichment ($\delta \text{‰}$ vs. PDB) after ingestion of linoleic acid (18:2n-6) or 9*cis*,12*trans*-18:2 (12*trans*-18:2) (A), or after ingestion of α -linolenic acid (18:3n-3) or 9*cis*,12*cis*,15*trans*-18:3 (15*trans*-18:3) (B) by human subjects.

oleic acid ($14.0 \pm 4.1\%$) when expressed as the fraction of tracer recovered ($P < 0.001$; Fig. 2).

DISCUSSION

In the present work, we have looked at the $^{13}\text{CO}_2$ production from linoleic and α -linolenic acids as well as from two of their main *trans* geometric isomers in apparently healthy young men. To our knowledge, this article is the first report of the postprandial oxidation in human volunteers of linoleic and α -linolenic acids given as TG, which is the dietary form for fatty acid ingestion. As a means of comparison, it would be rather difficult to draw a general





Fig. 2. Excretion of $^{13}\text{CO}_2$ during the 8 h after the ingestion of linoleic acid (18:2n-6), 9*cis*,12*trans*-18:2 (12*trans*-18:2), α -linolenic acid (18:3n-3), and 9*cis*,12*cis*,15*trans*-18:3 (15*trans*-18:3) by human subjects. Because the amounts of tracer given as 18:2 and 18:3 fatty acids were different, $^{13}\text{CO}_2$ excretion was normalized for the amount of tracer ingested. * Significantly different from the value obtained after ingestion of the three other fatty acids given as TG ($P < 0.05$).

conclusion from the previous rare studies of the oxidation of linoleic and α -linolenic acids in humans (15–18), considering that these were carried out under conditions that were different from each other and from ours. These studies used nondietary forms of the fatty acids (free fatty acids or methyl esters). Hence there may have been low bioavailability, as has already been suggested for the methyl esters (19). In addition, some were performed in men but others were carried out in women or lactating women, using either uniformly labeled or carboxyl-labeled fatty acids. Moreover, to account for incomplete recovery of labeled CO₂ from the bicarbonate pool, some authors used a correction factor to evaluate the oxidation of labeled tracers (15, 16, 20), whereas others did not (17, 18). The exact value of the correction factor remains controversial, from 0.5 to 0.9 (21). Therefore, our data were not corrected by this factor. Finally, duration of CO₂ collection and dietary conditions varied from one study to the other.

Nevertheless, despite all the differences, our figures for PUFA oxidation are of the same order of magnitude as most previously published figures. Most importantly, we confirm that α -linolenic acid is used for energy production in humans to a greater extent than is linoleic acid when the PUFA were given as TG, their naturally occurring dietary form. This difference has already been reported in rats (9, 11, 12) and in humans (17), although with labeled free fatty acids. This suggests that the digestibility of the TG is not involved in the difference observed between oxidation of the two PUFA. Assuming a ratio of linoleic acid to α -linolenic acid in the diet of between 5 and 10, the greater catabolism of α -linolenic acid may consistently increase the difference in the availability of these two essential fatty acids for their conversion into long-chain PUFA, which is generally considered to characterize the “essential” nature of linoleic and α -linolenic acids. The most important result is the greater catabolism of *trans*-18:2 compared with linoleic acid. This is consistent with the results we obtained in rats (12). The difference in oxidation may be due to selective formation of activated substrates (via the acyl-carnitines) for the β -oxidation pathway, or to different affinities for carrier proteins, such as albumin.

In conclusion, the present work demonstrates that the *trans* isomers of essential fatty acids are used for energy production in humans at least as well as their *cis* counterparts. Because these *trans* PUFA are also converted into higher derivatives (2, 5–7, 22), it is conceivable that their metabolic effects might be modulated by their bioavailability for this pathway, which in turn may be dependent of their use for energy production. 

We thank Nurse Liliane Morin (Laboratoire de Nutrition Humaine de Clermont-Ferrand, France) for excellent and skillful technical assistance. This work was supported by a grant from the European Commission (FAIR contract 95-0594, entitled Nutritional and Health Impact of *trans* Polyunsaturated Fatty Acids in European Populations), and by a PhD fellowship from INRA and Lesieur Alimentaire (L.B.).

Manuscript received 22 January 2001 and in revised form 13 February 2001.

REFERENCES

- Ackman, R. G., S. N. Hooper, and D. L. Hooper. 1974. Linolenic acid artefacts from the desodorization of oils. *J. Am. Oil Chem. Soc.* **51**: 42–49.
- Grandgirard, A., J. L. Sébédio, and J. Fleury. 1984. Geometrical isomerization of linolenic acid during heat treatment of vegetable oils. *J. Am. Oil Chem. Soc.* **61**: 1563–1568.
- Sébédio, J. L., A. Grandgirard, and J. Prévost. 1988. Linoleic acid isomers in heat treated sunflower oils. *J. Am. Oil Chem. Soc.* **65**: 362–366.
- Chen, Z. Y., G. Pelletier, R. Hollywood, and W. M. N. Ratnayake. 1995. *Trans* fatty acids isomers in Canadian human milk. *Lipids.* **30**: 15–21.
- Beyers, E. C., and E. A. Emken. 1991. Metabolites of *cis,trans*, and *trans,cis* isomers of linoleic acid in mice and incorporation into tissue lipids. *Biochim. Biophys. Acta.* **1082**: 275–284.
- Ratnayake, W. M. N., Z. Y. Chen, G. Pelletier, and D. Weber. 1994. Occurrence of 5*c,8c,11c,15t*-eicosatetraenoic acid and other unusual polyunsaturated fatty acids in rats fed partially hydrogenated canola oil. *Lipids.* **29**: 707–714.
- Berdeaux, O., J. L. Sébédio, J. M. Chardigny, J. P. Blond, T. Mairrot, J. M. Vatièle, D. Poullain, and J. P. Noël. 1996. Effects of *trans* n-6 fatty acids on the fatty acid profile of tissues and liver microsomal desaturation in the rat. *Grasas Aceit.* **47**: 86–99.
- Coots, R. H. 1964. A comparison of the metabolism of *cis,cis*-linoleic, *trans,trans*-linoleic and a mixture of *cis,trans* and *trans,cis*-linoleic acids in the rat. *J. Lipid Res.* **5**: 473–476.
- Leyton, J., P. J. Drury, and M. A. Crawford. 1987. Differential oxidation of saturated and unsaturated fatty acids *in vivo* in the rat. *Br. J. Nutr.* **57**: 383–393.
- Anderson, R. L. 1967. Oxidation of the geometric isomers of Δ^9 -octadecenoic acid by rat liver mitochondria. *Biochim. Biophys. Acta.* **144**: 18–24.
- Cunnane, S. C., and M. J. Anderson. 1997. The majority of dietary linoleate in growing rats is beta-oxidized or stored in visceral fat. *J. Nutr.* **127**: 146–152.
- Bretillon, L., J. M. Chardigny, J. L. Sébédio, D. Poullain, J. P. Noël, and J. M. Vatièle. 1998. Oxidative metabolism of (1-¹⁴C) mono-*trans* isomers of linoleic and α -linolenic acids in the rat. *Biochim. Biophys. Acta.* **1390**: 207–214.
- Loreau, O., A. Maret, D. Poullain, J. Chardigny, J. L. Sebedio, B. Beaufre, and J. P. Noel. 2000. Large scale syntheses of *cis* and *trans* [1-¹³C]-fatty acids. I. Preparation of (9*Z*,12*Z*)- and (9*Z*,12*E*)-[1-¹³C]-octadeca-9,12-dienoic acid. *Chem. Phys. Lipids.* **106**: 65–78.
- Redden, P. R., X. Lin, and D. F. Horrobin. 1996. Comparison of the grignard deacylation TLC and HPLC methods and high resolution ¹³C-NMR for the *sn*-2 positional analysis of triacylglycerols containing gamma-linolenic acid. *Chem. Phys. Lipids.* **79**: 9–19.
- Jones, P. J. H., P. B. Pencharz, and M. T. Clandinin. 1985. Whole body oxidation of dietary fatty acids: implications for energy utilization. *Am. J. Clin. Nutr.* **42**: 769–777.
- Demmelair, H., M. Baumheuer, B. Koletzko, K. Dokoupil, and G. Kratl. 1998. Metabolism of U-¹³C-labeled linoleic acid in lactating women. *J. Lipid Res.* **39**: 1389–1396.
- DeLany, J. P., M. M. Windhauser, C. M. Champagne, and G. A. Bray. 2000. Differential oxidation of individual dietary fatty acids in humans. *Am. J. Clin. Nutr.* **72**: 905–911.
- Vermunt, S. H. F., R. P. Mensink, M. M. G. Simonis, and G. Hornstra. 2000. Effects of dietary α -linolenic acid on the conversion and oxidation of ¹³C- α -linolenic acid. *Lipids.* **35**: 137–142.
- Yang, L. Y., A. Kuksis, and J. J. Myher. 1990. Intestinal absorption of menhaden and rapeseed oils and their fatty acid methyl esters in the rat. *Biochem. Cell. Biol.* **68**: 480–491.
- Binnert, C., C. Pachiaudi, M. Beylot, D. Hans, J. Vandermander, P. Chantre, J. P. Riou, and M. Laville. 1998. Influence of human obesity on the metabolic fate of dietary long- and medium-chain triacylglycerols. *Am. J. Clin. Nutr.* **67**: 595–601.
- Clugston, G. A., and P. J. Garlick. 1983. Recovery of infused [¹⁴C] bicarbonate as ¹⁴CO₂ in man. *Clin. Sci.* **64**: 231–233.
- Bretillon, L., J. M. Chardigny, J. P. Noël, and J. L. Sébédio. 1998. Desaturation and chain elongation of 1-¹⁴C mono-*trans* isomers of linoleic and α -linolenic acids in perfused rat liver. *J. Lipid Res.* **39**: 2228–2239.