



Crystal structure of a four-copper laccase complexed with an arylamine: insights into substrate recognition and correlation with kinetics

Thomas Bertrand, Claude Jolival, Pierre Briozzo, Eliane Caminade, Nathalie Joly, Catherine Madzak, Christian Mougin

► To cite this version:

Thomas Bertrand, Claude Jolival, Pierre Briozzo, Eliane Caminade, Nathalie Joly, et al.. Crystal structure of a four-copper laccase complexed with an arylamine: insights into substrate recognition and correlation with kinetics. *Biochemistry*, 2002, 41 (23), pp.7325-7333. 10.1021/bi0201318S0006-2960(02)00131-9 . hal-02677382

HAL Id: hal-02677382

<https://hal.inrae.fr/hal-02677382>

Submitted on 20 Jan 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - ShareAlike 4.0 International License

Crystal structure of a four-copper laccase complexed with an arylamine: insights into substrate recognition and correlation with kinetics^{II,∇}

Thomas Bertrand,[‡] Claude Jolival,[§] Pierre Briozzo,^{II,*} Eliane Caminade,[§] Nathalie Joly,^{II}
Catherine Madzak[⊥] and Christian Mougin[§]

[‡]Laboratoire d'Enzymologie et Biochimie Structurales, UPR 9063 du CNRS, 91198
Gif-sur-Yvette Cedex, France.

[§]Laboratoire de Phytopharmacie et Médiateurs Chimiques, INRA, route de Saint-Cyr, 78026
Versailles Cedex, France.

^{II}Laboratoire de Chimie Biologique, Institut National Agronomique Paris-Grignon, 78850
Thiverval-Grignon, France.

[⊥]Laboratoire de Génétique Moléculaire et Cellulaire, INRA, 78850 Thiverval-Grignon,
France.

^{II}This work was supported by grants from the Centre National de la Recherche Scientifique
(UPR 9063), and the Institut National de la Recherche Agronomique (UR 258 and UMR 206).

[∇]Atomic coordinates: pdb accession number 1KYA

*Corresponding author. e-mail: briozzo@grignon.inra.fr

Running title: Structure of a four-copper laccase with an arylamine

Abbreviations: ABTS: 2,2'-azino-bis(3-ethylbenzthiazoline-6-sulfonic acid); CPB: citrate/phosphate buffer; EPR: electron paramagnetic resonance; 2HF: *N',N'*-dimethyl-*N*-(2-hydroxyphenyl)urea; 2HF-4Cl: *N',N'*-dimethyl-*N*-(4-chloro-2-hydroxyphenyl)urea; 2HF-5Cl: *N',N'*-dimethyl-*N*-(5-chloro-2-hydroxyphenyl)urea; 4HF: *N',N'*-dimethyl-*N*-(4-hydroxyphenyl)urea; 4HF-5Cl: *N',N'*-dimethyl-*N*-(5-chloro-4-hydroxyphenyl)urea; HPLC: high performance liquid chromatography; IEF: Isoelectric focusing; Lac-Cc: the *Coprinus cinereus* laccase (pdb accession number 1A65); LacIIIb: the *Trametes versicolor* laccase (sequence accession number AF414109); MS: mass spectrometry.

ABSTRACT

Laccases are multi-copper oxidases that catalyze the oxidation of a wide range of phenols or arylamines, and their use in industrial oxidative processes is increasing. We purified from the white rot fungus *Trametes versicolor* a laccase which exists as five different isozymes, depending on glycosylation. The 2.4 Å resolution structure of the most abundant isozyme of the glycosylated enzyme was solved. The four copper atoms are present, and it is the first crystal structure of a laccase in its active form. The crystallized enzyme binds 2,5-xyldine, which was used as a laccase inducer in the fungus culture. This arylamine is a very weak reducing substrate of the enzyme. The cavity enclosing 2,5-xyldine is rather wide, allowing the accommodation of substrates of various sizes. Several amino-acid residues make hydrophobic interactions with the aromatic ring of the ligand. In addition, two charged or polar residues interact with its amino group. The first one is an histidine which also coordinates the copper that functions as the primary electron acceptor. The second is an aspartate conserved among fungal laccases. The purified enzyme can oxidize various hydroxylated compounds of the phenylurea family of herbicides, that we synthesized. These phenolic substrates have better affinities at pH 5 than at pH 3, which could be related to the 2,5-xyldine binding by the aspartate. This is the first structure of a multicopper oxidase complexed to a reducing substrate. It provides a model for engineering laccases either more efficient or with a wider substrate specificity.

Multicopper oxidases combine the four-electron reduction of dioxygen to water with the one-electron oxidation of four reducing substrate molecules. The well-defined members of this class are laccase, ascorbate oxidase and ceruloplasmin (for a review, see (1) and (2)). Laccases (*para*-diphenol:dioxygen oxidoreductases, EC 1.10.3.2) are generally glycosylated. Most are extracellular enzymes, and a given species may produce isozymes of both extra- and intracellular types. They are produced by various fungi, plants, and certain bacteria or insects. It has been proposed that plant laccases - in particular tree enzymes - catalyze during lignification the initial polymerization of monolignols by oxidatively coupling them into dimers and trimers. Then, peroxidases synthesize the extended polymeric lignins from oligolignols (1). In contrast, fungal laccases play a role in lignin degradation. Their *in vivo* functions also include fungal virulence, pigment formation and detoxification of phenols produced during lignin degradation (3).

Laccases contain four copper ions distributed into three sites, defined according to spectroscopic properties. The T1 site contains the type 1 blue copper (Cu1), whose tight coordination to a cysteine is responsible for an intense absorption band around 600 nm, giving its blue color to the enzyme. The T2 site contains a type 2 copper (Cu2) with a characteristic electron paramagnetic resonance (EPR). In the T3 site, the pair of strongly coupled type 3 coppers (Cu3a and Cu3b) is EPR-silent in the presence of dioxygen, indicative of a strongly antiferromagnetically coupled Cu pair bridged by a hydroxide. The mononuclear T1 site extracts electrons from the reducing substrate and mediates their transfer to the trinuclear T2/T3 center where molecular oxygen is reduced.

Laccases substrates are mostly phenols or arylamines. As these enzymes have a low substrate specificity, they can oxidize a variety of substrates, either of natural or industrial origin. Fungal laccases are marketed for bleaching in textile and dye industry. Other potential applications include paper industry, enzymatic conversion of chemical intermediates, or

oxidative transformation of environmental pollutants. Among the herbicides used for weed control, phenylurea derivatives have been extensively spread in the environment since their discovery in the 1950s. They are used both on cultivated (selective germination of cereal crops) and uncultivated (maintenance of roads, railways) areas. There is a growing concern over the potential of herbicides to contaminate soils, and surface or ground water. Among them, the chlorinated aromatic compounds tend to accumulate because of their low solubility in water. Moreover, several chlorophenols are carcinogenic. However, microorganisms can degrade them in soils. One possible initial step in this transformation pathway proceeds *via* the hydroxylation of the aromatic ring. This was reported in the case of the herbicide metoxuron (4), and also for the metabolisation of diuron (5) and monuron (6) by plants. In soils, photolysis of phenylurea herbicides exposed to sunlight can lead to ring hydroxylation for monuron (7), and in particular to the 2- and 4-hydroxylation of the aromatic ring for fenuron (8). Such a functionalization of the aromatic ring considerably extends the possible metabolic pathways. However, chlorinated hydroxyphenylurea herbicides are very difficult to degrade by aerobic microorganisms: these compounds are acutely toxic, as they uncouple oxidative phosphorylation.

White rot fungi are good candidates for further degradation of phenylureas in soil because they elude this toxicity problem. As most of their oxidative enzymatic systems involved in the transformation are extra-cellular, compounds do not need to be internalized prior to the degradation. These fungi can degrade a wide range of chlorinated phenols (9). In particular, laccases of *Trametes versicolor* have been reported to be involved in this process (10). We recently sequenced (accession number AF414109), purified and crystallized a laccase secreted by *T. versicolor* (11). Several sequences of laccases have been deposited for this basidiomycete. As the enzyme we studied shares 99.2 % sequence identity - only four residues are different - with the laccase III from *T. versicolor* (sequence accession number

D13372; (12)), we name it laccase IIIb (LacIIIb). Starting from such laccases, it would be of interest to design enzymes either more efficient in catalysis, or capable to oxidize a wider variety of reducing substrates.

However, there are not enough structural data to allow such an engineering, or even to understand laccase specificity and oxidation mechanism. The only known crystal structure of a laccase is that of the fungus *Coprinus cinereus* (Lac-Cc). In the two published models calculated either with data collected at room temperature (pdb code 1A65; (13)) or at 100 K (1HFU; (14)), the crystallized enzyme was trapped in a form devoid of the type 2 copper. This depletion completely inactivates laccases (15). Here we present the crystal structure of the active LacIIIb, allowing description of the trinuclear T2/T3 site. Furthermore, in all known structures of multicopper oxidases (ascorbate oxidase from Zucchini (16), human ceruloplasmin (17), and Lac-Cc) the putative cavity for the reducing substrate is empty. The structure presented here contains an arylamine, likely the laccase production inducer 2,5-xylydine (2,5-dimethylbenzenamine). This pinpoints key residues important for substrate binding.

As pointed above, the degradation of hydroxyphenylurea compounds by fungal laccases is of environmental interest. For this reason, and also to obtain additional information on enzyme/substrate interaction, we synthesized a series of isomers of such compounds. They differ by the respective positions of urea and hydroxyl substituents, and by the eventual presence of a chlorine substituent, on the aromatic ring. The kinetic parameters obtained for these derivatives are discussed in relation with the crystal structure.

EXPERIMENTAL PROCEDURES

Chemicals. Solvents used for high performance liquid chromatography (HPLC) analysis were purchased from Carlo Erba. 2,5-xylidine (2,5-dimethylbenzenamine), ABTS [2,2'-azinobis-(3-ethylbenzthiazoline-6-sulfonic acid)], and all chemicals and reagents used for the synthesis of hydroxyphenylurea compounds were purchased from Sigma-Aldrich, Acros, and Lancaster.

Production of the purified enzyme. The production and purification of LacIIIb have been described elsewhere (11). Briefly, the enzyme was produced from *T. versicolor* cultures induced by 2,5-xylidine. The purification included two steps of chromatography: firstly on a DEAE 52 anion exchange column, then on a hydrophobic interaction Phenyl sepharose column. This latter chromatography further improved the enzyme purity, as checked by sodium dodecyl sulfate-polyacrylamide gel electrophoresis, which gave a unique band.

Isoelectric focusing. Isoelectric focusing (IEF, (18)) was performed to check the eventual microheterogeneity of the enzyme. A vertical polyacrylamide Minigel system (Biorad) was utilized. The pH range of the ampholyte was 2.5 to 5 in a 5 % (w/v) polyacrylamide gel (prepared from Ready Mix IEF from Pharmacia). Proteins samples were mixed with the same volume of 60 % glycerol and 4 % Pharmalyte (Pharmacia). Focusing was performed at 200 V for one hour, then at 300 V for two additional hours. The resulting gel was rinsed and the pH gradient measured with a flat bottom pH electrode (Metrohm). The gel was then stained with 0.1 % Coomassie Blue R-250 in 10 % methanol and 0.5 % acetic acid for 10 min. Background destaining was performed by quick rinsing in 10 % methanol. Laccase activity of the different isoenzymes was assayed by incubating the gel in 50 mM citrate/phosphate buffer (CPB) pH 5 containing 1 % guaiacol (v/v). Enzymatically active protein bands appeared brown.

Crystallization, data collection and processing. All steps have been described elsewhere (11). Briefly, crystals were grown at 20°C in a sodium cacodylate buffer pH 6.5, using polyethylene glycol 8000 and zinc acetate as precipitants. A bundle of blue crystals appeared after a few days. Data were collected at 100K on a rotating anode generator, using glycerol as a cryoprotectant, and processed up to 2.4 Å resolution. Crystals belong to the monoclinic space group $P2_1$, with unit-cell dimensions $a=87.7$, $b=110.5$, $c=123.2$ Å, $\beta=103.4^\circ$. The unit cell contains four molecules per asymmetric unit.

Phasing, model building and structure refinement. The structure was solved by molecular replacement with AMoRe (19) in the resolution range 15.0-3.5 Å, using the laccase from *Coprinus cinereus* (pdb code 1A65) as the search model. This enzyme shares 57.7 % sequence identity with LacIIIb. The model included the entire polypeptide, without copper atoms or *O*-linked sugars. After rigid-body refinement, the correlation coefficient was 41.4 %, and the initial R factor 45.7 %.

Models were built using O (20). CNS (21) version 1.1 was used for refinement, using a maximum-likelihood strategy. After a few cycles of refinement, the initial electron density maps unambiguously revealed the presence of the four coppers in all molecules of the asymmetric unit. Clear density present in the active site was attributed to 2,5-xylydine. Bulk solvent correction and simulating annealing were used, as well as constraints corresponding to non-crystallographic symmetry (NCS). NCS was not applied to coppers, sugars and 2,5-xylydine. In the final steps, the model was refined using individual B factors. Water molecules were placed in residual density above 2.5 standard deviations, using the ‘*hbond*’ parameters of CNS. Side chains from a few residues had no clear density and were modeled as alanines. Details of the final model quality are given in Table 1.

Enzyme activity assays. Laccase activity was determined at 30°C by oxidation of ABTS into a stable cationic radical $ABTS^{\bullet+}$ (22) assayed at 420 nm. Ten microliters of the

enzyme solution were added to 990 μ l of 0.1 M CPB at pH 3 containing 1 mM ABTS and aerated by bubbling with air prior to the experiment to ensure oxygen saturation. In such conditions, an enzyme solution of one unit/ml was required to oxidize one micromole of ABTS per minute.

Synthesis of hydroxyphenylurea compounds. *N',N'*-dimethyl-*N*-(2-hydroxyphenyl)urea (2HF) and *N',N'*-dimethyl-*N*-(5-chloro-2-hydroxyphenyl)urea (2HF-5Cl) were prepared according to previously reported procedures (23), which describe the formation of 2-hydroxyphenylureas using a reaction between benzoxazolone (or corresponding 5-chloro substituted compounds) and alkylamines. *N',N'*-dimethyl-*N*-(4-chloro-2-hydroxyphenyl)urea (2HF-4Cl), *N',N'*-dimethyl-*N*-(4-hydroxyphenyl)urea (4HF) and *N',N'*-dimethyl-*N*-(5-chloro-4-hydroxyphenyl)urea (4HF-5Cl) were prepared according to a five-step procedure starting from a methyl hydroxybenzoate (23). This method was slightly adapted for the synthesis of the chlorinated compounds: the two last steps were replaced by a concomitant aminolysis reaction of both acetyl and isocyanate groups of the acetylchlorophenylisocyanates. This was achieved in an excess of liquid dimethylamine in either dry toluene or dichloromethane.

The structure of the compounds described here, and that of their precursors, were confirmed by ^1H - and ^{13}C -nuclear magnetic resonance spectroscopy, mass spectrometry and Infra-Red spectroscopy.

High performance liquid chromatography. HPLC analyses employed a Varian system including a 9010 pump and a UV-VIS 9050 detector. 20 μ l samples were analyzed using a reverse-phase Nova Pack C8 column (Waters). The flow rate throughout the analysis was 1 $\text{ml}\cdot\text{min}^{-1}$. UV absorbency of the eluent was measured at 254 nm. Several water/acetonitrile gradients were used:

Method I: hydroxyphenylurea derivatives were eluted by the following gradient: 0 to 3 min 100:0; 3 to 13 min ramped to 70:30; 13 to 14 min ramped to 0:100; 14 to 20 min held to 0:100.

Method II: the gradient for 2,5-xylydine analysis was: 0 to 15 min, 50:50; 15 to 17 min ramped to 0:100; 17 to 27 min, held at 0:100; 27 to 30 min, ramped to 50:50.

Determination of kinetic parameters of ABTS and hydroxyphenylurea derivatives.

Experiments were performed at 30°C in aerated 50 mM CPB at pH 3 and 5. Laccase concentrations, measured at 280 nm, ranged from 10^{-9} to 10^{-8} M in the assays, whereas substrate concentrations were in the range 0.01-2 mM. Time course of ABTS oxidation was followed for 2 min at 420 nm. Hydroxyphenylureas were incubated around 30 min in the presence of laccase and successive aliquots (0.9 ml) were removed. The substrate transformation was stopped by lowering the pH to 1-2 with 0.1 ml of 10 % trichloroacetic acid. Kinetic parameters were then determined using HPLC analysis (method I) to separate substrate and product compounds. In the range of substrate concentration used, the initial rates of hydroxylated phenylureas oxidation by laccase were consistent with Michaelis-Menten equation. The parameters obtained are listed in Table 2.

Transformation of 2,5-xylydine by LacIIIb. 2,5-xylydine (0.8 mM in 50 mM CPB pH 5, aerated prior to the experiment) was incubated with 0.25 unit of laccase at 30°C in darkness. A negative control without laccase was performed in parallel. Successive 1 ml aliquots were removed, filtered (Millipore 0.45 μ M) and analyzed by HPLC according to method II.

RESULTS

The global fold of the crystal structure. The structure is globular with approximate dimensions of 70 Å x 50 Å x 50 Å. It consists mainly of antiparallel β barrels (Fig. 1). There

are three cupredoxin-like domains: the T1 site (Cu1) belongs to one domain (lower part of Fig. 1), and the T2/T3 site (Cu2, Cu3a and Cu3b) is at the interface between the two other domains (upper part of Fig. 1). This topology is essentially the same as that from the monomer of Zucchini ascorbate oxidase (16). The overall structure is very close to that of the Cu2-depleted form of Lac-Cc (13). The electron density indicates glycosylation sites, and allows unambiguous determination of both the T1 and the trinuclear T2/T3 copper sites. As was described for ascorbate oxidase and Lac-Cc, there is a putative binding cavity for reducing substrates in the vicinity of Cu1. A clear electron density is present in this cavity, consistent with bound 2,5-xylydine.

The four N-glycosylation sites observed in the crystal structure. IEF experiments on the purified *T. versicolor* enzyme (Fig. 2) indicates five different isozymes. Their isoelectric points are rather close: 3.23 (for isozyme 1), 3.11 (isozyme 2), 3.0 (isozyme 3), 2.88 (isozyme 4), and 2.75 (isozyme 5). These laccase isozymes probably differ by their glycosylation. LacIIIb was expressed in fungus cultures, and not overexpressed in a different species (bacterium or yeast), and the purified enzyme was not deglycosylated before crystallization. This gave an opportunity to study its natural post-translational modifications. There are 6 putative *N*-glycosylation sites (consensus sequence Asn-X-Thr): asparagines 51, 54, 208, 217, 333 and 436. The density maps clearly indicates glycosylation on asparagines 54, 217, 333 and 436. In all cases the first *N*-acetyl-glucosamine could be modeled in the density. The glycosylation could be extended up to 4 supplementary osidic residues, depending on the site, and for a particular site on the molecule of the asymmetric unit.

The glycosylation observed in the crystal structure of LacIIIb is consistent with mass spectrometry experiments. Isozymes 1 and 2 of LacIIIb, the most abundant forms, were separated by cutting IEF bands, and proteolysed by trypsin. The resulting fragments were

attributed on the basis of their molecular weights in MALDI-TOF mass spectrometry (not shown). Fragments that could not be attributed suggested a post-translational modification. For instance, the glycosylation of Asn 436 explains why the expected 424-440 fragment remains absent in mass spectrometry experiments for isozyme 1. In contrast, this fragment is found with the isozyme 2. This suggests that the crystallized enzyme, glycosylated on Asn 436, corresponds to the isozyme 1, the most abundant one.

The trinuclear T2/T3 copper site. The trinuclear center is coordinated to eight histidines (Fig. 3a). Six of them coordinate Cu3a and Cu3b: histidines 66, 109 and 454 for Cu3a, histidines 111, 400 and 452 for Cu3b. In addition, the bridging oxygen atom from the hydroxide known to bind both Cu3a and Cu3b is equidistant from these two coppers. The oxygen-Cu3a distance is 1.8 to 1.9 Å, depending on the molecule of the asymmetric unit, and the oxygen-Cu3b distance is 1.8 to 2.0 Å. This results in a Cu3a-Cu3b distance of 3.75 Å. Cu2 is coordinated to histidines 64 and 398, and to a water molecule from the solvent. The Cu2-water distance varies notably among the four different molecules of an asymmetric unit (2.3 to 2.9 Å).

The culture inducer 2,5-xylidine is present in the cavity close to Cu1. The electron density indicates the presence of a ligand molecule in the putative cavity for the reducing substrate. Figure 4a shows that the omit map is in good accordance with the model of 2,5-xylidine. This arylamine was initially used in the fungus culture medium for laccase induction.

Given its presence at the active site, we checked if 2,5-xylidine could be a laccase substrate (Fig. 4b). After 20 hours of incubation in the presence of laccase (pH 5, 30°C), 38 % of the 2,5-xylidine was transformed, and significant new peaks appeared. The products corresponding to these new peaks had higher retention times, indicating that they were more hydrophobic than the arylamine substrate.

The T1 copper site and the neighboring cavity that binds 2,5-xylydine. Cu1 is coordinated to histidines 395 and 458, and to Cys 453 (Fig. 3b). The non coordinated Phe 463 residue is distant of 3.7 Å from the copper. This phenylalanine is homologous to the axial methionine coordinated to the copper of the T1 site in ascorbate oxidase. In LacIIIb structure, the histidine ring of His 458 (which is coordinated to Cu1) is close to the nitrogen from 2,5-xylydine: distances between enzyme and substrate nitrogens are less than 3.2 Å, consistent with a hydrogen bond, for all molecules of an asymmetric unit, except in molecule C. In this latter case, the distance is 4 Å.

The amino group of 2,5-xylydine is also H-bonded to a terminal oxygen of Asp 206 side chain. In addition, for molecule C (in which His 458 is not H-bonded to the arylamine) the second terminal oxygen is also H-bonded to 2,5-xylydine, making a bidentate interaction. Beside these hydrogen bonds, there are many hydrophobic protein-ligand interactions (Fig. 5, dark blue). They involve Phe 162, Leu 164 and Phe 265 residues, all belonging to two neighboring loops (above 2,5-xylydine in the orientation of Fig. 5a) which delineate the cavity. On the opposite side (below the ligand on Fig. 5a), two hydrophobic residues (phenylalanines 332 and 337) constitute the other wall of the cavity. This cavity is rather wide (approximate dimensions 10 x 10 x 20 Å), so that the observed ligand is not tightly buried in it.

Activity of the purified enzyme. Laccase-catalyzed oxidation of hydroxyphenylureas derivatives and of the non phenolic model substrate ABTS (shown on Fig. 6) was examined at pH 3 and 5. All kinetic parameters of hydroxyphenylureas are in the same order of magnitude: K_m values range from 0.22 to 4.4 mM, and k_{cat} from 32 to 290 s⁻¹ (Table 2). The highest oxidation rate (based on k_{cat}/K_m values) was for 4HF-5Cl at pH 5, the lowest for 2HF and 4HF at pH 3. Increasing the pH from 3 to 5 induces a significant increase in the affinity for each of the hydroxyphenylurea derivatives. As phenols are natural substrates for laccase, it

can be supposed that the nature and position of substituents on the phenolic ring is a key element of substrate docking in the active center. Among the hydroxyphenylureas studied, the derivative with a *para*-hydroxyl substituent, 4HF, shows the lowest affinity (specially at pH 3, with a K_m value of 4.4 mM), although this substitution position is expected to induce the weakest steric hindrance, as referred to the phenolic group. Actually, both at pH 3 and 5, the derivatives with an *ortho* substituent (as referred to the hydroxyl) have relatively good affinities, whatever the substituent is an urea (2HF has a lower K_m than 4HF) or a chlorine (4HF-5Cl has a lower K_m than 4HF). Such a result was also observed by Xu (24): all *ortho*-substituted phenols exhibited a better affinity than phenol itself for the recombinant laccase from *Trametes villosa*. At pH 3, the addition of a chlorinated substituent to the aromatic ring induces a positive effect on the binding affinity of the substrate. The largest effect is observed for 4HF-5Cl *versus* 4HF.

DISCUSSION

The trinuclear T2/T3 copper site: why is Cu2 labile? In the previously published structure of *C. cinereus* laccase, the extremely high mosaicity of crystals grown from the native enzyme prevented X-ray studies. Therefore, in order to get crystals suitable for data collection, enzymatic deglycosylation was necessary, but it caused the loss of Cu2 (25). On the opposite, with *T. versicolor* laccase a complete data set could be collected at 2.4 Å with the native enzyme, despite a high mosaicity of the crystals. This allows for the first time a complete description of the trinuclear copper center.

The distance between the two coppers of the T3 site is 3.75 Å. This is close to values observed for other multi-copper oxidases of known three-dimensional structure (around 3.7 Å for both ascorbate oxidase (16) and ceruloplasmin (17)). However, in the structure of Lac-Cc, the hydroxide bridging Cu3a no longer bridged Cu3b, and the Cu3a-Cu3b distance was larger (around 5 Å, (13), (14)). As was suggested by the authors, this longer distance probably reflects the absence of Cu2. Beside Cu2 depletion, modifications of the enzyme oxidation state can also increase the Cu3a-Cu3b distance and alter the binding of the oxygen bridging the Cu3 atoms. For instance in the case of ascorbate oxidase the Cu3a-Cu3b distance increases from 3.74 Å for the natural, oxidized form (16) to values ranging from 4.8 to 5.1 Å for the reduced, peroxide and azide forms of the enzyme, where no oxygen bridges Cu3a and Cu3b (26).

In *T. versicolor* laccase, either Cu3a or Cu3b is bound to three histidines and an oxygen atom, whilst Cu2 is coordinated to only two histidines, and to a water molecule. The same feature was found for the trinuclear copper site of ascorbate oxidase (pdb code 1AOZ), in which Cu2 is depletable. Moreover, in LacIIIb structure the position of the coordinated water molecule significantly varies among the four different molecules of the asymmetric unit, indicating that Cu2 is only weakly coordinated to this water. This gives a structural basis to the lability of Cu2.

The presence of 2,5-xylydine bound to the crystallized enzyme. An electron density consistent with 2,5-xylydine was found in the cavity close to the T1 copper site, indicating the presence of this arylamine in the crystals. Most inducers of fungal laccases are substrates of the enzyme to be produced, or substances structurally similar to these substrates. 2,5-xylydine, a usual laccase inducer (27), was used in the culture medium for *T. versicolor*. Incubation of 2,5-xylydine with the purified enzyme showed that this arylamine is a substrate, although very poor: comparative experiments were performed at pH 5 and 30°C with a same amount of

laccase (0.25 U/ml) and of the different substrates (0.8 mM). In such conditions, no significant transformation of 2,5-xylydine (less than 5 %) occurs after 4 hours, whereas both 2HF-5Cl and ABTS are completely transformed in less than ten minutes. It must also be kept in mind that the conditions chosen to test laccase-catalyzed oxidation of 2,5-xylydine were the most favorable. In contrast, in the mother liquor used for crystallization, the controlled temperature was 20°C. Moreover, the pH was 6.5, a value for which LacIIIb is almost inactive with hydroxyphenylurea substrates (not shown). Thus, the transformation of 2,5-xylydine should be very slow in the crystallization drop. Otherwise, the crystallization process could have selected a small subpopulation of the enzyme that bound 2,5-xylydine. In this case, the substrate would only be present at a low level in the purified enzyme preparation.

In the small electron-transporting cupredoxins, a main chain carbonyl oxygen coordinates the axial position of Cu1. An equivalent carbonyl is also present in multicopper oxidases, but is displaced away from the T1 site by a main chain rearrangement, resulting in the formation of a putative binding cavity for the reducing substrate (13). The structure of LacIIIb complexed to 2,5-xylydine is the first direct experimental evidence that the cavity close to Cu1 is indeed the substrate binding cavity.

Contribution to electron transfer analysis. The catalytic activity of laccases implies an electron transfer from Cu1 to the trinuclear T2/T3 site. This transfer may be through-bond, through-space or a combination of both. The LacIIIb structure provides two branched through-bond pathways starting from Cu1, both involving Cys 453 and a contiguous histidine. The first one leads to Cu3a through His 454, and the second one to Cu3b through His 452 (see Fig. 3). In the alternate through-space theory, it has been calculated that electron tunneling in proteins between redox centers that are distant of 14 Å or less does not imply the need for an optimal route through residues selected by the evolution (28). In *T. versicolor* laccase the

distance between Cu1 and Cu3a is 12.8 Å, that between Cu1 and Cu3b is 12.2 Å. Therefore, the structure is also consistent with a through-space transfer.

Concerning the electron transfer associated with the reduction of Cu1 by the substrate, the structure presented here gives information (Fig. 3b). The proximity of His 458, which coordinates Cu1, with the amino group of 2,5-xyldine, from which an electron is extracted by the enzyme, suggests that His 458 is the entrance door of this electron during its transfer to Cu1. This could explain the importance of site-directed mutagenesis of the contiguous sequence LEA (Leu 459-Glu 460-Ala 461 in LacIIIb) for the catalytic activity and pH dependence of fungal laccases (29). The Cu1-coordinating His 458 is highly conserved in all multicopper oxidases.

The cavity for the reducing substrate: possibility of a ligand-induced closure. The backbones of *T. versicolor* and *C. cinereus* laccases are extremely similar, except for the longer C-terminal region of Lac-Cc (Fig. 5a). The only significant difference involves the external loops surrounding the 2,5-xyldine binding site: the cavity is more closed in the case of LacIIIb. There are three possible explanations: i) an artefactual difference due to the different crystal packings of LacIIIb and Lac-Cc, possibly induced by the different glycosylation levels; ii) a species variation related to the sequence difference between *T. versicolor* and *C. cinereus* laccases; iii) an induced-fit closing movement of LacIIIb due to the presence of a bound ligand. At present, we have no way to suggest which explanation is the right one.

The cavity for the reducing substrate appears far more surrounded by hydrophobic side chains in LacIIIb than in Lac-Cc. The hydrophobic residues Leu 164 and Phe 265 of *T. versicolor* laccase (above 2,5-xyldine in the orientation of Fig. 5a) replace the corresponding polar (Gln 164) and charged (Lys 264) residues of *C. cinereus* enzyme. These two residues belong to loops which are closer in LacIIIb structure than in that of Lac-Cc. Moreover, the

hydrophobic residue Phe 162 of LacIIIb is larger, and closer from the 2,5-xylydine position, than the corresponding Ala 162 from *C. cinereus* enzyme (Fig. 5b). The opposite side of the ligand (below it on Fig. 5a) also shows a loop that is closer for LacIIIb. This loop contains the phenylalanines 332 and 337, which make hydrophobic interactions with the ligand, and are conserved in both enzymes. To summarize, the loops of *T. versicolor* laccase that appear more closed upon the substrate cavity contain 2,5-xylydine interacting residues. This is consistent with a possible ligand-induced fit of laccases.

Correlation of the structure with the pH dependence of hydroxyphenylureas affinities. For all five synthesized hydroxyphenylurea compounds the affinities reflected by K_m values are significantly better at pH 5 than at pH 3 (Table 2). No chemical deprotonation of the phenolic group of these substrates is expected in the 3-5 pH range. Thus, the observed effect of pH can be related to pK_a values of laccase side chains interacting with the oxidized substrate substituent: an amino substituent for 2,5-xylydine and other arylamines, an hydroxyl substituent for phenolic substrates. The structure indicates that the amino substituent of 2,5-xylydine is H-bonded to Asp 206 in all four molecules of the asymmetric unit. The typical pK_a value for an aspartate side chain is 3.9, allowing different protonation states for pH 3 and pH 5. At pH 5 the carboxylic function of Asp 206 is dissociated, so that its negative charge should stabilize the partial positive charge of the hydrogen extracted from the reducing substrate by the enzyme. Conversely, at pH 3 the side chain of Asp 206 is not dissociated, which is less favorable for the reduction process.

With ABTS there is no significant difference in K_m values between pH 3 and pH 5. ABTS differs from phenolic substrates because its oxidation does not involve proton transfer, but only a single electron transfer. Thus, as with ABTS no hydrogen can be extracted from the reducing substrate, the pH-dependent stabilizing role of Asp 206 is not observed. This is

consistent with the proposed essential role of this residue in the laccase-catalyzed oxidation of arylamine or phenol substrates. Asp 206 is well conserved among fungal laccases. It is also contiguous to Cys 205, a residue involved in a disulfide bond conserved in fungal laccases.

ACKNOWLEDGMENTS

We thank Joël Janin (CNRS UPR 9063) for helpful discussion about the manuscript.

REFERENCES

1. Solomon, E. I., Sundaram, U. M., and Machonkin, T. E. (1996) *Chem. Rev.* 96, 2536-2605.
2. McGuirl, M. A., and Dooley, D. (1999) *Curr. Opin. Chem. Biol.* 3, 138-144.
3. Gianfreda, L., Xu, F., and Bollag, J. M. (1999) *Bioremed. J.* 3, 1-25.
4. Briggs, G. G., and Ogilvie, S. Y. (1971) *Pestic. Sci.* 2, 165-168.
5. Roberts, T. R., Hutson, D. H., Lee, P. W., Nicholls, P. H., and Plimmer, J. R. (1998) in *Metabolic pathways of agrochemicals* (The Royal Society of Chemistry, Ed.), pp. 705-772, Cambridge, UK.
6. Frear, D. S., and Swanson, H. R. (1974) *Phytochemistry* 13, 357-360.
7. Tanaka, F. S., Wien, R. G., and Zaylskie, R. G. (1977) *J. Agric. Food Chem.* 25, 1068-1072.
8. Aguer, J. P., and Richard, C. (1996) *Pestic. Sci.* 46, 151-155.
9. Cameron, M. D., Timofeevski, S., and Aust, S. D. (2000) *Appl. Microbiol. Biotechnol.* 54, 751-758.
10. Ullah, M. A., Bedford, C. T., and Evans, C. S. (2000) *Appl. Microbiol. Biotechnol.* 53, 230-234.
11. Bertrand, T., Jolival, C., Caminade, E., Joly, N., Mougin, C., and Briozzo, P. (2002) *Acta Crystallog. sect. D* 58, 319-321.
12. Mikuni, J., and Morohoshi, N. (1997) *FEMS Microbiol. Lett.* 155, 79-84.
13. Ducros, V., Brzozowski, A. M., Wilson, K. S., Brown, S. H., Ostergaard, P., Schneider, P., Yaver, D. S., Pedersen, A. H., and Davies, G. J. (1998) *Nat. Struct. Biol.* 5, 310-316.

14. Ducros, V., Brzozowski, A. M., Wilson, K. S., Ostergaard, P., Schneider, P., Svendsen, A., and Davies, G. J. (2001) *Acta Crystallog. sect. D* 57, 333-336.
15. Reinhammar, B., and Oda, Y. (1979) *J. Inorg. Biochem.* 11, 115-27.
16. Messerschmidt, A., Ladenstein, R., Huber, R., Bolognesi, M., Avigliano, L., Petruzzelli, R., Rossi, A., and Finazzi-Agro, A. (1992) *J. Mol. Biol.* 224, 179-205.
17. Zaitseva, I., Zaitsev, V., Card, G., Moshkov, K., Bax, B., Ralph, A., and Lindley, P. (1996) *J. Biol. Inorg. Chem.* 1, 15-23.
18. Robertson, E. F., Dannelly, H. K., Malloiy, P. J., and Reeves, H. C. (1987) *Anal. Biochem.* 167, 290-294.
19. Navaza, J. (1994) *Acta Crystallog. sect. A* 50, 157-163.
20. Jones, T. A., Zou, J. Y., Cowan, S. W., and Kjeldgaard, M. (1991) *Acta Crystallog. sect. A* 47, 110-119.
21. Brünger, A. T., Adams, P. D., Clore, G. M., DeLano, W. L., Gros, P., Grosse-Kunstleve, R. W., and al., e. (1998) *Acta Crystallog. sect. D* 54, 905-921.
22. Wolfenden, B. S., and Wilson, R. L. (1982) *J. Chem. Soc. Perkin Trans. 2*, 805-812.
23. Jolival, C., Raynal, A., Caminade, E., Kokel, B., Le Goffic, F., and Mougin, C. (1999) *Appl. Microbiol. Biotechnol.* 51, 676-681.
24. Xu, F. (1996) *Biochemistry* 35, 7608-7614.
25. Ducros, V., Davies, G. J., Lawson, D. M., Wilson, K. S., Brown, S. H., Ostergaard, P., Pedersen, A. H., Schneider, P., Yaver, D. S., and Brzozowski, A. M. (1997) *Acta Crystallog. sect. D* 53, 605-607.
26. Messerschmidt, A., Luecke, H., and Huber, R. (1993) *J. Mol. Biol.* 230, 997-1014.
27. Fahraeus, G., Tullander, V., and Ljunggren, H. (1958) *Physiol. Plant.* 11, 631-643.
28. Page, C. C., Moser, C. C., Chen, X., and Dutton, L. (1999) *Nature* 402, 47-52.

29. Xu, F., Berka, R. M., Wahleithner, J. A., Nelson, B. A., Shuster, J. R., Brown, S. H., Palmer, A. E., and Solomon, E. I. (1998) *Biochem. J.* 334, 63-70.
30. Laskowski, R. A., MacArthur, M. W., Moss, D. S., and Thornton, J. M. (1993) *J. Appl. Crystallog.* 24, 946-950.
31. Kraulis, P. (1991) *J. Appl. Crystallog.* 24, 946-950.
32. Merritt, E. A., and Bacon, D. (1997) *Meth. Enzymol.* 277, 505-524.
33. Roussel, A., and Cambillau, C. (1989) *Silicon Graphics Geometry Partners Directory*, 77-78.
34. Esnouf, R. M. (1997) *J. Mol. Graph.* 15, 132-134.

Table 1: Refinement statistics

Resolution range (•)	35-2.4
R _{cryst} ^a (%)	25.3
R _{free} ^b (%)	27.6
Residues	1-499
Number of water molecules	791
R.m.s. deviations	
bond lengths (Å)	0.0075
bond angles (°)	1.47
Average B factor (• ²)	
protein	37.6
Cu (T1, T3)	46.8
Cu (T2)	74.6
2,5-xylidine	50.5
sugars	70.2
water molecules	42.1
Ramachandran statistics ^c (%)	
most favored regions	83.4
additional allowed regions	16.6

^a $R_{\text{cryst}} = \Sigma ||\text{Fobs}| - |\text{Fcalc}|| / \Sigma |\text{Fobs}|$, where |Fobs| and |Fcalc| are the observed and calculated structure factor amplitudes, respectively.

^b R_{free} is the same as R_{cryst} but calculated with a 5 % subset of all reflections that was never used in crystallographic refinement.

^c As evaluated by PROCHECK (30).

Table 2: Kinetic parameters of the purified LacIIIb with different substrates

substrate	K_m (mM)		k_{cat} (s^{-1})	
	pH 3	pH 5	pH 3	pH 5
ABTS	0.060	0.085	220	64
2HF	1.8	0.23	38	32
2HF-4Cl	1.6	0.38	290	140
2HF-5Cl	1.2	0.24	87	63
4HF	4.4	0.6	72	47
4HF-5Cl	0.74	0.22	82	97

FIGURE 1: The global fold of LacIIIb.

Ribbon representation showing the secondary structures. α helices are coloured dark blue, β strands are magenta. Copper atoms are indicated as cyan spheres, and labelled: 1 for the “blue” copper of the T1 site, 2 for the depletable copper of the T2 site, 3a and 3b for the coupled coppers of the T3 site. The four glycosylation sites are indicated by circles (full when sites are apparent in the orientation of the figure, empty when they are hindered). 2,5-xylydine is indicated in ball-and-stick with green sticks (carbon atoms are dark grey, the nitrogen atom is blue). Drawn with Molscript (31) and Raster3D (32).




FIGURE 2: Isoelectric focusing of the purified laccase.

Purified laccase samples (100 μ g per lane) were analyzed in the pH range from 2.5 to 5. Lane 3 contains 20 μ g of Sigma IEF standard protein mix. The two first lanes were stained with guaiacol to reveal laccase activity. Lanes 3 to 5 were stained for proteins with Coomassie R-250 blue. Lanes 1 and 4 show a laccase sample after the first purification step on a DEAE 52 column. In lanes 2 and 5, the sample was further chromatographed on a Phenyl Sepharose column.




FIGURE 3: The copper sites of LacIIIb and their environment.

a) The trinuclear T2/T3 site. Histidine side chains involved in copper binding (copper atoms are blue) are in sticks (C atoms are yellow, N atoms are cyan), and labelled. Oxygen atoms involved in copper coordination are red, and their label in parentheses indicates the corresponding molecule. Copper coordinations are indicated with dark lines. Dotted lines indicate interacting atoms distant from less than 3.2 Å. Drawn with Turbo (33).

b) The T1 site and the neighboring 2,5-xylydine ligand.

The side chains of the copper-coordinating histidines and cysteine are drawn. Also indicated are the side chains of the non coordinating Phe 463, and of Asp 206 residue that interacts with 2,5-xylydine (green model).





FIGURE 4: The 2,5-xylydine present in the putative reducing substrate cavity is a laccase substrate.

a) Electron density of the 2,5-xylydine ligand. Fo-Fc omit map (magenta) calculated in the absence of the ligand. The contour level is at 3 standard deviations of the map. The refined model of 2,5-xylydine is shown. The figure was prepared with Bobscript (34) and Raster3D.

b) Transformation of 2,5-xylydine by laccase at pH 5. HPLC analysis (method II). Peak A corresponds to 2,5-xylydine at the beginning of experiment. Peak B is observed after 20 hours incubation with laccase. Peak C is the negative control (without enzyme) after 20 hours incubation.




FIGURE 5: Superposition of the laccases from *Trametes versicolor* (LacIIIb) and from *Coprinus cinereus* (Lac-Cc).

a) Overall view. The LacIIIb backbone is represented as a thick dark blue trace, that of Lac-Cc as a thin red trace. Coppers are cyan for Lac-Cc, and yellow for LacIIIb. The positions of the residues interacting with 2,5-xylydine are marked with full circles, and labelled. Drawn with Molscrip and Raster3D.

b) Magnification of the 2,5-xylydine and neighboring interacting residues of the reducing substrate cavity. Side chains of LacIIIb residues are dark blue. Corresponding residues of Lac-Cc are red. Drawn with Turbo.




FIGURE 6: Schematic representation of LacIIIb substrates assayed for kinetic parameters determination. Drawn with ChemDraw.

