



HAL
open science

Nucleotide diversity of the ZmPox3 maize peroxidase gene: relationships between a MITE insertion in exon 2 and variation in forage maize digestibility

Carine Guillet-Claude, Christelle Birolleau-Touchard, Doménica Manicacci, Peter Rogowsky, Joan Rigau, Alain Murigneux, Jean-Pierre Martinant, Yves Y. Barrière

► To cite this version:

Carine Guillet-Claude, Christelle Birolleau-Touchard, Doménica Manicacci, Peter Rogowsky, Joan Rigau, et al.. Nucleotide diversity of the ZmPox3 maize peroxidase gene: relationships between a MITE insertion in exon 2 and variation in forage maize digestibility. *BMC Genetics*, 2004, 5, pp.19. 10.1186/1471-2156-5-19 . hal-02678277

HAL Id: hal-02678277

<https://hal.inrae.fr/hal-02678277>

Submitted on 31 May 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Research article

Open Access

Nucleotide diversity of the *ZmPox3* maize peroxidase gene: Relationships between a MITE insertion in exon 2 and variation in forage maize digestibility

Carine Guillet-Claude*^{1,2}, Christelle Birolleau-Touchard²,
Domenica Manicacci³, Peter M Rogowsky⁴, Joan Rigau⁵, Alain Murigneux⁶,
Jean-Pierre Martinant⁶ and Yves Barrière²

Address: ¹Centre de recherche en Biologie Forestière, Université Laval, Sainte-Foy, Québec, G1K 7P4, Canada, ²Unité de Génétique et d'Amélioration des Plantes Fourragères, INRA, Route de Saintes, Lusignan, F86600, France, ³Station de Génétique Végétale, UMR INRA-UPS-CNRS-INAPG, Ferme du Moulon, Gif sur Yvette, F91190, France, ⁴Reproduction et Développement des Plantes, UMR 5667 INRA-CNRS-ENSL-UCBL, IFR128 BioSciences Lyon-Gerland, ENS-Lyon, 46 allée d'Italie, Lyon, F69364, France, ⁵Departamento Genética Molecular, Institut de Biologia Molecular de Barcelona, Consejo Superior d'Investigaciones Científicas, Jordi Girona 18–26, Barcelona, 08034, Spain and ⁶Biogemma S.A.S., Campus Universitaire des Cézeaux, 24 avenue des Landais, Aubiere, F63170, France

Email: Carine Guillet-Claude* - cguillet@rsvs.ulaval.ca; Christelle Birolleau-Touchard - birollea@mons.inra.fr;
Domenica Manicacci - manicacc@moulon.inra.fr; Peter M Rogowsky - peter.rogowsky@ens-lyon.fr; Joan Rigau - jrlgmr@ibmb.csic.es;
Alain Murigneux - alain.murigneux@biogemma.fr; Jean-Pierre Martinant - jean-pierre.martinant@biogemma.fr;
Yves Barrière - barriere@lusignan.inra.fr

* Corresponding author

Published: 16 July 2004

Received: 03 June 2004

BMC Genetics 2004, 5:19 doi:10.1186/1471-2156-5-19

Accepted: 16 July 2004

This article is available from: <http://www.biomedcentral.com/1471-2156/5/19>

© 2004 Guillet-Claude et al; licensee BioMed Central Ltd. This is an Open Access article: verbatim copying and redistribution of this article are permitted in all media for any purpose, provided this notice is preserved along with the article's original URL.

Abstract

Background: Polymorphisms were investigated within the *ZmPox3* maize peroxidase gene, possibly involved in lignin biosynthesis because of its colocalization with a cluster of QTL related to lignin content and cell wall digestibility. The purpose of this study was to identify, on the basis of 37 maize lines chosen for their varying degrees of cell wall digestibility and representative of temperate regions germplasm, *ZmPox3* haplotypes or individual polymorphisms possibly associated with digestibility.

Results: Numerous haplotypes with high diversity were identified. Frequency of nucleotide changes was high with on average one SNP every 57 bp. Nucleotide diversity was not equally distributed among site categories: the estimated π was on average eight times higher for silent sites than for non-synonymous sites. Numerous sites were in linkage disequilibrium that decayed with increasing physical distance. A *zmPox3* mutant allele, carrying an insertion of a transposable element in the second exon, was found in lines derived from the early flint inbred line, F7. This element possesses many structural features of miniature inverted-repeat transposable elements (MITE). The mutant allele encodes a truncated protein lacking important functional sites. An ANOVA performed with a subset of 31 maize lines indicated that the transposable element was significantly associated with cell wall digestibility. This association was confirmed using an additional set of 25 flint lines related to F7. Moreover, RT-PCR experiments revealed a decreased amount of corresponding mRNA in plants with the MITE insertion.

Conclusion: These results showed that *ZmPox3* could possibly be involved in monolignol polymerisation, and that a deficiency in *ZmPox3* peroxidase activity seemingly has a negative effect on cell wall digestibility. Also, genetic diversity analyses of *ZmPox3* indicated that this peroxidase could be a relevant target for grass digestibility improvement using specific allele introgressions.

Background

Because of its high energy content and good ingestibility, silage maize is a major source of forage for dairy cattle. Its energy value depends mostly on the breakdown of cell walls by rumen micro-organisms. Lignin interferes with the digestion of cell wall polysaccharides by acting as a physical barrier to microbial enzymes. In forage crops, lignin content, lignin structure and cross-linking between cell wall components influence digestibility [1].

Lignins are formed through dehydrogenative polymerization of two or three monolignols. Grass lignins are primarily made of two major units; guaiacyl (G, monomethoxylated on the aromatic ring) and syringyl units (S, di-methoxylated), resulting of the polymerization of coniferyl and sinapyl alcohols, respectively. The last step in lignin biosynthesis transport of the monolignols to the cell wall thus involves the oxidation of monolignols within the plant cell wall matrix. The dehydrogenation to monolignols radicals is attributed to different classes of enzymes, such as peroxidases (oxidoreductases, EC 1.11.1.7), laccases (oxidoreductases, EC 1.10.3.2) and (poly)phenol oxidases [2,3]. However, the effects of modifying these enzymes on cell wall lignification and digestibility have not been investigated extensively in grasses. In tobacco, down-regulation of a peroxidase led to transformants with a reduced lignin content of up to 40–50 % [4]. Down-regulation of an anionic peroxidase in transgenic aspen was associated with lower lignin content and modified lignin composition [5].

In plants, peroxidases belong to a multigene family. In maize, 13 peroxidase genes are described in The Maize Genetics and Genomics Database (<http://www.maizegdb.org> [6]). Three maize cDNAs coding for three different peroxidases (*ZmPox1*, *ZmPox2* and *ZmPox3*) were isolated from a 9-day old tip-less root library [7]. Expression patterns of these peroxidases suggested that only *ZmPox2* and *ZmPox3* were involved in lignification. *ZmPox3* was mapped by Biogemma SAS (A Murigneux, pers. com.) in the bins 6.05/07, and therefore co-localizes with a major QTL cluster for lignification and cell wall digestibility [8,9].

The breeding of forage crops for higher digestibility can involve specific genetic resources devised through genetic engineering of lignin biosynthesis [1]. In addition, the identification of natural allelic forms of candidate genes that correlate with higher digestibility, and their subsequent use in marker assisted selection (MAS) schemes, may become an interesting breeding tool. Gene sequence diversity has been primarily studied to understand the impact of selection during maize domestication and to identify the levels and patterns of genetic variation in a large sample of maize loci [10-12]. However, Thornsberry

et al [13] and Palaisa *et al* [14] demonstrated that nucleotide diversity analysis of candidate genes for flowering date and endosperm colour, respectively, allowed for the identification of alleles responsible for variation of these quantitative agronomic traits. Recently, in *Eucalyptus globulus*, Poke *et al* [15] found polymorphisms which could alter enzyme function in cinnamoyl CoA reductase (*CCR*) and cinnamyl alcohol dehydrogenase 2 (*CAD2*).

Allelic variation of the *ZmPox3* peroxidase was studied in a set of 37 inbred lines or ecotypes representative of germplasm currently used in breeding programs and displaying a large range of variation in cell wall digestibility. A *zmPox3* mutant allele, carrying an insertion of a transposable element in the second exon, was found in five lines and one ecotype. This element possesses many features of the MITE (miniature inverted repeat transposable element) class of mobile element [16]. This MITE insertion was further investigated to elucidate the relationship between this peroxidase mutant allele and cell wall digestibility.

Results

ZmPox3 gene diversity

The *ZmPox3* gene, amplified for 37 diverse maize inbred lines or ecotypes (Table 1), is approximately 1.7 kb long, and the coding region contains two small introns of about 127 and 111 bp (Fig. 1A). The number of polymorphic sites and indel events for the complete *ZmPox3* maize gene is reported in Table 2, along with two estimators of nucleotide diversity π [17] and θ [18]. Values are given for the complete sequence, as well as 3'UTR, introns, silent, synonymous and non-synonymous sites. Fourteen diverse haplotypes were found. There were 31 SNPs (i.e. an average of one SNP every 57 bp), and 17 indels which representing 20% of the whole sequence length. Of the 17 indels identified, 29% were single-bp, 23% were double-bp and 11% contained three-bp. Nucleotide diversity was not equally distributed among site categories (Table 2). Non-coding regions had a higher diversity, especially in the introns as compared to the coding region, where non-synonymous sites showed very low diversity. The estimated π value was on average eight times higher for silent sites than for non-synonymous sites. Introns were particularly rich in SNPs and indels. Intronic regions had an average of one SNP every 14 bp while exons had only one SNP every 82 bp. The two *ZmPox3* introns have 15 indels varying in size from 1 to 12 bp, while the 3'UTR region has 3 indels with size ranging from 2 to 5 bp.

For five inbred lines (F7, and its progenies F226, F227, F324 and F7012) and one ecotype (Québec28), the only indel found in the coding region was identified in the second exon of the *ZmPox3* gene. This insertion corresponded to a transposable element sharing many features

Table 1: Maize germplasm used in the association study.

Line	Origin and Pedigree	Cell wall digestibility
Flint		
F1	Lacaune (France)	3.5
F2	Lacaune (France)	3.5
F7	Lacaune (France)	4
F226	Du101 × F7	4
F227	F7 × F120	3.5
F7012	F226 × F227	4.5
F4	Etoile de Normandie (France)	5
F66	Sost (France)	3
Ep1	Lizaragote (Spain)	4
F120	F22 × Ep1 ³ (with Fc22 from Chavannes)	3
Du101	related to Jaune de Bade	3.5
F64	Argentina P1186-223	4
F564	F7 × F64	4.5
F286	F7 × F564	4
F324	related to F226, F227 and F286	5
Line212	Private Biogemma line	1
Line16	Private Biogemma line	3.5
Early dent		
F113	Spooner473 = (W37a × W37) × (W47 × EK43)	4
W117	W643 (from Golden Krug) × Minnesota13	4
F271	Co125 (unknown origin) × W103 (from Golden Glow)	1
F288	related to US early dent (7/8) and Blanc de Chalosses (1/8)	3
MBS847	Iodent	2.5
F7025	Iodent × F113	2.5
Late dent		
B14	Iowa Stiff Stalk Synthetic (C1)	1.5
B73	Iowa Stiff Stalk Synthetic (BS13C5)	1.5
Mo17	Lancaster	2.5
Lan496	Lancaster related	2
W64A	Wf9 (from Reid Yellow dent) × CI 187-2 (from Krug Reid)	2.5
Wis93-3520	Wisconsin dent material	4
Wis94-443	Wisconsin dent material	4
DE811	Related to US late dent (BSSS, Lancaster, ...)	1.5
Ecotypes		
	Rottaler silomais (Flint, Germany)	-
	Noordlander VC145 (Flint × dent, The Netherlands)	-
	Sibiriacka (Flint, Russia)	-
	Rainbow flint (Canada)	-
	Polar dent (unknown)	-
	Québec28 (Flint, Canada)	-

of the MITE class of mobile element [16]. This element was short (321 bp), had 15 bp imperfect inverted repeats and was flanked by a 5 bp (CTCAG) direct repeat generated upon insertion.

Amino acid changes

According to the amino acid change classifications of Li *et al* [19] and Grantham [20], of the five non-synonymous SNPs found in *ZmPox3*, two changes were conservative (Val/Leu and Phe/Leu) and three were moderately conservative (Ser/Thr, Ala/Ser and Ala/Thr). The N terminal signal peptide (residue 1 to 22), and the active site as well

as the heme-ligand site were not affected by any of these five amino acid replacements [7]. However, out of the 9 putative N-glycosylation sites identified [7] two were affected by one moderately conservative and one conservative amino acid changes.

Description of the *ZmPox3* MITE insertion

The MITE insertion in the second exon disrupts the gene 129 bp downstream of the 3' splice site of the first intron (Fig. 1A) by introducing a stop codon 75 bp after the beginning of the insertion (Fig. 1A). The conceptual translation of the mutant allele results in a truncated protein of




Figure 1
 Organisation of the *ZmPox3* gene. A) Schematic diagram of the *ZmPox3* gene showing the MITE insertion in the second exon (Arrows indicate positions of the primers used to amplify the gene and to genotype the MITE insertion. Boxes represent the exons). B) Translation products (Protein from the normal allele and truncated protein from the allele disrupted by the insertion of MITE element).

137 residues (Fig. 1B) the product of the translation of exon 1, the first third of exon 2 and the beginning of the MITE element. Since important functional sites (eg. heme-ligand signature and putative N-glycosylation sites) are absent in the deduced truncated protein, maize inbred lines carrying this mutant allele are very likely deficient in *ZmPox3* peroxidase activity.

Linkage disequilibrium and recombination

There were many occurrences of significant linkage disequilibrium (LD) among *ZmPox3* polymorphic sites (20.6% of the tested LD are significant at $P < 0.001$) (Fig. 2). After correction for multiple tests using the Bonferroni procedure, 11.7 % of LD remained significant ($P < 0.001$). LD concerned sites mainly located in the first intron. The

transposable element (S0465) was in strong LD with six polymorphic sites located in the first two exons and in intron 1 and 2 (Fig. 2). These polymorphisms in the vicinity of the insertion site may have been linked to the mutation and inherited as such. LD decayed with increasing physical distance according to Remington's model ([21], adjustment $F(1.350) = 457, P < 0.0001$) (Fig. 3). Three recombination events were detected in *ZmPox3* when assessed as the minimum number of recombination events using the algorithm proposed by Hudson and Kaplan [22].

Relationship between *ZmPox3* alleles and cell wall digestibility

ANOVA performed for a subset of 31 maize lines, for which digestibility values are known (Table 1), indicated that the insertion of the transposable element (MITE) in the second exon was significantly associated with cell wall digestibility ($P = 0.032$). This insertion characterized a group of five inbred lines related to F7 and having a high cell wall digestibility (F7, F7012, F324, F227 and F226). Based on the assumption that the lack of *ZmPox3* peroxidase activity might lead to reduced or modified lignification, comparison of cell wall digestibility was investigated in 25 lines related to F7 and did or did not carry the *zmPox3* mutant allele (Table 3). The cell wall digestibility was significantly higher in the 11 lines with the *zmPox3* mutant allele. Lines having the MITE insertion were also more related to F7 than the lines having the normal allele, but this difference was not significant and the MITE effect remained significant when the percentage of F7 genome was added as a co-variable (Table 3). Moreover, it appeared that two genetic backgrounds were favorable to cell wall digestibility in this set of early flint lines, independently of MITE insertion ; F7 (the Lacaune ecotype) and the germplasm from Argentina into which was bred F64 and is now found in lines F564, F286, F324, F7064 and F7065. Lines having a high cell wall digestibility (value ≥ 4), but not carrying the mutant allele, were all related to the Argentina germplasm (F286 and F564). In addition, line F324 carrying both the MITE insertion and Argentina germplasm, had a very high cell wall digestibility.

Table 2: Nucleotide polymorphism of the *ZmPox3* gene.

	Coding region			Non coding region		All silent sites	TOTAL
	Synonymous	Non synonymous	Introns	3'UTR	Total		
No of sites	272.12 *	788.88 *	238	135	373	645.12	1766
S SNP (singl)	8	5 (2)	17	1	18	26	31 (2)
S Indel (singl.)	0	1 **	14 (2)	3	17 (2)	17 (2)	18 (2)
π	0.008	0.0018	0.032	0.0036	0.02	0.015	0.0075
θ	0.007	0.0015	0.02	0.002	0.013	0.01	0.005




Figure 2

Linkage disequilibrium (LD) matrix among *ZmPox3* informative polymorphisms (SNPs and indels) for 37 maize sequences. The top part of graph indicates the position of each polymorphic site along the three genes sequenced regions (exon as black boxes). *P* values are the probabilities obtained using Bonferroni correction on Fisher's exact test. Polymorphic sites showing no LD are excluded from the matrix representation.

Expression profiles of the *ZmPox3* gene and its mutant allele

To determine whether the mutant allele was transcribed, the *ZmPox3* expression in F7012 (mutant *zmPox3* allele) and Lan496 (normal *ZmPox3* allele) were compared. Forward primer U19S1 located at the 5' end of the gene was combined either with primer U19R1 situated upstream or with primer U19MITEAS situated downstream of the MITE insertion in F7012 (Fig. 1A). The use of internal (18S) and external (pAW109) controls confirmed that both RNA quantity and RT-PCR were homogenous for

samples obtained from stems with surrounding leaf sheaths of the upper or basal part of the plant (Fig. 4A, lines 2 and 3). No difference in signal intensity was observed between F7012 and Lan496 for the upstream part of *ZmPox3* (Fig. 4A, line 4). On the contrary, the bands spanning the MITE insertion were barely visible in F7012 (Fig. 4A, line 5). This decrease for the mutant line could be explained by rapid RNA degradation of the non-translated part of the mutant allele, though we cannot exclude a decreased PCR efficiency due to possible secondary MITE structures. For experiments on leaf blades,




Figure 3
Decay of linkage disequilibrium with distance between nucleotide sites for *ZmPox3*. Curve shows non linear regression of r^2 on distance, by using model described in Remington *et al* [21]. The regression coefficient $b1$ is 0.00462

the signal was weaker in F7012 than in Lan496 for the upstream part of *ZmPox3*. This unexpected result was difficult to explain and may be due to the developmental differences of F7012 and Lan496 leaves. Nevertheless, a substantial difference between the bands reflecting the upstream and the MITE-spanning parts was also observed in F7012.

To obtain a more detailed expression profile of *ZmPox3*, its expression was studied using stem segments composed of nodes and the adjacent inter-nodes from F2 line at silking stage. *ZmPox3* expression was higher in lower-nodes and inter-nodes than in upper-nodes (Fig. 4B). Intense expression occurred in nodes close to the ear which developed on node 11. In the ear leaf, expression was very low 10 days after pollination, higher after 21 days and low again 30 days after pollination. *ZmPox3* peroxidase expression appeared important in lignifying tissue, but seemingly not in the earlier stage of lignification of adult plants. In comparison, O-methyl transferase (*AldOMT*) expression was stronger in older nodes and inter-nodes and extended much further to the younger nodes and inter-nodes at the top of the plant. In contrast, its expression was lower than that of *ZmPox3* in ear leaves (Fig. 4B). These data confirmed that the two genes are involved in different steps of the lignification process in maize.

Discussion

ZmPox3 nucleotide diversity was comparable to that previously reported for other nuclear genes in maize, and even slightly higher. For instance Tenaillon *et al* [23] and Rafalski [24] found one SNP approximately every 48 and 130 bp in the 3'UTR and coding regions, respectively. In *ZmPox3*, we counted on average one SNP every 57 bp.

Moreover, *ZmPox3* had a nucleotide diversity unequally distributed among site categories. There were 26 SNPs for all the silent sites and only five amino acids replacements for 357 residues (1.4%). Although the low frequency of non-synonymous polymorphisms suggested that at least some coding region of these genes are constrained, classical tests of neutrality showed no evidence of selection (data not shown). However, previous molecular diversity studies in maize have shown that departure from neutrality is rare [23]. A signature of positive selection could be revealed by an allele with unusually long-range LD [25]. In *ZmPox3* however, LD decayed very rapidly with increasing physical distance, within 200 bp on average. This result is consistent with recent studies showing that LD typically decays rapidly within individual maize loci [10,21,23].

A moderately conservative amino acid changes, located in the second exon, affected a putative N-glycosylation site. This SNP was characteristic of a cluster of four high digestibility lines (F564, EP1, Wis94-443 and Wis93-3520) and was associated with cell wall digestibility (ANOVA; $P = 0.03$). It would be interesting to determine whether this amino acid replacement affects enzymatic activity. For example, in rice, Larkin and Park [26] have reported that two SNPs which caused amino acid substitutions in the *waxy* gene affected the endosperm apparent amylose content. The MITE insertion was also shown to be associated with cell wall digestibility (ANOVA; $p = 0.032$). Furthermore, a survey of an additional set of 25 lines related to F7 revealed significant association between MITE and cell wall digestibility. This MITE element of 321 bp is 93% identical to the 316 bp insertion observed in the *B-M033* allele of the *b* gene, a regulatory gene involved in anthocyanin synthesis in maize [27]. It is also very similar to a transposon in intron 4 (92% identity) of the maize β -D-glucosidase gene (accession U60560), and to a 316 bp insertion in the 5'UTR region of the maize *waxy* gene (89% identity) [28]. It has been suggested that MITE insertions in non-coding regions such as the 5'UTR or 3'UTR of transcribed maize genes, provide regulatory sequences involved in transcription initiation or mRNA stability and are thus important tools of evolution [16]. Thornsberry *et al* [13] found significant association between flowering time traits and the 485 bp insertion in the 5' non coding region of the *Dwarf8* gene caused by a putative MITE element. To our knowledge, the insertion of a MITE element in the second exon of *ZmPox3* has never been previously reported in the coding region of maize genes. This insertion very likely resulted in transcription of a truncated and non-functional *ZmPox3*. Thereby, the high digestibility of numerous maize lines carrying the *zmPox3* mutant allele could be considered as the result of a deficiency in the activity of this peroxidase probably involved in the lignin biosynthesis.




Figure 4
 Expression profile of *ZmPox3*. Total RNA was isolated from the indicated tissues and amplified by RT-PCR. Amplification products were visualized by agarose gel electrophoresis. RNA transcribed from plasmid pAW109 was used as an external loading control. The *ZmPox3* forward primer (U19S1) was combined with a reverse primer upstream (U19R1) or downstream (U19MITEAS) of the MITE insertion. PCR cycles used to evaluate *ZmPox3* varied between panel A and panel B (indicated by numbers to the right of the gel pictures). A) *ZmPox3* and *AldOMT* expression in lines F7012 (mutant allele) or Lan496 (normal allele). The stem with the leaf sheaths of 50 day old plants was divided into a basal part (4 inter-nodes) and an upper part (ending at the base of the tassel). In addition a 30 cm long leaf blade was harvested. B) *ZmPox3* and *AldOMT* expression in different parts of line F2 (normal allele). At flowering the stem with the leaf sheaths was cut at the base of every node yielding units of node (N) and adjacent inter-node (i) with numbering starting at the base of the plant. In addition the ear leaves were harvested at defined stages expressed in days after pollination (DAP).

As reported by de Obeso *et al* [7], *ZmPox3* mRNA was detected at low levels in lignifying tissues of young maize roots. Our RT-PCR experiments highlighted that the expression of the *ZmPox3* gene was weaker in the first steps of stem lignification and in younger parts of the stem, than in older parts of the stem. These results, the well-known role of laccases and peroxidases at the final steps of lignin biosynthesis [7] and the identification of a *ZmPox3* ortholog in the *Zinnia elegans* model system of lignifying tissues (S. Guillaumie, pers. com.) reinforce the possibility of an involvement of this peroxidase in maize monolignol polymerisation. The expression profile was clearly distinct from that of *AldOMT* indicating that lignification involves precise spatial and temporal regulation

of the numerous genes involved in the different steps of this complex process.

Conclusions

Analysis of allelic diversity in relation with cell wall digestibility values likely validated *ZmPox3* as a plausible candidate gene for silage maize digestibility improvement. This peroxidase could have a direct effect on plant cell wall digestibility, even more so given that this peroxidase is colocalized with a cell wall digestibility and lignification QTL. The comparison between *ZmPox3* alleles that differ mostly by the presence or absence of the MITE element offers an opportunity to evaluate the impact of this kind of mutation event, as it was extensively done for

Table 3: Comparisons of cell wall digestibility in 25 maize lines related to F7 with and without the MITE insertion in *ZmPox3*.

Lines	Pedigree	% F7	<i>ZmPox3</i> MITE	Cell wall digestibility
F7	Lacaune ecotype	100	present	4
F192	F7 × F2	50	present	3
F226	Du101 × F7	50	present	3.5
F227	F7 × F120	50	present	3
F283	F226 × F227	50	present	3.5
F324	(F282 × F283) × F286	62.5	present	5
F7012	F226 × F227	50	present	4.5
F7032	Early flint synthetic L	25	present	3.5
F7064	F7012 × (BSSS × F286)	43.7	present	4.5
F7065	F7012 × (BSSS × F286)	43.7	present	5
LGFS	Early flint synthetic P	12.5	present	3.5
CPI718	Private line	25	absent	3
CPI622	Private line	25	absent	3
F131	F7 × CH10	50	absent	2.5
F286	F7 × F564	75	absent	4.5
F268	(F215 × F192) × (F2 × F160)	12.5	absent	1.5
F564	F7 × F64 (F64 = Argentina flint)	50	absent	4
F7023	Early flint synthetic T	35	absent	1.5
LGI1	Private line	25	absent	3
LGI2	Private line	25	absent	2
LGI9	Private line	12.5	absent	1.5
LGD3	Private line	20	absent	2.5
R2n02	Private line	50	absent	3.5
R2n12	Private line	25	absent	2.5
R2n13	Private line	25	absent	3
Mean 11 lines with MITE present		48.9		3.9
Mean 14 lines with MITE absent		32.5		2.7
F value		4.3 ns		12.4 **
F value with % F7 as covariable		-		6.6 **

brown-midrib alleles [29]. Thus, either the targeted introduction of the mutant allele in different genetic backgrounds or the down-regulation in transgenic lines could confirm the usefulness of *ZmPox3* MITE disruption in breeding maize for higher digestibility. Moreover, because it seems to be a rare event, the search for the *ZmPox3* MITE insertion in genetic resources could also help in determining the phylogeny of maize lines and introductions to Europe.

Methods

Plant material

Thirty-seven inbred lines were chosen in order to display a large variation in maize cell wall digestibility and to represent a diversified sample of genotypes used both in European breeding programs and representative of commonly used US germplasm (Table 1). Maize plants were grown in vermiculite supplemented with a nutrient solution under a 16/8 h light/dark regime for 10 days. Because an abnormal *zmPox3* gene was first found in F7012, and then in different parents of F7012 which were all related

to the INRA flint inbred F7, the existence of this mutant allele was then investigated in 15 other private and public lines related to F7 (Table 3). RT-PCR investigations of *ZmPox3* expression were done i) on successive nodes and inter-nodes n° 5 to 14 in plants of line F2 harvested at silking date, ii) on ear leaves of line F2 harvested 10, 21 and 40 days after silking, under field conditions, and iii) in basal and upper plant parts (stem and sheaths), and in the largest expanded leaves of lines Lan496 and F7012 harvested 50 days after sowing in a greenhouse.

DNA extraction, primer design and PCR amplification

Genomic DNA was isolated from young maize leaves using the plant DNAEASY miniprep kit (Qiagen).

A pair of primers was designed based on published cDNA sequence (accession number AJ401276): U19S1 (forward; 5'-GACGAAGCGGCACTGCTTGCCTTACCA-3') and U19AS1 (reverse; 5'-TGCCACAGTAACAAGCGAGCTTACCAAGA-3'), respectively complementary to positions 1-29 and to 1198-1170 of the maize *ZmPox3* cDNA sequence.

Table 4: Pairs of primers used to obtain the *ZmPox3* gene and to genotype the MITE insertion.

	U19S1 U19ASI	U19MITES U19MITEAS	U19MITES U19ASI	U19S1 U19MITEAS
normal allele	1.4 kb	none	none	05 kb
mutant allele	1.7 kb	0.3 kb	1.2 kb	0.8 kb
heterozygote	1.4 kb 1.7 kb	0.3 kb	1.2 kb	0.5 kb 0.8 kb

Amplified fragments were about 1.4 to 1.7 kb long, and encompassed the coding region and 3'UTR (Table 4 and Fig. 1A).

A pair of primers flanking the MITE insertion was designed (Fig. 1A) as U19MITES (forward; 5'-GGCACTGGAGGCTCAGGGTGTGTT-3') and U19MITEAS (reverse; 3'-AGGAGACAACGCCGGGCAC-5'). If a maize line had the mutant allele, a 0.3 kb fragment corresponding to the MITE element was amplified. On plants having the normal allele, no amplification occurred (Table 4 and Fig. 1A). Combinations between pairs of primers U19S1 / U19ASI and U19MITES / U19MITEAS allowed us to distinguish between homozygote and heterozygote genotypes (Table 4). The procedure to detect the *zmPox3* mutant allele has been also described in detail in the French patent FR0302954.

PCR amplification reactions were performed in 50 µl containing 100 mM tris-HCl, 500 mM KCl, 15 mM MgCl₂ and 0.01% gelatin, 200 µM of each dNTP, 0.2 µM of 5' oligo, 0.2 µM of 3' oligo and 2.5 units of REDTaq DNA Genomic Polymerase (Sigma). 100 ng of genomic DNA was used as template. To amplify the *ZmPox3* gene, the program was 5 min at 95°C followed by 30 cycles of 30 s at 95°C, 30 s at 60°C, and 1.5 min at 72°C, followed by 5 min at 72°C. To genotype the MITE insertion in *ZmPox3*, the PCR amplification protocol was slightly modified and consisted of 5 min at 95°C followed by 25 cycles of 30 s at 95°C, 30 s at 65°C, and 30 s at 72°C, followed by 5 min at 72°C.

DNA sequencing

Sequencing was performed for each PCR fragments in both directions by Isoprism (France) and MWG-Biotech (Germany). Ecotype Québec28 was heterozygous at *ZmPox3* locus and the two alleles have been sequenced. The sequences containing singletons were checked by re-amplifying genomic DNA and partially re-sequencing the appropriate alleles. Sequences were aligned using CLUSTALW [30]. Sequence data from this article have been deposited with the EMBL/GenBank Data Libraries

under accessions n° AY500781-AY500812; AY508159-AY508163 and AY508516.

RT-PCR analysis

For isolation of total RNA, tissues were placed together with a stainless steel bead in 2 ml Eppendorf tubes and frozen in liquid nitrogen. Using pre-cooled holders the tissues were ground to powder in a Mixer Mill MM300 (Qiagen) by shaking two times 30 sec. The ground, frozen tissues were vortexed with 1 ml TRIzol® Reagent (Invitrogen) at room temperature until melted. The aqueous phase resulting from a 10 min centrifugation at 18000 g and 4°C was re-extracted with 200 µl chloroform at room temperature. The RNA was precipitated with 500 µl isopropanol for 10 min at room temperature. The RNA pellet obtained by 10 min centrifugation at 18000 g and 4°C was washed with 1 ml 70% ethanol, dried and resuspended in 30 µl RNase free water. After treatment with RNase free DNase and inactivation of the DNase according to the instructions of the supplier (AMBION) the RNA was quantified in a spectrophotometer at 260 nm. Approximately 5 µg of total RNA were reverse transcribed using random hexamers (Amersham) and reverse transcriptase without RNaseH activity (Fermentas). The 20 µl reverse transcription reaction also contained 2.5 × 10⁵ copies of GeneAmplicon pAW109 RNA (Applied Biosystems). The obtained cDNA was diluted 50 times in water and 5 µl used for amplification by PCR in a volume of 20 µl.

Part of *ZmPox3* was amplified using primer U19S1 in combination with either primer U19R1 (5'-CGTCAGGTTGCCTACCGTGTGTCGATCAGCAC-3') situated 84 bp upstream of the MITE insertion or primer U19MITEAS located downstream of the insertion. The constitutively expressed 18S rRNA gene (primers 5' CCATCCCTCCGTAGTTAGCTTCT 3' and 5' CCTGTCCGCCAAGGCTATATAC 3') was used as an internal control of RNA quantity and GeneAmplicon pAW 109 RNA (primers 5' CATGTCAAATTTCACTGCTTCATC 3' and 5' TGACCACCCAGCCATCCTT 3') as positive control of the RT-PCR efficiency. In order to get semi-quantitative results, the

number of cycles of the PCR reactions was adjusted for each gene to obtain barely visible bands in agarose gels. Aliquots of the PCR reactions were loaded on agarose gels and stained with ethidium bromide.

Evaluation of maize silage digestibility

In vitro cell wall digestibility of lines (*per se* values) was investigated through different multi-year and multi-local experiments managed by the different partners. Values of cell wall digestibility were estimated as described in Rousset *et al* [8] through the DINAGZ criterion (*in vitro* digestibility of the "non starch (ST), non soluble carbohydrates (SC) and non crude protein (CP) part") based on the enzymatic solubility of the whole plant [31]. Data were gathered in order to obtain an index score of cell wall digestibility for each line from 1 (40 %) to 5 (60 %), according to the DINAGZ trait, whose heritability is most often close to 0.75 [1]. Lines F271 and F4, possessed the lowest and highest cell wall digestibility in this set of maize lines, respectively

Data analysis

DNA sequences were analyzed using DnaSP [32]. Levels of nucleotide diversity were estimated as mean pairwise differences (π ; [17]) and number of segregating sites (θ ; [18]). The minimum number of intragenic recombination events was estimated using the four-gamete test [22]. Insertions and deletions (indels) were excluded from the estimates. The significance of pairwise linkage disequilibrium (LD) among polymorphic sites (SNPs and indels) was tested using Fisher's exact test excluding non-informative sites (singletons), and corrected for multiple analyses using the Bonferroni procedure [33]. The decay of LD with physical distance along genes was evaluated by non-linear regression (PROC NLIN in SAS software, SAS institute 1999) following Remington's model [21] that considers potential low mutation rate and adjustment for sample size.

Each informative polymorphic site (SNPs and indels) was examined independently for an association with variation in cell wall digestibility using an ANOVA. The set of 31 lines for which digestibility values are known was used (Table 1).

A single factor variance analysis (MITE presence or absence) was used to compare the average digestibility value of lines, either having or not having the *zmPox3* mutant allele, in the set of 25 lines related to F7 (Table 3). MITE effect was similarly investigated after adding the percentage of F7 genome in each line (expected value according to the pedigree) as a co-variable.

List of abbreviations

bp – base pairs, Indel – insertion/deletion, kb – kilo bases, MITE – miniature inverted-repeat transposable element, PCR – polymerase chain reaction, QTL – quantitative trait loci, SNP – single nucleotide polymorphism, UTR – untranslated region.

Authors' contributions

CGC carried out most of the experimental studies and data analysis and participated in the discussion and preparation of the manuscript. CBT participated in the DNA sequencing and data analysis. DM carried out linkage disequilibrium measurement and participated in the discussion. PR carried out RT-PCR experiments. JR carried out the *ZmPox3* cloning. AM carried the *ZmPox3* mapping and participated in the design. JPM participated in the design and the coordination of the study. YB contributed to the design and coordination of the study and participated in the discussion and preparation of the manuscript.

Acknowledgements

This work was supported by grants from the Genoplante maize program, coordinated by Alain Charcosset (INRA Le Moulon) and Alain Murigneux (Biogemma, Les Cézeaux). We thank Sabine Guillaumie and Marie Fourmann for comments on the manuscript. We thank Jeremy l'Homedet, Richard Printemps and Stéphanie Barraud for skilful technical assistance. We also thank Pascal Condamine and Agnes Massonneau for their contribution to the RT-PCR analysis. Limagrain Genetics, Pau-Euralis and RAGT Semences are thanked for providing seeds of lines related to F7 and information on their cell wall digestibility. We are very grateful to Pierre-Philippe Claude for encouragement and help during manuscript preparation.

References

1. Barrière Y, Guillet C, Gofner D, Pichon M: **Genetic Variation and Breeding Strategies for Improved Cell Wall Digestibility in Annual Forage Crops. A Review.** *Anim Res* 2003, **52**:193-228.
2. Boerjan W, Ralph J, Baucher M: **Lignin biosynthesis.** *Annu Rev Plant Biol* 2003, **54**:519-546.
3. Boudet AM: **Towards an understanding of the supramolecular organization of the lignified wall.** *The plant cell wall* Edited by: J Rose. Blackwell publishing; 2003:155-182.
4. Blee KA, Choi JW, O'Connell AP, Schuch W, Lewis NG, Bolwell GP: **A lignin-specific peroxidase in tobacco whose antisense suppression leads to vascular tissue modification.** *Phytochemistry* 2003, **64**:163-176.
5. Li Y, Kajita S, Kawai S, Katayama Y, Morohoshi N: **Down-regulation of an anionic peroxidase in transgenic aspen and its effect on lignin characteristics.** *J Plant Res* 2003, **116**:175-182.
6. **The Maize Genetics and Genomics Database** [<http://www.maizegdb.org>]
7. de Obeso M, Caparros-Ruiz D, Vignols F, Puigdomenech P, Rigau J: **Characterisation of maize peroxidases having differential patterns of mRNA accumulation in relation to lignifying tissues.** *Gene* 2003, **309**:23-33.
8. Rousset V, Gibelin C, Fontaine AS, Barrière Y: **Genetic Analysis in Recombinant Inbred Lines of Early Dent Forage. II- QTL Mapping for Cell Wall Constituents and Cell Wall Digestibility from *per se* Value and Top Cross Experiments.** *Maydica* 2002, **47**:9-20.
9. Méchin V, Argillier O, Hébert Y, Guingo E, Moreau L, Charcosset A, Barrière Y: **Genetic Analysis and QTL Mapping of Cell Wall Digestibility and Lignification in Silage Maize.** *Crop Sci* 2001, **41**:690-697.

10. Clark RM, Linton E, Messing J, Doebley JF: **Pattern of diversity in the genomic region near the maize domestication gene *tb1*.** *Proc Natl Acad Sci USA* 2004, **101**:700-707.
11. Ching A, Caldwell KS, Jung M, Dolan M, Smith OS, Tingey S, Morgante M, Rafalski AJ: **SNP frequency, haplotype structure and linkage disequilibrium in elite maize inbred lines.** *BMC Genet* 2002, **3**:19.
12. Buckler ES 4th, Thornsberry JM: **Plant molecular diversity and applications to genomics.** *Curr Opin Plant Biol* 2002, **5**:107-111.
13. Thornsberry JM, Goodman MM, Doebley J, Kresovich S, Nielsen D, Buckler ES 4th: **Dwarf8 polymorphisms associate with variation in flowering time.** *Nat Genet* 2001, **28**:286-289.
14. Palaisa KA, Morgante M, Williams M, Rafalski A: **Contrasting effects of selection on sequence diversity and linkage disequilibrium at two phytoene synthase loci.** *Plant Cell* 2003, **15**:1795-1806.
15. Poke FS, Vaillancourt RE, C. Elliot R., Reid BR: **Sequence variation in two lignin biosynthesis genes, cinnamoyl CoA reductase (CCR) and cinnamyl alcohol dehydrogenase (CAD2).** *Mol Breed* 2003, **12**:107-118.
16. Wessler SR, Bureau TE, White SE: **LTR-retrotransposons and MITEs: important players in the evolution of plant genomes.** *Curr Opin Genet Dev* 1995, **5**:814-821.
17. Tajima F: **Evolutionary relationship of DNA sequences in finite populations.** *Genetics* 1983, **105**:437-460.
18. Watterson GA: **On the number of segregating sites in genetic models without recombination.** *Theor Popul Biol* 1975, **7**:256-276.
19. Li WH, Wu CI, Luo CC: **A new method for estimating synonymous and nonsynonymous rates of nucleotide substitution considering the relative likelihood of nucleotide and codon changes.** *Mol Biol Evol* 1985, **2**:150-174.
20. Grantham R: **Amino acid difference formula to help explain protein evolution.** *Science* 1974, **185**:862-864.
21. Remington DL, Thornsberry JM, Matsuoka Y, Wilson LM, Whitt SR, Doebley J, Kresovich S, Goodman MM, Buckler ES 4th: **Structure of linkage disequilibrium and phenotypic associations in the maize genome.** *Proc Natl Acad Sci U S A* 2001, **98**:11479-11484.
22. Hudson RR, Kaplan NL: **Statistical properties of the number of recombination events in the history of a sample of DNA sequences.** *Genetics* 1985, **111**:147-164.
23. Tenailon MI, Sawkins MC, Long AD, Gaut RL, Doebley JF, Gaut BS: **Patterns of DNA sequence polymorphism along chromosome 1 of maize (*Zea mays ssp. mays* L.).** *Proc Natl Acad Sci U S A* 2001, **98**:9161-9166.
24. Rafalski A: **Applications of single nucleotide polymorphisms in crop genetics.** *Curr Opin Plant Biol* 2002, **5**:94-100.
25. Rafalski A, Morgante M: **Corn and humans: recombination and linkage disequilibrium in two genomes of similar size.** *Trends Genet* 2004, **20**:103-111.
26. Larkin PD, Park WD: **Association of waxy gene single nucleotide polymorphism with starch characteristic in rice (*Oryza sativa* L.).** *Mol Breed* 2003, **12**:335-339.
27. Selinger DA, Chandler VL: **Major recent and independent changes in levels and patterns of expression have occurred at the *b* gene, a regulatory locus in maize.** *Proc Natl Acad Sci USA* 1999, **96**:15007-15012.
28. Spell ML, Baran G, Wessler SR: **An RFLP adjacent to the maize waxy gene has the structure of a transposable element.** *Mol Gen Genet* 1988, **211**:364-366.
29. Marita JM, Vermerris W, Ralph J, Hatfield RD: **Variations in the cell wall composition of maize brown midrib mutants.** *J Agric Food Chem* 2003, **51**:1313-1321.
30. Thompson JD, Higgins DG, Gibson TJ: **CLUSTAL W: improving the sensitivity of progressive multiple sequence alignment through sequence weighting, positions-specific gap penalties and weight matrix choice.** *Nucleic Acids Res* 1994, **22**:4673-4680.
31. Aufrère J, Michalet-Doreau B: **In vivo digestibility and prediction of digestibility of some by-products.** *ECC Seminar Belgique*; 1983.
32. Rozas J, Sanchez-DelBarrio JC, Messeguer X, Rozas R: **DnaSP, DNA polymorphism analyses by the coalescent and other methods.** *Bioinformatics* 2003, **19**:2496-2497.
33. Sokal RR, Rohlf FJ: **Biometry.** Edited by: Ed W H Freeman. San Francisco; 1981.

Publish with **BioMed Central** and every scientist can read your work free of charge

"BioMed Central will be the most significant development for disseminating the results of biomedical research in our lifetime."

Sir Paul Nurse, Cancer Research UK

Your research papers will be:

- available free of charge to the entire biomedical community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours — you keep the copyright

Submit your manuscript here:
http://www.biomedcentral.com/info/publishing_adv.asp

