



HAL
open science

Genotype-level variation in lifetime breeding success, litter size and survival of sheep in scrapie-affected flocks

Margo E. Chase-Topping, Loeske E.B. Kruuk, Daniel Lajous, Suzanne Touzeau, Louise Matthews, Geoff Simm, James D. Foster, Rachel Rupp, Francis Eychenne, Nora Hunter, et al.

► To cite this version:

Margo E. Chase-Topping, Loeske E.B. Kruuk, Daniel Lajous, Suzanne Touzeau, Louise Matthews, et al.. Genotype-level variation in lifetime breeding success, litter size and survival of sheep in scrapie-affected flocks. *Journal of General Virology*, 2005, 86 (4), pp.1229-1238. 10.1099/vir.0.80277-0 . hal-02683319

HAL Id: hal-02683319

<https://hal.inrae.fr/hal-02683319>

Submitted on 23 Jan 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

1 **Genotype-level variation in lifetime breeding success, litter size**
2 **and survival of sheep in scrapie affected flocks**

3
4 Margo E. Chase-Topping¹, Loeske E.B. Kruuk^{1,2}, Daniel Lajous³, Suzanne
5 Touzeau⁴, Louise Matthews¹, Geoff Simm⁵, James D. Foster⁶, Rachel Rupp³,
6 Francis Eychenne³, Nora Hunter⁶, Jean-Michel Elsen³, and Mark E.J. Woolhouse¹

7
8 ¹ Centre for Tropical Veterinary Medicine, University of Edinburgh, Easter Bush,
9 Roslin, Midlothian EH25 9RG, UK

10 ² Institute of Cell, Animal and Population Biology, University of Edinburgh,
11 Edinburgh, EH9 3JT

12 ³ Institut National de la Recherche Agronomique, Station d'Amélioration
13 génétique des Animaux, BP 27, 31326 Castanet-Tolosan Cedex, France

14 ⁴ Institut National de la Recherche Agronomique, Unité de Mathématiques et
15 Informatique appliquées, 78352 Jouy-en-Josas Cedex, France

16 ⁵ Sustainable Livestock Systems Group, Scottish Agricultural College, Bush
17 Estate, Penicuik, Midlothian EH26 0PH, UK

18 ⁶ Institute for Animal Health Neuropathogenesis Unit, Ogston Building, West
19 Mains Road, Edinburgh EH9 3JF, UK

20

21

22

23

1 Corresponding Author:

2

3 Dr. Margo E. Chase-Topping

4 Centre for Tropical Veterinary Medicine (CTVM)

5 Royal (Dick) School of Veterinary Studies

6 The University of Edinburgh

7 Easter Bush, ROSLIN, Midlothian

8 Scotland, EH25 9RG, UK

9 phone / fax : +44 (0) 131 650 7263/445 5099

10 Email: margo.chase@ed.ac.uk

11

12

13 Running head: Genotype variation in scrapie affected flocks

14

15

16 **Total number of words in body of text: 4826**

17 **Total number of tables and figures: 8**

18 **Total number of words in summary: 149**

1 **SUMMARY**

2

3 Five different sheep flocks with natural outbreaks of scrapie were examined to
4 determine associations between individual performance (lifetime breeding success
5 (LBS), litter size and survival) and scrapie infection or PrP genotype. Despite
6 different breed composition and forces of infection, consistent patterns were
7 found among the flocks. Regardless of the flock, scrapie infected sheep produced
8 on average 34% fewer offspring than non-scrapie infected sheep. The effect of
9 scrapie on LBS appears to be a function of lifespan as opposed to fecundity.
10 Analysis of litter size revealed no overall or genotype differences among the 5
11 sheep flocks. Survival, however, depends on the individual's scrapie status
12 (infected or not) and its PrP genotype. Susceptible genotypes appear to perform
13 less well in LBS and life-expectancy even if they are never affected with clinical
14 scrapie. One possible explanation for these results is the effect of pre-clinical
15 scrapie. Additional evidence supporting this hypothesis is discussed.

16

1 **INTRODUCTION**

2

3 Scrapie is a transmissible spongiform encephalopathy (TSE), a category of fatal
4 and incurable diseases that includes bovine spongiform encephalopathy (BSE),
5 chronic wasting disease (CWD), transmissible mink encephalopathy (TME),
6 feline spongiform encephalopathy (FSE), Kuru and variant Creutzfeldt-Jakob
7 disease (vCJD). Scrapie has been reported world-wide and affects many sheep
8 producing regions (Dawson *et al.* 1998). It has been present in the sheep
9 population of Britain since the mid-18th century (Parry, 1983; Stamp, 1962) and
10 remains widespread throughout the country.

11

12 Despite recent detailed studies of scrapie outbreaks within individual sheep flocks
13 (Elsen *et al.*, 1999; Hunter *et al.*, 1996; Hunter *et al.* 1997) and comparative
14 epidemiological analysis on multiple sheep flocks (Redman *et al.* 2002), key
15 determinants of epidemiological and transmission dynamics of sheep scrapie are
16 still poorly understood. In recent years considerable progress has been made in
17 establishing the genetics of susceptibility of scrapie (Dawson *et al.* 1998; Hunter
18 *et al.* 1997). It is known that resistance or susceptibility is largely under genetic
19 control (Hunter, 1997), however, the effects of PrP genotype on scrapie
20 susceptibility can vary between flocks and breeds of sheep (Dawson *et al.* 1998)
21 and can also depend on scrapie isolates (Goldmann *et al.* 1994). To date, there
22 have been few detailed within-flock studies of the effects of variation in PrP
23 genotype at the individual level during natural scrapie outbreaks. Many studies

1 have been performed to determine the genetic status and variability of PrP
2 genotype of sheep breeds in different countries (e.g. Germany: Drogemuller *et al.*
3 2001; Italy: Vaccari *et al.* 2001 and Spain: Acin *et al.* 2004), and a few studies
4 have examined the PrP genotype profile of individual flocks (e.g. Baylis *et al.*
5 2000). Previous research that has examined genotype-level associations within
6 flocks have generally focused on the relationship with scrapie infection, including
7 incubation time (Goldmann *et al.*, 1991, 1994) and age of onset of scrapie (Baylis
8 *et al.* 2002; Bossers *et al.* 1996; Clouscard *et al.* 1995; Elsen *et al.* 1999). Despite
9 the extensive amount of research that exists on scrapie infection no study has
10 attempted to quantify the effect of scrapie on significant performance parameters
11 such as lifetime breeding success, litter size or survival. A few analyses have
12 examined PrP genotype-level associations with performance parameters (e.g.
13 Brandsma *et al.* 2004: litter size and 135 days weight; Barillet *et al.* 2002: dairy
14 production traits, Prokopova *et al.* 2002: lean growth rate; and de Vries *et al.*
15 2004: muscle mass, liveweight gain, wool quality and fat depth). Overall these
16 studies have found no significant association between PrP genotype and the trait
17 examined, although some association between the resistant ARR and depth of
18 muscle mass was found in German black-headed mutton sheep (de Vries *et al.*
19 2004). However, these studies examined the traits in the absence of scrapie
20 infection, with a view to determining the effect of breeding for resistance, rather
21 than the population dynamic and population genetic implications of a natural
22 scrapie outbreak within a flock.

23

1 As with all TSEs, scrapie has a long incubation period between infection and
2 onset of typical clinical signs. Although there is no explicit evidence to date for
3 effects of pre-clinical scrapie, it has been identified as a possible cause for
4 unexplained mortality within flocks (McLean *et al.* 1999). Furthermore, the focus
5 of research on outbreaks of scrapie in sheep flocks has been on scrapie cases, and
6 no study has considered individuals that did not develop clinical signs. Genotype-
7 related differences in the performance of sheep manifesting no signs of scrapie
8 may indicate the presence of pre-clinical scrapie within the flock. Identification
9 and quantification of this phenomenon may result in changes in the incidence of
10 scrapie deaths and the overall impact of scrapie as a disease within sheep.

11

12 In this paper, we focus on differences in individual performance associated with
13 scrapie infection or PrP genotype in five different sheep flocks with natural
14 outbreaks of scrapie. An outbreak of scrapie should exert substantial selection
15 pressures against those PrP alleles associated with susceptibility. We illustrate the
16 force of this selection by quantifying the effect of scrapie on individual fitness,
17 assessed through estimates of individual lifetime breeding success (LBS).

18 Differences in LBS due to scrapie are expected within each flock. Such
19 differences may be the result of differential longevity and/or differential
20 fecundity. We examine each component across five different sheep flocks.

21 Measures of individual lifetime breeding success, litter size and survival are used
22 to quantify: (1) the impact of scrapie; and (2) differences between PrP genotypes
23 in scrapie and non-scrapie infected sheep.

1

2 **METHODS**

3

4 **Study Flocks**

5

6 Data were generated from five outbreaks of natural scrapie (Table 1). Three of the
7 outbreaks were in flocks maintained by the Institute for Animal Health
8 Neuropathogenesis Unit (NPU), one in a flock maintained by the Scottish
9 Agricultural College (SAC) and one in a flock maintained by the Institut National
10 de la Recherche Agronomique (INRA) (Table 1). All flocks were maintained for
11 research purposes. The origins and histories of the flocks are described in greater
12 detail elsewhere (Elsen *et al.* 1999; Hunter *et al.* 1996, 1997; Redman *et al.* 2002).

13

14 **Field Data**

15

16 The following data are available for almost all individual sheep in each flock: date
17 of birth; pedigrees; date of death or removal from flock; cause of death or reason
18 for removal. Scrapie was suspected based on clinical signs, including loss of
19 condition and rubbing. Suspect scrapie cases were confirmed by histopathological
20 detection of vacuolation of brain tissue. Only confirmed cases of scrapie were
21 used in the analysis.

22

23 For three of the outbreaks, the SAC Suffolk, the NPU Cheviot II and the INRA
24 Romanov; there was some information on PrP genotypes, established by

1 sequencing PCR products or using oligonucleotide probes, as previously
2 described (Elsen *et al.* 1999; Hunter *et al.* 1996, 1997). Data from these three
3 flocks were used to examine genotype variation in lifetime breeding success, litter
4 size and survival. For the INRA Romanov flock, genotype data were available for
5 all animals in the flock since the onset of scrapie in 1993 whereas the genotyped
6 individuals in the SAC Suffolk and the NPU Cheviot II consisted of most scrapie
7 cases and approximately 50% of the non-scrapie infected sheep in each flock. As
8 such, the focus of the genotype variation analysis was on the INRA Romanov
9 flock. However, where possible, corresponding data were presented for the NPU
10 Cheviot II and SAC Suffolk flocks.

11

12 **Statistical Analysis**

13

14 Data within each database was standardised to suit the analysis that was to be
15 performed. For all flocks experimental, non-breeding animals and all males were
16 excluded from the analysis. In the INRA Romanov flock the breeding practices
17 with males were different: replacement sires were not used for long and
18 experimental animals were mostly males and culled according to protocol. Males
19 were therefore removed from the other flocks to standardise the data. Statistical
20 analysis was first performed on each flock to determine differences associated
21 with scrapie status (scrapie infected vs non-scrapie infected). For flocks with
22 genotype information (INRA Romanov, NPU Cheviot II and SAC Suffolk)
23 individuals were categorised as either susceptible (genotypes that were affected by
24 scrapie) or non-affected (genotypes that were not affected by scrapie or scrapie

1 infection was low or suspect). Susceptible and non-affected genotypes within each
2 flock are listed in Table 2.

3
4 **Variation in lifetime breeding success.** Lifetime breeding success (LBS) was
5 calculated as the total number of live offspring produced by each breeding female,
6 with and without scrapie, in each flock. Data analysis included all cohorts
7 involved in the outbreak (Table 1) with the exception of the INRA Romanov
8 flock. Data collection in the INRA Romanov flock is ongoing therefore there are
9 living females that have yet to produce all their offspring. As such, lifetime
10 breeding data is not available for these animals. Therefore, the analysis in the
11 INRA Romanov flock was restricted to cohorts born between 1986 (first cohort
12 involved in the outbreak) and 1993, excluding those which died prior to the 1993-
13 1999 outbreak.

14
15 Mean (\pm SE) lifetime breeding success was estimated for each flock. Differences
16 in the LBS of scrapie infected and non-scrapie infected sheep within each flock
17 were analysed using a Student t-test. To determine if there were any differences in
18 the effect of scrapie on LBS across the five flocks a comparison was performed
19 using a Generalised Linear Model with negative binomial errors (S-Plus Version
20 6.0) and the significance of the flock*status interaction was assessed from the
21 change in deviance on dropping that term from the model, distributed as $\chi^2_{(4)}$. For
22 flocks with genotype information, two analyses were performed to examine
23 differences in LBS; the first examined differences between scrapie infected and
24 non-scrapie infected individuals within and across susceptible genotypes and the

1 second examined the LBS of non-scrapie infected individuals, looking for
2 differences between susceptible and non-affected genotypes. Both analyses were
3 performed using a General Linear Model (GLM) (SAS Version 8.2).
4
5 **Variation in litter size** . The size of all litters produced throughout the scrapie
6 outbreak was calculated for scrapie and non-scrapie dams in each flock. Data
7 from all flocks were standardised to include all litters born within the years of the
8 scrapie outbreak (Table 1).
9
10 Differences between scrapie and non-scrapie individuals and between PrP
11 genotypes in the number of live lambs per litter (“litter size”) produced by dams at
12 each breeding event were tested. Linear mixed effects models with dam identity
13 fitted as a random effect were used to account for the repeated measures made on
14 individual sheep over multiple breeding attempts. PrP genotype or scrapie status
15 was used as a fixed effect. Models were fitted with Poisson errors using the
16 procedure glmmPQL (S-Plus Version 6). For all flocks, we initially tested for
17 effects of breeding year (as a multilevel factor) and dam age (as a quadratic
18 function). These variables, if significant, ($P < 0.05$) were retained in the models, as
19 follows: NPU Cheviot II, dam age; SAC Suffolks, breeding year; INRA
20 Romanovs, NPU Cheviot I and NPU Suffolks, dam age and breeding year.
21 Analyses of associations between litter size and PrP genotype were restricted to
22 the INRA Romanov and SAC Suffolk flocks due to insufficient genotype data in
23 the other flocks.
24

1 **Variation in survival** . Survival analyses were performed on the female
2 population considering the age at removal from flock as the survival
3 measurement. Removal includes animals that died naturally as well as those
4 culled for non-experimental reasons. Data analysed included only cohorts which
5 were exposed to scrapie (Table 1). All survival analyses were performed using
6 Proc lifetest and Proc Phreg (SAS Version 8.2). Median life expectancies (\pm 95%
7 confidence intervals) were calculated using survival data censored for sheep
8 culled at less than 1 year of age and those still alive. Data were stratified by
9 genotype (VRQ/VRQ, ARQ/VRQ, ARQ/ARQ, Non-affected) and scrapie status
10 (scrapie infected, non-scrapie infected). The following null hypotheses were
11 tested in Proc Lifetest: (1) there are no differences in the overall mean life
12 expectancy of scrapie infected versus non-scrapie infected individuals within each
13 of the 5 flocks; and (2) there are no differences in the mean life expectancy of
14 non-scrapie infected individuals among the susceptible and non-affected
15 genotypes in the NPU Cheviot II, SAC Suffolk and INRA Romanov flocks.
16 Differences between survivorship curves were tested using Kaplan-Meier
17 estimator and the log-rank test. Significance was set at $p \leq 0.05$, and where
18 multiple comparisons were performed the Bonferroni correction was applied.
19
20 In addition to the Kaplan-Meier procedure, Cox proportional hazard models were
21 run using Proc Phreg (SAS Version 8.2) to determine the significance of any
22 variables other than genotype in the survivorship of non-scrapie infected
23 individuals. Selection of variables was made by looking for significant changes in

1 the log likelihood (χ^2) after using a hierarchical method of variable selection
2 (Collett, 2003). The following variables were tested for significance and model
3 improvement: year of birth, mode of feeding (maternal vs. artificial), and breeding
4 status (breeder, non-breeder). Genotype was added into the model last after other
5 significant variables were adjusted for. Significance was set at $p \leq 0.05$. Goodness
6 of fit of all models was examined by looking at the residuals.

7
8 **Variation in cause of removal.** Managers of the INRA Romanov flock kept
9 records on the reason for removal from the flock in addition to the date of
10 removal. The data can be grouped into the following three categories: Poor Health
11 (e.g., mastitis, arthritis, septicaemia, lungs, diarrhoea, toxaemia), Accidental (e.g.,
12 drowning, fracture, wound) and Management (e.g., culled for meat, sold, age-
13 related culling). Such data may provide information to indicate whether or not
14 there are any removals that may be attributed to pre-clinical scrapie. We
15 hypothesise that effects of pre-clinical scrapie would result in sheep with the
16 susceptible genotypes being removed significantly more for health-related causes
17 than sheep with non-affected genotypes. To test this hypothesis we examined the
18 causes of removal in the three most susceptible genotypes (ARQ/VRQ,
19 VRQ/VRQ and ARQ/ARQ) as well as the non-affected genotypes. Comparisons
20 of the number of removals of susceptible and non-affected genotypes within each
21 removal category were made using a χ^2 test or Fishers Exact test (if $n < 5$).
22 Analysis of frequency data was carried out in StatXact (Version 5.0). Statistical
23 significance was set at $p \leq 0.05$.

24

1 **RESULTS**

2

3 **Variation in lifetime breeding success**

4

5 **Association with scrapie status.** For both scrapie and non-scrapie infected sheep
6 the LBS was highest in the INRA Romanov sheep and lowest in the NPU Cheviot
7 I sheep (Table 3). For all flocks the lifetime breeding success of females that
8 developed scrapie was significantly lower than non-scrapie infected sheep
9 ($p \leq 0.001$), with the exception of the NPU Cheviot II flock ($n=10$, Table 3).

10 However, the power to detect differences in LBS within the NPU Cheviot II flock
11 was low .

12

13 Despite differences between flocks in the average number of offspring, the
14 percentage difference in the LBS between scrapie and non-scrapie infected ewes
15 was similar across all 5 flocks, with the scrapie ewes producing on average 34%
16 fewer offspring (Table 3). Combining the data from all five flocks, no significant
17 interaction between flock and scrapie status was found ($p=0.637$), implying no
18 difference between flocks in the reduction in breeding success due to scrapie.

19

20 **Association with PrP genotype.** For the INRA Romanov flock, we compared
21 LBS in scrapie and non-scrapie infected sheep within each of the three susceptible
22 genotypes (ARQ/ARQ, ARQ/VRQ and VRQ/VRQ) (Fig. 1). The INRA Romanov
23 flock had genotype information on all the scrapie infected sheep ($n=202$) and the

1 majority (67%; n=330/491) of non-scrapie infected sheep. Amongst susceptible
2 genotypes a GLM revealed significant effects of both status (scrapie infected
3 versus non-scrapie infected: $F_{1,360}=50.36$, $p<0.001$) and of genotype (ARQ/ARQ,
4 ARQ/VRQ and VRQ/VRQ; $F_{2,360}$, $p=0.004$) on LBS. There was no interaction
5 between the two factors ($p=0.709$), indicating that the proportionate reduction in
6 LBS due to scrapie did not differ between genotypes. As observed across the
7 entire flock, the LBS of scrapie infected sheep was significantly less than non-
8 scrapie infected sheep. Regardless of status, multiple comparison tests (with
9 Bonferroni correction) revealed that the LBS of VRQ/VRQ was significantly
10 lower than both ARQ/ARQ ($p=0.003$) and ARQ/VRQ ($p=0.022$) but there was no
11 significant difference between ARQ/ARQ and ARQ/VRQ ($p=0.815$).

12

13 Considering only non-scrapie sheep, there were differences between susceptible
14 and non-affected genotypes. A GLM analysis revealed significant genotype
15 effects ($F_{3,325}=3.70$, $p=0.012$). Multiple comparison (with Bonferroni correction)
16 revealed that the LBS of non-scrapie infected VRQ/VRQ sheep was significantly
17 less than the non-affected genotypes ($p=0.022$) and only marginally not
18 significantly different from the ARQ/ARQ non-scrapie infected sheep ($p=0.071$).
19 No other comparison was significant or approaching significance ($p>0.10$).

20

21 The NPU Cheviot II flock had genotype information on all scrapie infected sheep
22 (n=10), however, very few (18%; n=41/225) non-scrapie infected sheep were
23 genotyped. Despite the small sample size a GLM analysis of status (scrapie

1 infected versus non-scrapie infected) and genotype (ARQ/VRQ versus
2 VRQ/VRQ) was performed amongst susceptible genotypes. There was no
3 interaction between the two factors ($p=0.696$) and no significant status ($p=0.083$)
4 or genotype differences ($p=0.057$) although genotype tended towards significance,
5 with the LBS of VRQ/VRQ sheep less than that of sheep with the ARQ/VRQ
6 genotype. Considering only non-scrapie infected sheep, comparison of the LBS of
7 the three susceptible and non-affected genotypes revealed significant genotype
8 effects ($p=0.002$). Multiple comparisons (with Bonferroni correction) revealed
9 that the LBS of non-scrapie infected VRQ/VRQ sheep was significantly less than
10 the non-affected genotypes ($p=0.0036$).

11

12 Within the SAC Suffolk flock there was only one susceptible genotype
13 (ARQ/ARQ). As with the NPU Cheviot II flock, genotyping information was
14 limited. All scrapie infected sheep were genotyped, however, only 39%
15 ($n=211/537$) of the non-scrapie infected sheep were genotyped. Despite the small
16 sample size, a one-way ANOVA on differences in the LBS of scrapie infected
17 versus non-scrapie infected amongst ARQ/ARQ genotypes revealed no significant
18 difference between scrapie infected and non-scrapie infected sheep within the
19 susceptible genotype ARQ/ARQ ($p=0.563$). Considering only non-scrapie
20 infected sheep, there were significant differences between susceptible
21 (ARQ/ARQ) and non-affected genotypes ($p<0.001$) where ARQ/ARQ sheep had a
22 significantly lower LBS than the non-affected sheep.

23

1 **Variation in Litter Size**

2

3 **Association with scrapie status.** The largest litter sizes were observed in the
4 INRA Romanov flock and the smallest in the NPU Cheviot I flock (Table 4).

5 There were no significant differences between the size of litters from scrapie
6 infected and non-scrapie infected dams in each flock (Table 4).

7

8 **Association with PrP genotype.** Amongst susceptible genotypes in the INRA
9 Romanov and SAC Suffolk flocks, there were no differences in litter size between
10 sheep that developed scrapie and those that did not (INRA: $F_{1,302}=0.973$, $p=0.325$;
11 SAC: $F_{1,76}=1.584$, $p=0.212$). Considering only sheep that never developed scrapie,
12 there were also no significant differences between non-affected and susceptible
13 genotypes (INRA: $F_{1,693}=0.90$, $p=0.346$; SAC: $F_{1,76}=1.584$, $p=0.212$).

14

15 **Variation in Survival**

16

17 **Association with scrapie status.** For all 5 flocks there was a significant reduction
18 in the survival time (age at removal) of scrapie infected individuals relative to
19 non-scrapie infected individuals (Table 5). The INRA Romanov had the largest
20 difference between median survival of scrapie infected and non-scrapie infected
21 sheep (4.3 years) whereas the NPU Cheviot I had the lowest (1.4 years).

22

1 **Association with PrP genotype (non-scrapie infected sheep only).** For the
2 INRA Romanov flock both the Kaplan Meier ($\chi^2=39.23$, $df=3$, $p<0.001$; Figure
3 2A) and Cox proportional hazards genotype-only model revealed significant
4 differences among the 4 genotype groups ((VRQ/VRQ = ARQ/VRQ) <
5 (ARQ/ARQ = non-affected)) in the age at removal of non-scrapie infected sheep.
6 As such, the following groups of genotypes were formed: highly susceptible
7 (VRQ/VRQ + ARQ/VRQ) and other (ARQ/ARQ + non-affected). This was done
8 to increase the power of the analysis as the sample size of VRQ/VRQ non-scrapie
9 infected individuals was very low. Diagnostic checks on the Cox proportional
10 hazards model with covariates revealed a violation of the assumption of
11 proportional hazards. This appeared to be the result of increased risk of early
12 death for the highly susceptible VRQ/VRQ and ARQ/VRQ individuals after 2
13 years. As such, a piecewise Cox model was applied, comparing age at removal for
14 the different genotype groups (highly susceptible, other) before and after 2 years.
15 The results show that there is a significant genotype effect even after adjustment
16 for significant variables: year of birth, breeding status, and breeding status by
17 genotype interaction (Table 6), however, only for individuals after 2 years. There
18 was no difference in the risk of removal between the genotype groups prior to 2
19 years. Sheep with the highly susceptible genotypes, VRQ/VRQ and ARQ/VRQ,
20 had a 14x higher risk of an early death.
21
22 For the NPU Cheviot II Flock both the Kaplan Meier ($\chi^2=23.7$, $df=2$, $p<0.001$;
23 Figure 2B) and Cox proportional hazards genotype-only model revealed

1 significant differences among the 3 genotype groups (VRQ/VRQ, ARQ/VRQ and
2 non-affected) in the age at removal of nonscrapie infected sheep. The risk of early
3 death for sheep with genotype VRQ/VRQ was 4.2x higher than for non-affected
4 sheep ($p < 0.001$). The risk of early death for sheep with genotype ARQ/VRQ was
5 2.7x higher than for non-affected sheep ($p = 0.001$). The only other variable that
6 was significant was year of birth. Addition of this variable did not change the
7 significance of genotype in the model.

8

9 In the SAC Suffolk flock both the Kaplan Meier ($\chi^2 = 3.90$, $df = 1$, $p = 0.048$; Figure
10 2C) and Cox proportional hazards genotype-only model revealed significant
11 differences among the 2 genotype groups (ARQ/ARQ and non-affected) in the
12 age at removal of nonscrapie infected sheep. The risk of early death for sheep
13 with genotype ARQ/ARQ was 1.5x higher than non-affected sheep but the
14 significance was marginal ($p = 0.049$). However, adjusting for significant variables
15 (i) year of birth and (ii) breeding status revealed that differences between
16 genotypes ARQ/ARQ and non-affected were significant ($p = 0.010$).

17

18 **Variation in cause of removal**

19

20 For all 3 flocks examined there were genotype differences in the life expectancy
21 of the sheep. Overall, sheep with highly susceptible genotypes did not live as long
22 as sheep with non-affected and/or less susceptible genotypes. Examination of the
23 distribution of age at death from scrapie (Figure 2A-C) revealed similarity

1 between the three flocks. The peak in scrapie deaths approximates the point at
2 which 50% of the susceptible yet non-scrapie infected animals in the flock are
3 being removed (Figure 2A-C). For example, in the INRA Romanov flock mean
4 age of scrapie deaths is approximately 2 years of age, with all scrapie deaths
5 occurring before age 4. In the survival graph for non-scrapie infected deaths all
6 VRQ/VRQ and ARQ/VRQ die within 4 years, whereas the less susceptible
7 ARQ/ARQ and non-affected genotypes have a maximum lifespan of 9 years
8 (Figure 2A). A similar pattern can be observed for VRQ/VRQ and ARQ/VRQ
9 sheep in the NPU Cheviot II flock and the ARQ/ARQ in the SAC Suffolk flock.
10

11 For the INRA Romanov flock the presence of the VRQ allele appears to be a
12 significant factor in the age at removal of non-scrapie infected sheep in flocks
13 affected by scrapie. The cause of this lower mean life expectancy in ARQ/VRQ
14 and VRQ/VRQ sheep in the presence of scrapie suggests pre-clinical scrapie
15 amongst the most susceptible genotypes. To explore this hypothesis further we
16 examined the causes of death in non-scrapie infected sheep with the highly
17 susceptible genotypes (VRQ/VRQ and ARQ/VRQ) versus other non-scrapie
18 infected sheep with ARQ/ARQ and non-affected genotypes . A greater proportion
19 of animals with the highly susceptible genotypes were removed for health-related
20 reasons ($\chi^2=41.11$, $df=1$, $p<0.001$), whereas animals with ARQ/ARQ and non-
21 affected genotypes were more likely to be removed for management reasons
22 ($\chi^2=38.56$, $df=1$, $p<0.001$). There was no significant difference between the

1 genotype groups for the proportion of animals removed for accidental causes
2 ($p > 0.05$).

3

4

5 **DISCUSSION**

6

7 We have used detailed individual-level analyses of outbreaks of natural scrapie in
8 five sheep flocks to quantify the effects of scrapie and of PrP genotype on
9 individual fitness. Despite different breed composition and scrapie incidence, we
10 found consistent patterns in lifetime breeding success, litter size and sheep
11 survival among the flocks.

12

13 There were significant differences in lifetime breeding success (LBS) of scrapie
14 and non-scrapie infected sheep within the 4 flocks where there was sufficient data
15 to examine the comparison, with scrapie sheep producing on average 34% fewer
16 offspring than non-scrapie infected sheep. However, despite differences in the
17 average LBS measured in each flock, there was no evidence of any difference
18 between flocks in proportionate reduction in LBS due to scrapie. There is
19 therefore no indication of any variation between sheep breeds in loss of fitness
20 due to scrapie infection. In addition to the overall effect of scrapie, there were also
21 genotype differences in the LBS of scrapie and non-scrapie sheep which
22 correlated with the susceptibility of the genotype ($VRQ/VRQ < ARQ/VRQ <$
23 $ARQ/ARQ < \text{non-affected}$). This could only be examined in detail for the INRA

1 Romanov flock, but a similar pattern was apparent in the NPU Cheviot II and
2 SAC Suffolk flocks.

3

4 The effect of scrapie on lifetime breeding success appears to be a function of
5 lifespan as opposed to fecundity. Analysis of litter size revealed no overall or
6 genotype differences among the 5 sheep flocks. However, significant differences
7 in survival of sheep were identified in this study. In general, age at removal from
8 the flock depends on individual status (i.e. scrapie infection) and PrP genotype.

9 For the five flocks examined, the median age at which scrapie infected sheep were
10 removed from the flock was significantly less than that for non-scrapie infected
11 sheep. Reduced survival in scrapie sheep was expected based on previous research
12 where lower life expectancies were observed for the most susceptible sheep in the
13 flocks (Bossers *et al.* 1996; Clouscard *et al.* 1995; Elsen *et al.* 1999;
14 Thorgeirsdottir *et al.* 2002). As such, differences in the survival of scrapie
15 affected sheep was not analysed in detail in this study. The focus of the survival
16 analysis in this study was on non-scrapie infected sheep. The results of the
17 Survival analysis and Cox Proportional Hazard model indicated significant
18 genotype differences in the pattern of survival among the non-scrapie infected
19 individuals for the flocks examined. Even when adjustment is made for significant
20 covariates, there was an increased risk of removal associated with susceptible
21 genotypes. For the INRA Romanov flock this seemed to depend on genotype or
22 genotype susceptibility. VRQ/VRQ and ARQ/VRQ genotype individuals had

1 significantly lower life-expectancies, whereas the life-expectancy of ARQ/ARQ
2 genotyped sheep were not significantly different from non-affected sheep.
3

4 The distribution of removals from each flock approximates the age distribution of
5 scrapie deaths. This distribution suggests that although scrapie was not diagnosed,
6 these sheep were removed because of scrapie that was not detected or other
7 health-related causes associated with scrapie incubating within the sheep. Reports
8 from other field studies are inconsistent. McLean *et al.* (1999) reported having
9 more sheep die of unknown causes on scrapie affected farms than scrapie-free
10 farms. Baylis *et al.* (2002) also observed in scrapie-affected sheep flocks a
11 number of sheep that were found dead of unknown causes (8% of entire flock) but
12 there was not a significant association with scrapie risk. In a recent study,
13 however, a high prevalence of scrapie (6%) was observed amongst sheep that
14 were found-dead in Shetland where scrapie is very common (Humphry *et al.*
15 2004).

16

17 For the INRA Romanov flock a significantly higher proportion of ARQ/VRQ and
18 VRQ/VRQ sheep died of poor health in comparison to ARQ/ARQ and the non-
19 affected genotypes. One would have expected that if removals were the result of
20 pre-clinical scrapie that sheep with the ARQ/ARQ genotype would also have a
21 high proportion of removal as a result of health-related illness. It appears that
22 there may be a deleterious effect of the presence of the VRQ allele in the presence
23 of scrapie in the flock. Unfortunately there were no equivalent data from the other

1 flocks with which to test this idea. The results of this study suggest that, across
2 different flocks of different sheep breeds, susceptible PrP genotypes appear to
3 perform less well in overall fecundity and life-expectancy even if they do not
4 contract scrapie. This effect is more apparent in the most susceptible genotype:
5 VRQ/VRQ performed consistently worse in relation to lifetime breeding success
6 and survival even amongst apparently uninfected individuals.

7
8 There are two possible explanations for these findings. The first is that susceptible
9 genotypes are in relatively poorer condition and are removed at younger ages.
10 Unfortunately lack of data makes this hypothesis difficult to examine, although
11 research to date suggests that there are no PrP genotype-related performance traits
12 (Roden *et al.* 2001; Barillet *et al.* 2002; Brandsma *et al.* 2004; DeVries *et al.*
13 2004). The second hypothesis is that they are suffering from effects of pre-clinical
14 scrapie, which is manifesting itself in terms of reduced lifespan even though
15 typical clinical signs of scrapie are yet to develop. If this hypothesis were true we
16 might expect: (1) most deaths in years 2-4 when most scrapie cases occur; and (2)
17 the cause of death for susceptible genotypes to be different (i.e. more health-
18 related). Both expectations are confirmed by the results reported within this
19 paper, although the results for the susceptible genotype ARQ/ARQ in the INRA
20 Romanov flock are not as clear. Physiological evidence of pre-clinical scrapie
21 does exist. Changes in behaviour that appear to consistently precede clinical signs
22 have been observed (Parry, 1983). Studies have shown that there was reduced
23 rumination in sheep with scrapie and cattle with BSE (Austin & Simmons, 1993).

1 This reduced rumination may provide an explanation for the observations of loss
2 of weight or body condition that has been reported for scrapie (Clark & Moar,
3 1992), BSE (Wilesmith *et al.* 1992) and chronic wasting disease (Williams &
4 Young, 1982).

5

6 Scrapie has become the target of control measures and eradication programs
7 world wide. The identification of infected sheep is crucial for the success of these
8 programs. After initial infection, the disease has a long incubation period during
9 which time infected sheep may be able to transmit disease to non-infected sheep.
10 Evidence of scrapie can now be detected in sheep before the clinical signs occur
11 (e.g. Schreuder *et al.* 1998) but it is unknown whether or not sheep are affected
12 during this 'pre-clinical' phase. This study has suggested the possibility that
13 reduced lifespan in susceptible PrP genotypes may be the result of pre-clinical
14 scrapie. If pre-clinical scrapie does exist amongst susceptible genotypes we may
15 underestimate levels of scrapie-related mortality in sheep flocks. The results
16 presented here highlight the need for further research on performance of different
17 sheep PrP genotypes both in the presence and absence of scrapie.

18

19 **ACKNOWLEDGEMENTS**

20

21 This work was funded primarily by the Biotechnology and Biological Sciences
22 Research Council. The authors would also like to thank the contributions from the
23 Wellcome Trust and the Royal Society. SAC receives financial support from the

1 Scottish Executive Environment and Rural Affairs Department. Special thanks to
2 Darren Shaw for assistance with the illustrations in the manuscript.

3

4 **REFERENCES**

5

6 **Acín, C., Martín-Burriel, I., Goldmann, W., & 7 other authors (2004).** Prion
7 protein gene polymorphisms in healthy and scrapie-affected Spanish sheep. *J.*
8 *Gen. Virol.* **85**, 2103-2110.

9

10 **Austin, A.R. & Simmons, M.M. (1993).** Reduced rumination in bovine
11 spongiform encephalopathy and scrapie. *Vet. Rec.* **132**, 324-325.

12

13 **Barillet, F., Andréoletti, O., Palhiere, I. & 12 other authors (2002).** Breeding
14 for scrapie resistance using *PrP* genotyping in the French dairy sheep breeds,
15 Proceedings of the 7th World Congress on Genetics Applied to Livestock
16 Production, Montpellier, France, 2002, pp. 19-23.

17

18 **Baylis, M., Houston, F., Goldmann, W., Hunter, W. & McLean, A.R. (2000).**
19 The signature of scrapie: differences in the PrP genotype profile of scrapie-
20 affected and scrapie-free UK sheep flocks. *Proc.R. Soc. Lond. B Biol. Sci.* **267**,
21 2029-2035.

22

1 **Baylis, M., Goldmann, W., Houston, F., Cairns, D., Chong, A., Ross, A.,**
2 **Smith, A., Hunter, N. & McLean, A.R. (2002).** Scrapie epidemic in a fully PrP-
3 genotyped sheep flock. *J. Gen. Virol.* **83**, 2907-2914.
4
5 **Bossers, A., Schreuder, B.E.C., Muileman, I.H., Belt, P.B.G.M. & Smits,**
6 **M.A. (1996).** PrP genotypes contributes to determining survival times of sheep
7 with natural scrapie. *J. Gen. Virol.* **77**, 2669-2673.
8
9 **Brandsma, J.H., Janss, L.L.G. & Visscher, A.H. (2004).** Association between
10 PrP genotypes and litter size and 135 days weight in Texel sheep. *Livest. Prod.*
11 *Sci.* **85**, 59-64.
12
13 **Clark, A.M. & Moar, J.A.E. (1992).** Scrapie – a clinical assessment. *Vet. Rec.*
14 **130(17)**, 377-378.
15
16 **Cloucard, C., Beaudry, P., Elsen, J-M., Milan, D., Dussaucy, M., Bounneau,**
17 **C., Schelcher, F., Chatelain, J., Launay, J.M. & Laplanche, J.L. (1995).**
18 Different allelic effects of the codons 136 and 171 of the prion protein gene in
19 sheep with natural scrapie. *J. Gen. Virol.* **76**, 2097-2101.
20
21 **Collett, D. (2003).** *Modelling Survival Data in Medical Research*, 2nd edn.
22 Chapman and Hall. London.
23

1 **Dawson, M., Hoinville, L.J., Hosie, B.D. & Hunter, N. (1998).** Guidance on the
2 use of PrP genotyping as an aid to the control of clinical scrapie. *Vet. Rec.* **142**,
3 623-625.
4

5 **De Vries, F., Borchers, N., Hamann, H., Drogemuller, C., Reinecke, S.,**
6 **Lupping, W. & Distl, O. (2004).** Associations between prion protein genotype
7 and performance traits of meat breeds of sheep. *Vet. Rec.* **155(5)**,140-143.
8 **Drogemuller, C., Leeb, T. & Distl, O. (2001).** PrP genotype frequencies in
9 German breeding sheep and the potential to breed for resistance to scrapie. *Vet.*
10 *Rec.* **149**, 349-352.
11

12 **Elsen, J.M., Amigues, Y., Schelcher, F., Ducrocq, V., Andréoletti, O.,**
13 **Eychenne, F., Khang, J.V.T., Poivey, J.P., Lantier, F. & Laplanche J.L.**
14 **(1999).** Genetic susceptibility and transmission factors in scrapie: detailed
15 analysis of an epidemic in a closed flock of Romanov. *Arch. Virol.* **144**, 431-445.
16

17 **Goldmann, W., Hunter, N., Benson, G., Foster, J.D. & Hope J. (1991).**
18 Different scrapie-associated fibril proteins (PrP) are encoded by lines of sheep
19 selected for different alleles of the *Sip* gene. *J. Gen. Virol.* **72**, 2411-2417.
20

21 **Goldmann, W., Hunter, N., Smith, G., Foster, J. & Hope, J. (1994).** PrP
22 genotype and agent effects in scrapie – change in allelic effects with different
23 isolates of agent in sheep, a natural host of scrapie. *J. Gen. Virol.* **75**, 989-995.

1

2 **Humphry, R.W., Clark, A.M., Begara-McGorum, I. & Gunn, G.J. (2004).**
3 Estimation of scrapie prevalence in cull and found-dead sheep on the Shetland
4 Islands. *Vet. Rec.* **156**, 303-304.

5

6 **Hunter, N., Foster, J.D., Goldmann, W., Stear, M.J., Hope, J. & Bostock, C.**
7 **(1996).** Natural scrapie in a closed flock of Cheviot sheep occurs only in specific
8 PrP genotypes. *Arch. Virol.* **141**, 809-824.

9

10 **Hunter, N., Moore, L., Hosie, B.D., Dingwall, W.S. & Greig, A. (1997).**
11 Association between natural scrapie and PrP genotype in a flock of Suffolk sheep
12 in Scotland. *Vet. Rec.* **140**, 59-63.

13

14 **Hunter, N. (1997).** PrP genetics in sheep and the applications for scrapie and
15 BSE. *Trends Microbiol.* **5**, 331-334.

16

17 **McLean, A.R., Hoek, A., Hoinville, L.J. & Gravenor, M.B. (1999).** Scrapie
18 transmission in Britain: a recipe for a mathematical model, *Proc.R. Soc. Lond. B*
19 *Biol. Sci.* **266**, 2531-2538.

20

21 **Parry, H.B. (1983).** *Scrapie disease in sheep.* Academic Press, London.

22

- 1 **Prokopova, L., Lewis, R.M., Dingwall, W.S. & Simm, G. (2002).** Scrapie
2 genotype: a correlation with lean growth rate? Proceedings of the 7th World
3 Congress on Genetics Applied to Livestock Production, Montpellier, France,
4 2002, pp. 19-23.
- 5
- 6 **Redman, C.A., Coen, P.G., Matthews, L., Lewis, R.M., Dingwall, W.S.,**
7 **Foster, J.D., Chase-Topping, M.E., Hunter, N. & Woolhouse, M.E.J. (2002).**
8 Comparative epidemiology of scrapie outbreaks in individual sheep flocks.
9 *Epidemiol. Infect.* **128**, 513-521.
- 10
- 11 **Roden, J. A., Haresign, W., & Anderson, J.M.L. (2001).** Analysis of *PrP*
12 genotype in relation to performance traits in Suffolk sheep, British Society of
13 Animal Science: Proceedings of the Annual Meeting, pp. 45.
- 14
- 15 **Schreuder, B.E.C., van Keulen, L.J.M., Vromans, M.E.W., Langeveld, J.P.M.**
16 **& Smits, M.A. (1998).** Tonsillar biopsy and PrPSv detection in the preclinical
17 diagnosis of scrapie. *Vet. Rec.* **142(21)**, 564-568.
- 18
- 19 **Stamp, J.T. (1962).** Scrapie: a transmissible disease of sheep. *Vet. Rec.* **74**, 357-
20 362.
- 21
- 22 **Thorgeirsdottir, S., Georgsson, G., Reynisson, E., Sigurdarson, S. &**
23 **Palsdottir, A. (2002).** Search for healthy carriers of scrapie: an assessment of

1 subclinical infection of sheep in an Icelandic scrapie flock by three diagnostic
2 methods and correlation with PrP genotypes. *Arch. Virol.* **147**, 709-722.

3

4 **Vaccari, G., Petraroli, R., Agrimi, U., Eleni, C., Perfetti, M.G., Di Bari, M.A.,**
5 **Morelli, L., Ligios, C., Busani, L., Nonno, R. & Di Guardo, G. (2001).** PrP
6 genotype in Sarda breed sheep and its relevance to scrapie. *Arch. Virol.* **146**,
7 2029-2037.

8

9 **Wilesmith, J.W., Hoinville, L.J., Ryan, J.B.M. & Sayers, A.R. (1992).** Bovine
10 spongiform encephalopathy – aspects of the clinical picture and analysis of
11 possible changes 1986-1990. *Vet. Rec.* **130**, 197.

12

13 **Williams, E.S. & Young, S. (1982).** Spongiform encephalopathy of rocky
14 mountain elk. *J. Wildl. Dis.* **18**, 465.

1 **Figure Legends**

2

3 Fig. 1. Differences in Lifetime Breeding Success (LBS) within scrapie susceptible
4 genotypes (ARQ/ARQ, ARQ/VRQ, VRQ/VRQ) and non-affected genotypes in
5 the INRA Romanov flock.

6

7 Fig. 2. Foreground: Survivorship. Age at removal for female non-scrapie infected
8 sheep. A. INRA Romanov flock for susceptible (VRQ/VRQ, ARQ/VRQ,
9 ARQ/ARQ) and non-affected genotypes; B. NPU Cheviot II flock for the 2
10 susceptible (VRQ/VRQ and ARQ/VRQ) and non-affected genotypes; C. SAC
11 Suffolk flock for the susceptible (ARQ/ARQ) genotype and the non-affected
12 genotypes. Background: Distribution of the age of scrapie deaths for females in A.
13 the INRA Romanov flock; B. the NPU Cheviot II flock; and C. the SAC Suffolk
14 flock. VRQ/VRQ: bold, black; VRQ/ARQ: bold, grey; ARQ/ARQ: normal, black;
15 Non-affected: normal, grey.

Table 1. Demographic characteristics of the study flocks with outbreaks of natural scrapie. Outbreak, calendar years over which cases of natural scrapie were observed. Cohorts, birth cohorts involved in the outbreak of natural scrapie.

Flock	Organisation	Country	Breed	Research	Outbreak (years)	Cohorts (years)	Range of flock size per year *	No. sheep*	No. cases*
NPU Cheviot I	Institute of Animal Health, Neuropathogenesis Unit (NPU)	Scotland	Cheviot	Scrapie	1970-1982	1967-1978	273-751	1321	137
NPU Cheviot II	Institute of Animal Health, Neuropathogenesis Unit (NPU)	Scotland	Cheviot	Scrapie	1986-1994	1982-1994†	304-653	1604	33
NPU Suffolk	Institute of Animal Health, Neuropathogenesis Unit (NPU)	Scotland	Suffolk	Scrapie	1959-1982	1956-1980	43-597	1658	710
SAC Suffolk	Scottish Agricultural College (SAC)	Scotland	Suffolk	Meat	1990-1996	1988-1994	198-760	2489	108
INRA Romanov	Institut National de la Recherche Agronomique (INRA)	France	Romanov	Fecundity Meat Scrapie‡	1993-1999	1986-1999†	390-792	5841	448

*, all values except INRA Romanov are from Redman *et al.* 2002. INRA Romanov data was calculated from INRA database.

†, data collection in these flock are ongoing. For the purpose of this research the database was closed in 1994 and 1999 for the NPU Cheviot II and INRA Romanov flocks respectively.

‡, research on scrapie began in 1993 after the first case of scrapie was observed in the flock.

Table 2. Susceptible and non-affected genotypes within the INRA Romanov, NPU Cheviot II and SAC Suffolk flocks. Susceptible genotypes are presented in order of decreasing susceptibility. Scrapie susceptibility, expressed as % of genotype affected, is shown in brackets.

NPU Cheviot II		SAC Suffolk		INRA Romanov	
Susceptible	Non-affected	Susceptible	Non-affected	Susceptible*	Non-affected [†]
VRQ/VRQ (56%)	AHQ/AHQ	ARQ/ARQ (58%)	ARQ/ARH	VRQ/VRQ (76%)	AHQ/AHQ
ARQ/VRQ (33%)	AHQ/VRQ		ARR/ARH	ARQ/VRQ (52%)	AHQ/VRQ
	ARQ/AHQ		ARR/ARQ	ARQ/ARQ (42%)	ARQ/AHQ
	ARQ/ARQ		ARR/ARR		ARR/AHQ
	ARR/AHQ				ARR/ARQ
	ARR/ARQ				ARR/ARR
	ARR/ARR				ARR/VRQ
	ARR/VRQ				

*, Data on scrapie susceptibility from Elsen *et al.* 1999.

[†], Includes some suspect scrapie cases in all genotypes except ARR/ARR and ARR/AHQ (Elsen *et al.* 1999)

Table 3. Summary of the Lifetime Breeding Success (LBS) of scrapie and non-scrapie infected females in each flock, t statistic and corresponding p-value to test for differences between the two categories and the difference between scrapie and non-scrapie infected individuals within each flock. Cohorts used in the analysis are in brackets.

Flock	Status	n	Lifetime Breeding Success			t	p	% Difference
			Mean	SE	Range			
NPU Cheviot I (1967-1978)	Non-scrapie	208	2.9	0.1	1 - 8	5.14	<0.001	-27.0
	Scrapie	66	1.8	0.1	1 - 5			
NPU Cheviot II (1982-1994)	Non-scrapie	225	4.8	0.2	1 - 20	1.19	0.237	-34.0
	Scrapie	10	3.3	0.6	1 - 7			
NPU Suffolk (1956-1980)	Non-scrapie	191	3.2	0.2	1 - 10	5.43	<0.001	-40.0
	Scrapie	270	2.1	0.1	1 - 7			
SAC Suffolk (1988-1994)	Non-scrapie	537	4.8	0.1	1 - 27	4.17	<0.001	-31.0
	Scrapie	56	3.5	0.4	1 - 17			
INRA Romanov (1986-1993)	Non-scrapie	491	16.5	0.4	1 - 40	8.76	<0.001	-38.0
	Scrapie	202	10.3	0.6	1 - 37			

Table 4. Summary of the mean size of all litters born to scrapie and non-scrapie infected dams in each flock during the scrapie outbreak (years are in brackets). n_d is number of dams; n_l is number of litters. F statistic is from generalised linear mixed effects model with dam identity as random effect and Poisson errors (with corresponding degrees of freedom, df, and p-value).

Flock	Status	n_d (n_l)	Mean Litter Size		F (df)	p
			Mean	SE		
NPU Cheviot I (1970-1982)	Non-scrapie	248 (472)	1.34	0.02	0.060 (1,311)	0.807
	Scrapie	65 (90)	1.23	0.04		
NPU Cheviot II (1986-1994)	Non-scrapie	242 (566)	1.62	0.02	0.001 (1,233)	0.993
	Scrapie	10 (19)	1.58	0.12		
NPU Suffolk (1959-1982)	Non-scrapie	274 (870)	1.78	0.02	2.383 (1,535)	0.123
	Scrapie	292 (606)	1.65	0.02		
SAC Suffolk (1990-1996)	Non-scrapie	694 (1561)	1.76	0.02	0.455 (1,749)	0.500
	Scrapie	57 (107)	1.81	0.09		
INRA Romanov (1993-1999)	Non-scrapie	547 (917)	3.28	0.03	1.430 (1,661)	0.232
	Scrapie	114 (150)	3.43	0.08		

Table 5. Median survival times (\pm 95% CI) for scrapie and non-scrapie infected sheep in each flock. Cohorts used in the analysis are in brackets.

Flock	Status	Median	95% CI	χ^2	p
NPU Cheviot I (1970-1778)	Scrapie	2.24	2.14 – 2.49	129	<0.001
	Non-scrapie	3.80	3.53 – 4.46		
NPU Cheviot II (1986-1994)	Scrapie	2.41	2.17 – 3.32	31	<0.001
	Non-scrapie	6.07	5.74 – 6.57		
NPU Suffolk (1959-1980)	Scrapie	2.82	2.67 – 2.90	226	<0.001
	Non-scrapie	5.04	4.35 – 5.26		
SAC Suffolk (1990-1996)	Scrapie	2.63	2.07 – 2.60	164	<0.001
	Non-scrapie	4.64	4.19 – 4.89		
INRA Romanov (1993-1999)	Scrapie	1.76	1.75 – 1.82	484	<0.001
	Non-scrapie	6.02	5.35 – 6.30		

Table 6. Piecewise Cox Proportional Hazard model for mean life expectancy of non-scrapie infected sheep in the INRA Romanov flock. Risk ratio, $\exp(\text{parameter estimate})$. 95% CI, $\exp(\text{parameter estimate} \pm 1.96 (\text{SE}))$. YOB, year of birth. NB, non-breeder. Other, ARQ/ARQ and non-affected genotypes. Highly susceptible, VRQ/VRQ and ARQ/VRQ genotypes. Baseline, genotype other, breeder, YOB 1993.

Model 2: Genotype + covariates						
Variable	Parameter	SE	Wald Chi square	p	Risk ratio	95% CI
Non-Breeder (NB)	1.566	0.252	38.57	<0.001	4.79	2.92 – 7.85
YOB 1994	0.022	0.203	0.01	0.913	1.02	0.687 – 1.52
YOB 1995	0.093	0.262	0.13	0.722	1.10	0.656 – 1.84
YOB 1996	0.302	0.227	1.77	0.183	1.35	0.867 – 2.11
YOB 1997	-0.249	0.293	0.73	0.394	0.78	0.439 – 1.38
YOB 1998	-1.197	0.376	10.2	0.001	0.30	0.145– 0.631
YOB 1999	-0.495	0.364	1.85	0.174	0.61	0.298 – 1.24
Non-Breeder, Other	-0.888	0.366	5.89	0.015	0.41	0.201 – 0.843
Highly susceptible < 2yrs	0.400	0.351	1.29	0.255	1.49	0.749 – 2.97
Highly susceptible > 2yrs	2.702	0.320	71.3	<0.001	14.91	7.96 – 27.9

