

Nutrient dynamics in a chronosequence of douglas-fir (Pseudotsuga menziesii (Mirb.) Franco) stands on the Beaujolais mounts (France). 2. Quantitative approach

Renato Marques, Jacques J. Ranger, Sylvain Villette, André Granier

▶ To cite this version:

Renato Marques, Jacques J. Ranger, Sylvain Villette, André Granier. Nutrient dynamics in a chronosequence of douglas-fir (Pseudotsuga menziesii (Mirb.) Franco) stands on the Beaujolais mounts (France). 2. Quantitative approach. Forest Ecology and Management, 1997, 92 (1-3), pp.167-197. 10.1016/S0378-1127(96)03913-8. hal-02685561

HAL Id: hal-02685561 https://hal.inrae.fr/hal-02685561

Submitted on 1 Jun2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Forest Ecology and Management 92 (1997) 167-197

Forest Ecology and Management

Nutrient dynamics in a chronosequence of Douglas-fir (*Pseudotsuga menziesii* (Mirb). Franco) stands on the Beaujolais Mounts (France). 2. Quantitative approach

Renato Marques, Jacques Ranger *, Sylvain Villette, André Granier

INRA-Centre de Nancy, Equipe Cycles Biogeochimiques, Champenoux, 54280 Seichamps, France

Accepted 5 August 1996

Abstract

Nutrient fluxes were determined in a chronosequence of Douglas-fir stands in the Beaujolais Mounts (France). Annual and seasonal variations occurred during the 3 years of investigation; fluxes were generally highest in autumn-winter.

Atmospheric inputs were among the mean values from a monitoring network of forest ecosystems in France. Nutrient outputs from the soil profile were higher than average and occurred mainly during vegetation dormancy.

Mean input-output budgets were negative for N, S, K, Ca and Mg, characterising an imbalance of the site chemistry dynamics. The P budget was positive. Most of the nutrient output from the ecosystem occurred as losses in the drainage water. These losses were related to excess nitrification and consecutive cation mobilisation throughout the soil profile.

Surface water, however, had a neutral pH and very low nitrate and aluminium contents, which may have been buffered by the subsoil.

Budgets differed between stands and tended to be more negative in the youngest stand. Part of this behaviour was related to stand age and part to the former land use of plots. Theoretical budgets were calculated for forest rotation lengths of 20, 40 and 60 years; it was concluded that shorter rotations would increase nutrient losses. The trend of a decrease in budget deficits with stand age suggests that the effect of vegetation change will be reduced at the next rotation but the impact of stand development may remain.

Predicted nutrient budgets for a second 60 year Douglas-fir rotation suggested that available Ca in the soil would be depleted and that this depletion would be even more drastic if whole tree harvesting were adopted.

In conclusion, Douglas-fir stands introduced changes in soil function that may impoverish the soil if present trends remain the same over the next forest rotations. The maintenance of sustainability will require nutrient input by fertilisation.

Keywords: Douglas-fir; Nutrients; Biogeochemical cycle; Input-output budgets; Dynamics; Chronosequence; Gravitational soil solutions

^{*} Corresponding author. Tel.: 33 83 39 40 68; fax: 33 83 39 40 69.

^{0378-1127/97/\$17.00} Copyright © 1997 Elsevier Science B.V. All rights reserved. PII \$0378-1127(96)03913-8

1. Introduction

Douglas-fir was first introduced in France 150 years ago. It is presently one of the most intensively planted species in the country owing to its high potential for wood production and its well appreciated wood quality.

Sustainable management of these stands needs to be demonstrated, not only in terms of maintaining the present forest production level but also with respect to maintaining soil fertility and surface water quality. "Wise management of production forests is essential if we are to ensure that future generations will be able to derive the same, if not more, resources from the land than were derived by preceding generations" (Beets et al., 1994).

Land occupation by new vegetation may result in dramatic changes in soil dynamics (Page, 1970; Feger, 1988; Ranger and Nys, 1994). Thus, a research programme was initiated in 1991 by selecting a chronosequence of Douglas-fir stands able to represent ecosystem dynamics throughout a whole forest rotation.

The first results described the dynamics of biomass and nutrient accumulation in the stand (Ranger et al., 1995). Relevant conclusions were obtained by comparing Douglas-fir with other species or by testing different silvicultural scenarios, i.e. changing the rotation length or changing the intensity of forest harvesting. Nutrient fluxes through the ecosystem were then investigated to calculate the input–output budget of the ecosystem. The first step was to qualify the chemistry of solutions transferred within these Douglas-fir stands and the processes involved in the acquisition of the chemistry and on the dynamics of these solutions (Marques and Ranger, 1996).

The objective of the present paper is to quantify the main fluxes of the biogeochemical cycle in the forest stands and to establish input-output budgets for individual stands and for different lengths of forest rotation. The results are discussed by focusing on the possible impact of Douglas-fir on soil and surface water quality.

2. Materials and methods

Only the essentials are indicated here as a detailed description of the site and instruments has already

been published in a companion paper (Marques and Ranger, 1996).

2.1. Site characterisation

Three stands aged 20, 40 and 60 years representing a forest chronosequence were chosen on the Beaujolais Mounts in France $(46^{\circ}30'N, 4^{\circ}38'E)$.

Stand characteristics were: 922, 350 and 225 trees ha⁻¹; 14.3, 28.2 and 36.3 m average height; 18.1, 33.3 and 52.1 cm average DBH; 250, 670 and 1100 m³ ha⁻¹ average volume of total stemwood. The 20and 40-year-old stands presented no understorey; the understorey of the 60-year-old stand was mainly composed of blackberry (*Rubus fruticosus* L.). The nutrient content and mean current immobilisation of stands (excluding understorey) is summarised in Table 1 (Ranger et al., 1995). Mean root distribution for the three stands is as follows: 33% in the 0–15 cm layer, 25% in the 15–30 cm layer. 28% in the 30–60 cm layer and 14% below 60 cm.

The site has an elevation of 750 m, a mean annual temperature of 7°C and mean annual rainfall of 1000 mm, which is more or less evenly distributed throughout the year with a slight maximum in summer.

2.2. Soil characterisation

Soils were typic Dystrochrepts (Soil Survey Staff, 1975). Soil texture was loam (40- and 60-year-old stands) and silt-loam (20-year-old stand) in the top 80 cm layer, and loam (all stands) in the deepest horizons. The organic matter content of the upper soil horizons was rather high in all stands and somewhat higher in the 20-year-old stand. Organic nitrogen followed the same tendency. Total Ca and Mg were somewhat higher and K lower in the soil of the 20-year-old stand than in the older stands. The pH showed no differences between stands. The young stand had higher exchangeable Ca²⁺ and Mg²⁺ concentrations in the upper soil horizons than the other stands but also higher concentrations of exchangeable Al^{3+} . The 'base' cation saturation in the deepest soil layers was highest in the 20-year-old stand.

The storage of total and of available elements in the organic and mineral soil are presented in Table 1.

				•											
Р	K	Ca	Mg	z	s	۲. ۲.	×	Ca	Mg	z	S	Ь	ĸ	Ca	Mg
15	156	100	18	320	46	25	270	189	28	469	46	41	208	391	42
/ear ⁻¹)															
0.4	6.0	1.7	0.5	9.2	1.3	0.7	6.9	5.6	0.8	6.2	-0.4	0.5	-3.0	5.8	0.4
1.6	11.9	13.1	1.8	11.1	1.5	0.9	7.1	5.3	0.9	9.3	1.6	1.1	- 1.7	14.0	0.8
4	5	42	5	149	шu	×	21	48	8	88	шu	9	10	53	4
23	120	105	75	524	шu	35	195	06	90	500	шu	36	413	145	138
622	185	230	49	ши	ШU	344	180	127	26	шu	m	280	360	256	53
1384	298	617	143	шш	mm	521	308	187	41	nm	шu	378	552	360	82
649	310	377	129	шш	nm	387	396	265	124	шu	шu	322	783	454	195
1411	423	764	223	шш	шш	564	524	325	139	шu	шu	420	975	558	224
0.9	78.3	5.8	22.0	5.3	шu	0.0	123.5	3.3	20.5	6.8	шu	1.3	153.8	3.1	24.0
1.6	158.5	12.5	45.0	6.3	шu	1.7	277.7	7.4	52.8	7.9	шu	2.3	359.4	10.0	66.5
Example 1	15 year ⁻¹) 5 0.4 4 23 622 1384 649 1411 1.6	year ⁻¹) 5 0.4 6.0 3 1.6 11.9 23 120 622 185 1384 298 649 310 1411 423 0.9 78.3 1.6 158.5	$\begin{array}{cccccccccccccccccccccccccccccccccccc$	$\begin{array}{c ccccccccccccccccccccccccccccccccccc$	$\begin{array}{c ccccccccccccccccccccccccccccccccccc$	$ \begin{array}{c ccccccccccccccccccccccccccccccccccc$	$ \begin{array}{c ccccccccccccccccccccccccccccccccccc$								

Nutrient storage and immobilisation in the compartments of the forest stands and soils

Table 1

^b N expressed as total organic N; P extracted following Duchaufour and Bonneau (1959); Ca and Mg extracted by 1 N KCl and K by 1 N NH₄Cl; L, litter; F+H, decomposed leaves and humus.

° N expressed as total organic N: P, K, Ca and Mg extracted by HF dissolution.

nm, not measured.

169

2.3. Measurement of atmospheric input and seepage water

Nutrient fluxes were measured in four compartments of the ecosystem following the classical model for forest plot studies. Compartments were: (i) open area, (ii) canopy, (iii) forest floor and (iv) soil (subdivided into four layers in the 40- and 60-year-old stands, and into three layers in the 20-year-old stand).

Daily bulk precipitation in the open area (Pt) was collected by an automatic rain gauge. Flux through the canopy was considered to be total throughfall plus stemflow (TS), which were sampled, respectively, by double gutters under the canopy (three replicates per plot) and collars around tree trunks (ten replicates per plot). Stemflow was locally large and often overflowed containers (60-120 1), but it only represented 3% of TS when the total plot area was considered. Forest floor solutions (FF) were collected by three sets of nine narrow lysimeters. Gravitational soil solutions from 15 cm (P-15), 30 cm (P-30), 60 cm (P-60) and 120 cm depth (P-120) were sampled by zero-tension plate lysimeters (four replicates per level) in the 40- and 60-year-old stands and only from the top three levels in the 20-year-old stand.

Collection and measurements started in July 1992 and ended in June 1995.

2.4. Solution analyses

After volume measurement and sampling, the solutions were carried to the laboratory, filtered (0.45 μ m), maintained at 4°C and analysed as quickly as possible. Measurements of pH were made with a single-rod pH electrode (XEROLIT[®]) connected to a Mettler DL21 pH meter. Nitrate, ammonium and chloride were measured by colorimetry (Technicon autoanalyser II) and total Si, S, P, K, Ca, Mg, Mn, Na, Fe and Al by ICP emission spectroscopy (JY 38 Plus spectrometer). Total organic carbon (TOC) was measured during the first year on an TCM 480 (Erba Science) apparatus and subsequently on a TOC 5050 (Shimadzu).

2.5. Water fluxes

Fluxes in the first two compartments, *Pt* and *TS*, were directly available from field measurements.

Fluxes through the forest floor and within the soil, which were measured by lysimeters, were inadequate for estimating soil water seepage because it was not possible to determine the volume of soil actually drained by the lysimeters.

A water flux model (Villette, 1994) was then used to quantify fluxes through the ecosystem. This model was first developed by Breda (1994). Daily rainfall was available from field measurements; model fluxes were estimated daily and cumulated over a 4 week period to coincide with the water chemistry data. Assumptions for model calculations were mainly based on earlier investigations by Aussenac and Boulangeat (1980) and Granier (1987). Parametrisation was adapted to the present study according to the structure of stands and to the local climate conditions.

The general equation of the model is

$$Pt = In + TR + ET_u + \Delta S_{soil} + DR$$

where Pt is incident precipitation, In is canopy interception, TR is transpiration of trees, ET_u is evapotranspiration of the understorey, ΔS_{soil} is soilwater storage variation and DR is drainage at 60 or 120 cm depth. Pt was mainly obtained from local measurements but occasionally had to be corrected, using its relationship with throughfall when monthly rainfall was lower than monthly throughfall (this happened during some snow events).

Interception was calculated by the equation

$$In = Pt - TS$$

TS in the model was estimated by computing throughfall (THR) and stemflow (STF) separately with the following equations calibrated for the stands

$$\frac{\text{THR}}{Pt} = \frac{0.87 \times (Pt-1)^{1.7}}{2.8 + (Pt-1)^{1.7}}$$
$$\frac{\text{STF}}{Pt} = \frac{0.025 \times (Pt-1)^3}{2 + (Pt-1)^3}$$

TR was not measured directly but estimated by the relationship with potential evapotranspiration (PET) calculated from the Penman equation

$$TR = 0.65 \times PET$$

The coefficient 0.65 was derived from sap-flow measurements and a water balance obtained on Dou-

glas-fir stands in the Lorraine region by Aussenac and Boulangeat (1980) and Granier (1987); PET was provided by the Meteorological Station of 'Tarare-Les Sauvages' situated 50 km further south at the same elevation. Granier (1987) observed that this coefficient varied with the relative extractable water (REW) in the soil. REW is defined as

$$REW = \frac{\text{extractable soil water}}{\text{maximum available soil water}}$$

Maximum available soil water (Ws) was defined as

$$W_{s} = (M_{c} - M_{pF4,2}) \times D_{a} \times (1 - S_{p}) \times \rho$$

where M_c is soil moisture at field capacity, $M_{pF4,2}$

is soil moisture at pF = 4.2, D_a is apparent density of fine earth, S_p is volumetric proportion of stones and ρ is the density of water (1 kg l⁻¹).

Under soil water limiting conditions (water stress), TR was assumed to decrease linearly with REW, according to observations made in four Douglas-fir stands (Aussenac and Boulangeat, 1980)

 $TR = 0.65 \times PET \times REW/0.4$

The limitation of TR when foliage is wet was also taken into account in the model.

 ET_{u} was estimated for each stand from the estimated global radiation decrease between open area



Fig. 1. Schematic water fluxes after the model developed by Villette (1994).

 (Rg_0) and under tree cover (Rg_u) according to the following formula

$$ET_{\rm u} = {\rm PET} \times \frac{Rg_{\rm u}}{Rg_{\rm o}} \times {\rm REW}_{30}$$

 Rg_{u} and Rg_{o} were measured with a lux meter;

 REW_{30} is the relative extractable water in the upper 30 cm soil layer.

Actual evapotranspiration of stands (*ET*), which did not appear in the general equation, was calculated as: $ET = TR + ET_u + In$.

 $\Delta S_{\rm soil}$ was the variation of the available water



Fig. 2. (a) Relationship between measured water (1) from lysimeters and estimated water fluxes (mm) in the soil. (b) Relationship between measured water from P-30 lysimeters and estimated water fluxes in the soil at 30 cm depth. As sampling was done every 4 weeks, there are some months where sampling occurred twice.



content in each soil layer during the water flux, i.e. final water content minus initial water content.

Water fluxes were calculated by considering each compartment as a black-box and fluxes resulted from the balance between gain and loss in each box (Fig. 1).

Input-output calculations in the overground compartments (Pt and TS) assumed that input equalled Pt and output was equal to $In + ET + ET_u$.

Forest floor water flux was considered as $TS - ET_{w}$.

Mineral soil was divided into four layers corresponding to the lysimeter depths for characterising individual budgets: 0-15, 15-30, 30-60 and 60-120cm. Input to the 0-15 cm soil layer was the drainage flux from forest floor and so on for the next soil layer down to a depth of 60 or 120 cm. Water penetration into each soil horizon was portioned out into a quick flux along the macroporosity and a slow diffusion flux along the microporosity. Water losses by evapotranspiration were assumed to be proportional to the distribution of roots in each soil layer which was evaluated from grid measurements on two soil pits per stand. If the budget was positive then drainage (DR) occurred, if negative then the soil reservoir was being emptied by the plants.

Significant correlation between measured and calculated values of Pt and TS (n = 37; $r_{20} = 0.95$, $r_{40} = 0.97$, $r_{60} = 0.97$) and very close absolute values were observed. Estimated soil water fluxes showed a time pattern variation rather well correlated with water volume obtained from the lysimeters (Fig. 2(a)Fig. 2(b)) but direct comparison of absolute values was not possible. Nevertheless, any errors in the estimations would not affect comparisons between stands since they would apply to all stands equally.

2.6. Nutrient fluxes

Element fluxes were obtained by weighting concentrations with the calculated water fluxes from the model. Seasonal and annual fluxes were the total amount of fluxes at the end of each season or year.

Seasons were defined as follows: summer, July to September; autumn, October to December; winter, January to April; spring, May to late June.

2.7. Partitioning of atmospheric input

Total atmospheric deposition (TD) to forest ecosystems are separated into wet deposition (Pt), dry deposition (DD), and cloud and fog droplet deposition (Ulrich, 1983; Lovett and Lindberg, 1984). During the 3 years of measurement cloud and fog were considered as negligible; Pt solutions, which were sampled as bulk precipitation, were considered as wet deposition in the calculations.

$$TD_{y} = Pt_{y} + DD_{y}$$
 $x =$ chemical element (1)

Canopy interaction processes (Ci), i.e. canopy leaching and canopy uptake, alter the solution chemistry down through the canopy and are implicit in throughfall and stemflow measurements.

$$TS_x = Pt_x + DD_x \pm Ci_x \tag{2}$$

Lack of reliable measurement methods for DDand Ci fluxes made it necessary to use the calculation approach first described by Ulrich (1983) and adapted here. It was assumed that Ci was negligible or did not exist for certain elements (e.g. Na and S) which therefore could be used as tracers for calculating DD for other elements. Unlike Ulrich (1983) and Bredemeier (1988), we did not separate particulate and gas deposition and S-SO₄²⁻ was used as a tracer in addition to Na. This option was supported by the behaviour of elements in Pt solutions. Na⁺ was better correlated with H₂PO₄⁻-P, K⁺, Ca²⁺ and Mg²⁺, and SO₄²⁻-S with NH₄⁺-N and NO₃⁻-N. Thus, Na⁺ was used as a tracer to calculate DD for $H_2PO_4^--P$, K^+ , Ca^{2+} and Mg^{2+} , and $SO_4^{2-}-S$ was the element tracer for NH_4^+-N and $NO_3^{-}-N$. Dry deposition of Na^+ and $SO_4^{2-}-S$ was considered as TS - Pt. DD for the other elements was calculated with the following equations

$$DD_{x} = \left(\frac{TS - Pt}{Pt}\right)_{Na} \times Pt_{x}$$

$$x = H_{2}PO_{4}^{-} \cdot P, K^{+}, Ca^{2+} \text{ and } Mg^{2-} \qquad (3)$$

$$DD_{y} = \left(\frac{TS - Pt}{Pt}\right)_{SO_{4}} \times Pt_{y}$$

$$y = NH_{4}^{+} \cdot N \text{ and } NO_{3}^{-} \cdot N$$

Once all the *DD* values were obtained, canopy interactions were calculated using Eq. (2). A positive *Ci* value was considered as canopy leaching and a negative value as direct uptake from the canopy. This indicated the net result of the dominant processes occurring in the canopy. Thus, the rate of uptake (C_{up}) from total deposition and the rate of leaching (C_{lc}) by *TS* solutions were calculated as follows

$$C_{\rm up}(\%) = -\left(\frac{-Ci}{TD}\right) \times 100 \text{ and}$$
$$C_{\rm lc}(\%) = \left(\frac{Ci}{TS}\right) \times 100 \tag{4}$$

2.8. Nutrient budgets

Nutrient budgets were established for the major cations and anions; two soil depths were successively considered as the output reference level: 60 cm depth for all stands and 120 cm depth for the 40and 60-year-old stands. The terms of these budgets were as follows

Input = Pt + DD + (weathering)

Because weathering could not be estimated reliably, it was not taken into account in the present input calculation.

Output = Biomass immobilisation + leaching out of

the soil (P-60 or P-120)

Biomass immobilisation (roots not included) represented the mean current immobilisation (MCI)



Fig. 3. Annual and seasonal variations of water input-output fluxes in the 40-year-old stand. As sampling was done every 4 weeks, there are some months where sampling occurred twice.

Table 2											
Annual	water	fluxes	(mm)	estimated	by	the	water	balance	model	(Villette,	1994)

	Pt	Т	S	In	PET	ET	TR	ETu	Drainag	ge ^a		
									P-15	P-30	P-60	P-120
20-year-old star	ıd											· · · · ·
1992/1993	821	604	19	198	569	542	331	12	529	455	350	290
1993/1994	1060	778	24	257	601	614	344	13	702	621	511	449
1994/1995	1193	923	28	242	671	633	377	13	850	769	654	581
Year average	1025	768	24	233	614	596	351	13	694	615	505	440
40-year-old star	nd											
1992/1993	821	604	19	198	569	567	331	38	459	382	308	265
1993/1994	1060	778	24	257	601	631	336	37	635	550	476	435
1994/1995	1193	923	28	242	671	618	339	37	798	718	635	595
Year average	1025	768	24	233	614	605	335	37	631	550	473	432
60-year-old star	nd											
1992/1993	821	604	19	198	569	591	331	62	453	360	265	242
1993/1994	1060	778	24	257	601	654	337	60	628	528	435	413
1994/1995	1193	923	28	242	671	641	340	58	787	694	598	572
Year average	1025	768	24	233	614	629	336	60	622	527	433	409
Annual average	for all sta	nds										
Ť	1025	768	24	233	614	610	341	37	649	564	470	427

^a P-15, P-30, P-60 and P-120 represent drainage from 15, 30, 60 and 120 cm depth, respectively.

Pt, bulk precipitation; T, throughfall; S, stemflow; In, interception; PET, Penman evapotranspiration; ET, estimated evapotranspiration; TR, transpiration; ET_u , evapotranspiration of understorey.



Fig. 4. (a) Mean seasonal water (mm) and NO_3-N , SO_4-S and TOC fluxes (kg ha⁻¹ per season) in the compartments of the forest stands. (b) Mean seasonal Ca, Mg, Al and K fluxes (kg ha⁻¹ per season) in the compartments of the forest stands. (c) Mean seasonal Si, Mn, Na and Cl fluxes (kg ha⁻¹ per season) in the compartments of the forest stands. (d) Mean seasonal P, NH₄-N and H fluxes (kg ha⁻¹ per season) in the compartments of the forest stands.



Fig. 4 (continued).





Fig. 4 (continued).

which was calculated from the total nutrient content (Tnc) estimated for each stand (Ranger et al., 1995)

$$MCI (20-year-old stand) = \frac{(Tnc 20)}{20}$$
$$MCI (40-year-old stand) = \frac{(Tnc 40 - Tnc 20)}{20}$$
$$MCI (60-year-old stand) = \frac{(Tnc 60 - Tnc 40)}{20}$$

For the 40- and 60-year-old stands, the nutrient amount exported by thinning was included in the calculation of MCI. This amount was evaluated by applying the mean nutrient concentrations observed in the samples to the biomass exported by thinnings (data given by the Forest Management Service).

Budgets for rotations lasting of 20, 40 or 60 years were calculated as follows:

Reference level at 60 cm depth

20-year-old stand: average annual budget (20-year-old stand) \times 20 (A)

40-year-old stand: average annual budget (40-year-old stand) \times 20 + A (B)

60-year-old stand: average annual budget (60-year-old stand) \times 20 + B

Reference level at 120 cm depth

40-year-old stand: average annual budget (40-year-old stand) \times 40 (C)

60-year-old stand: average annual budget (60-year-old stand) \times 20 + C

3. Results

3.1. Water flux

Precipitation averaged 1025 mm per year. The first year sampled (1992/1993) was drier than the others with a total rainfall of 821 mm and a particularly dry winter period. Mean seasonal rainfall was lowest in summer, when the water flux at the bottom of the soil was very low (Fig. 3).

Canopy interception, throughfall and stemflow amounted to 23%, 75% and 2% of mean incident rainfall, respectively. The mean annual loss of water by *ET* and *TR* was 610 mm and 341 mm, respectively, i.e. 60% and 33% of the incident precipitation (Table 2).

Water fluxes decreased from Pt to P-120 and this decrease was more accentuated during spring and summer (Fig. 4(a)).

Net annual water loss at the soil bottom averaged 427 mm but a strong variability (range = 250-600 mm) related to rainfall input was observed. Stand variability was limited, less than 5% of the general mean.

3.2. Nutrient flux

Nutrient fluxes showed the following general trend: (i) they were low in the first two compartments (Pt and TS), (ii) increased in the upper soil layers (from FF to P-30) and then (iii) decreased with soil depth (Fig. 4(a)Fig. 4(b)Fig. 4(c)Fig. 4(d)).

Mean annual fluxes of the three stands are presented in Table 3. Yearly rainfall input to the ecosystem per hectare was 27.4 kg TOC, 5.9 kg Cl⁻, 5.5 kg SO₄²⁻-S, 4.8 kg NO₃⁻-N, 3.6 kg NH₄⁺-N, 3.1 kg Ca²⁺, 3.0 kg Na⁺, 2.0 kg K⁺, 1.0 kg H₂PO₄⁻-P and less than 1.0 kg for Si, Mg²⁺, Mn²⁺, H⁺, Al³⁺. and Fe²⁺.

TS solutions were enriched in some elements and impoverished in others as a consequence of canopy interactions. The partitioning of atmospheric input was used to estimate dry deposition, canopy leaching and direct uptake of nutrients by the canopy (Table 4). Mean total deposition (TD) was estimated at 12.5 kg S, 10.8 kg NH₄⁺-N, 8.2 kg NO₃⁻-N, 5.7 kg Ca²⁺. 5.3 kg Na⁺, 3.5 kg K⁺, 1.6 kg $H_2PO_4^-P$ and 1.0 kg Mg^{2+} ha⁻¹ year⁻¹. Some 58% of H₂PO₄⁻-P and 48% of NH⁺₄-N were in average taken up from the total deposition by the canopy. K-, Ca- and Mgleaching were respectively estimated at 78%, 36% and 52% of TS. Thus, fluxes reaching the soil surface (TS) were on average 103.0 kg TOC, 18.6 kg K^+ , 14.6 kg Cl⁻, 12.5 kg SO₄²⁻-S, 9.1 kg Ca²⁺, 8.6 kg NO₃⁻-N, 5.7 kg NH₄⁺-N, 5.3 kg Na⁺. 2.1 kg Mg^{2+} and less than 2.0 kg for Mn^{2+} , Si, $H_2PO_4^--P$. Fe^{2+} , Al^{3+} and H^+ ha⁻¹ year⁻¹ (Table 3).

The downward fluxes (FF) of all elements significantly increased, the highest nutrient fluxes were those of K⁺. Ca²⁺, and NO₃⁻-N, which averaged, respectively, 51.8, 45.0 and 43.6 kg ha⁻¹ year⁻¹.

Most of the fluxes then decreased with depth, but large amounts of nutrients flowed out of the deepest soil layer where mean fluxes amounted to 30.7 kg SO_4^{2-} -S, 26.6 kg NO_3^- -N, 14.8 kg K $^+$ and 14.5 kg Ca^{2+} ha^{-1} year $^{-1}$ (Table 3).

3.3. Seasonal and stand age variations

Seasonal patterns according to stand age are presented for some elements on Fig. 4(a)Fig. 4(b)Fig. 4(c)Fig. 4(d). Most elements reached their highest values in autumn and winter. During these seasons, the decreasing trend with depth did not occur; several elements showed higher fluxes at P-120 than in the upper layers. Elements like NO₃⁻-N, K⁺, Ca²⁺ and Mn^{2+} also showed large fluxes in summer. TOC, at the *FF* level, was elevated throughout the year.

Fluxes of most elements were highest in the 20-year-old stand, irrespective of season and compartment except for NO_3^--N , Ca^{2+} , Na^+ and Cl^- in *Pt* and *TS*.

Comparing fluxes was more complicated in the other two stands. In spring and summer, fluxes at all levels were low and generally less variable between stands than during the other seasons. In autumn and winter, Pt and TS did not differ between the 40- and

Table 3

Mean annual fluxes in the stands and average fluxes for all stands (values are in kg ha⁻¹ year⁻¹, except for water flux which is in mm) $\frac{1}{2}$

Compartment	water nux	INH ₄ -IN	NO ₃ -N	SU ₄ -S	H_2PO_4-P	ĸ	Ca	Mg	na	Mn	н	AI	re	51	CI	TOC
20-year-old sta	and															
Pt	1025	4.8	3.6	5.5	1.0	2.0	3.1	0.6	3.0	0.2	0.1	0.1	0.4	0.6	5.9	27.4
TS	792	3.8	4.0	11.2	0.8	27.4	6.5	2.0	3.7	1.9	0.0	0.2	0.5	1.1	12.4	98.7
FF	741	13.9	66.0	32.4	4.2	85.6	67.1	12.1	9.1	6.3	0.1	5.2	1.4	19.6	33.0	293.5
P-15	686	2.7	85.8	30.0	0.7	31.1	40.7	11.5	6.7	15.0	0.4	32.7	1.5	29.6	17.7	160.8
P-30	559	3.5	90.3	39.7	0.6	42.2	56.9	14.5	7.6	12.3	0.3	28.3	0.6	25.8	20.9	107.5
P-60	395	2.3	36.6	15.5	0.3	22.7	21.7	9.1	9.1	3.2	0.1	10.1	0.4	14.0	22.2	87.8
40-year-old sta	and															
Pt	1025	4.8	3.6	5.5	1.0	2.0	3.1	0.6	3.0	0.2	0.1	0.1	0.4	0.6	5.9	27.4
TS	792	6.0	10.1	13.5	0.6	15.0	10.7	2.2	5.3	2.1	0.2	0.4	0.6	0.6	14.6	106.0
FF	730	12.0	33.3	18.8	1.6	31.0	35.8	5.9	7.4	11.1	0.2	3.6	2.1	12.6	27.0	290.7
P-15	586	4.1	38.7	23.5	0.4	14.0	30.9	7.0	6.7	7.6	0.3	17.5	1.0	22.0	17.7	163.0
P-30	547	2.0	27.6	23.9	0.6	10.6	20.4	6.4	5.6	6.7	0.2	14.6	0.6	21.5	14.1	85.0
P-60	462	0.3	24.9	23.1	0.7	7.7	10.2	5.1	4.7	4.1	0.2	15.4	0.3	23.3	8.5	22.5
P-120	399	0.6	21.1	45.6	0.4	20.6	13.6	7.0	12.7	5.0	0.2	21.2	0.3	19.5	28.2	35.8
60-year-old sta	and															
Pt	1025	4.8	3.6	5.5	1.0	2.0	3.1	0.6	3.0	0.2	0.1	0.1	0.4	0.6	5.9	27.4
TS	792	7.3	11.8	12.7	0.7	13.4	10.1	2.2	6.8	1.5	0.2	0.5	0.6	0.6	16.8	104.2
FF	709	11.2	31.5	13.7	1.8	38.8	32.2	6.1	5.8	7.2	0.2	3.3	1.8	9.0	26.7	294.0
P-15	620	1.5	70.8	16.9	0.6	23.0	47.4	8.4	5.9	7.0	0.3	21.1	0.6	30.4	15.5	103.4
P-30	511	0.9	36.0	12.7	0.5	10.5	19.5	4.8	4.2	5.0	0.2	12.0	0.4	19.9	10.4	45.2
P-60	324	0.4	5.9	14.7	0.3	4.9	5.1	1.8	2.7	1.7	0.1	5.4	0.2	7.8	6.0	17.7
P-120	312	0.3	32.2	15.7	0.3	9.0	15.3	4.6	4.3	6.2	0.1	13.1	0.2	12.8	8.2	14.1
Average for al	ll stands															
Pt	1025	4.8	3.6	5.5	1.0	2.0	3.1	0.6	3.0	0.2	0.1	0.1	0.4	0.6	5.9	27.4
TS	792	5.7	8.6	12.5	0.7	18.6	9.1	2.1	5.3	1.8	0.1	0.4	0.6	0.8	14.6	103.0
FF	727	12.4	43.6	21.6	2.6	51.8	45.0	8.0	7.4	8.2	0.2	4.0	1.8	13.7	28.9	292.7
P-15	631	2.8	65.1	23.5	0.6	22.7	39.7	9.0	6.4	9.9	0.3	23.7	1.1	27.3	17.0	142.4
P-30	539	2.1	51.3	25.4	0.6	21.1	32.2	8.6	5.8	8.0	0.2	18.3	0.5	22.4	15.1	79.2
P-60	394	1.0	22.5	17.8	0.4	11.8	12.4	5.4	5.5	3.0	0.1	10.3	0.3	15.0	12.2	42.7
P-120 a	356	0.5	26.6	30.7	0.3	14.8	14.5	5.8	8.5	5.6	0.1	17.2	0.2	16.2	18.2	24.9

^a Average for the 40- and 60-year-old stands.

Pt. bulk precipitation; TS, throughflow + stemflow; FF, forest floor; P-15, P-30, P-60 and P-120, lysimeters at 15, 30, 60 and 120 cm depth.

NH4-N 1 992/1993 59 4.9 50 3.7 10 5.0 10 5.0 10 9.9 72 TD -6.1 72 TD -6.1	NO _X -N	2														y-irad-/w							
902/1993 ²¹ 4.9 13 3.7 10 5.0 10 5.0 10 - 6.1 * D * Clc			r L	×	Ca	Мg	R Z	N++HN	NO3-N	s	4	×	Ca	Mg 1	- P	N-+ HN	NO ₃ -N	s	۵.	ы	Ca	Mg	R'N
7 4.9 55 3.7 9D 5.0 1D 9.9 1D -6.1 KS TD -6.1 KGIC																							
⁵ 3.7 2D 5.0 7D 9.9 FS − TD − 6.1 & D &Clc	3.0	4.7	0.3	1.6	3.1	0.5	6.1	4.9	3.0	4.7	0.3	1.6	3.1	0.5	6.1	4.9	3.0	4.7	0.3	1.0	3.1	0.5	6.1
0D 5.0 1D 9.9 1S - TD - 6.1 #D #Clc	3.8	9.6	0.3	26.8	5.5	1.7	3.1	5.6	8.3	13.3	0.2	13.1	10.5	1.9		7.2	11.0	12.6	0.2	511.3	10.3	2.0	5.5
"D 9.9 rS - TD - 6.1 &D %Clc	3.0	4.8	0.2	0.9	I .8	0.3	1.1	8.8	5.3	8.5	0.3	1.7	3,4	0.5	5	8.1	6.4	7.9	0.6	2.9	5.7	0.9	3.6
rs - TD - 6.1 & D %Clc	6.0	9.6	0.5	2.5	4.9	0.8	3.1	13.7	8.3	13.3	0.7	3.3	6.5	1.0	₫`[13.0	9.7	12.6	0.9	4,4	8.7	Ŧ.	5.5
<i>й D</i> 7.Clc	- 2.2	0.0	-0.2	24.3	0.6	0.9	0.0	- 8.1	0.0	0.0	- 0.5	9.9	4.0	0.8	0.0	- 5.8	3.1	0.0	· 0.7	6.8	1.6	0.5	0.0
∦Clc		001		5	88	46	100		100	100		25	62	56	00		72	001		65	85	72	100
		c		16	2	54	0		0	0		75	38	1	c		28	0		61	15	28	0
7 Cup 62	36		46					50			72					4			5				
1993 / 1994																							
Dr 4.4	4.2	6.0	8.1	3.6	3.7	0.8	3.9	t.t		6.0	<u>,</u>	3.6	3.7	0.8	3.9	1 .4	<u>1</u>	6.0	8.1	3.6	3.7	0.8	3.9
rS 2.1	3.1	6.11	<u> </u>	26.2	6.3	5.1	4.3		8.6	13.9	1.0	15.9	10.8	2.3	6.4	5.8	6.01	12.7	1.0	12.9	9.4	2.2	7.5
DD 4.4		5.9	0.2	0.3	0.3	0.1	0.3	5.9	5.5	7.9	1.1	5.3	5.4	0.5	2.5	5.0	4.6	6.7	1.6	3.3	3.4	0.7	3.6
r <i>D</i> 8.8	8.3	11.9	1.9	3.9	4.0	0.9	4.3	10.3	9.7	13.9	6.5	5.9	6.1	1.3	6.4	t'6	8.8	12.7	4.5	6.9	1.1	1.5	7.5
FS = TD = -6.7	- 5.2	0.0	- 0.6	22.3	2.2	1.2	0.0	- 6.2		0.0	- 1.9	10.0	4.7	0.9	0.0	- 3.6	-	0.0	+ 1 .4	6.1	2.3	0.6	0.0
7 D		001		15	65	43	100			100		37	57	1 65	00		81	100		53	76	72	100
%Cle		Û		85	35	57	С			c		63	43 .	=	0		19	0		47	÷-	28	0
%Сир 76	63		32					60	=		65					30			70				
1994/1995																							
Pr 4.9	3.8	5.8	0.8	0.8	5.6	b .0	3.2	4.9	3.8	5.8	0.8	0.8	2.6	0.4	2.5	6.t	8.8	5.8	0.8	0.8	2.6	t'0	с і ~,
TS 5.7	5.2	12.0	0.7	1.62	7.6	 	3.9	8.4	13.3	13.3	0.5	15.9	10.7	2.3	5.5	0.0	13.4	12.9	0.8	16.0	10.8	4.2	7.3
DD 5.2	4.0	6.2	<u>5.0</u>	0.2	0.5	0.1	0.6	6.3	6't	7.5	0.6	0.6	6.I	0.3	2	6.0	4.6	7.0	1.1		t.5	5.0	1,1 1
TD 10.2	7.8	12.0	1.0	1.0	3.1	0.5	3.9	11.3	8.6	13.3	۲. I	+.	5.4	0.7	5.5	10.9	8.4	12.9	1.9	6.T	0 U	6.0	7.3
\mathcal{L}^{+} = \mathcal{L} = \mathcal{L} = \mathcal{L}	- 2.6	0.0	£ () -	28.2	4.5	8.1	0.0	- 2.9	4.7	0.0	- 1.0	14.5	6.2	1.7	0.0	6.1 -	5.1	0.0	- 1.1	14.2	×.	2	0.0
64 D		001		r.	41	5	100		65	100		6	4	29	8		62	100		21	56	37	100
容Clc 容		c		<i>1</i> 6	59	67	0		35	0		16	58	71	0		38	c		88	4	63	c
%Cup +1	33		38					56			67					×			27				
Annual average																							
Pr 4.8	3.6	5.5	1.0	2.0	3.1	0.6	3.0	4.8	3.6	5.5	1.0	0.5	1.1	06	3.0	4.8	3.6	5.5	0.1	0.2	3.1	9.6	3.0
TS 3.8	1.0	71	0.8	27.4	6.5	5.0	3.7	6.0	10.1	13.5	0.6	15.0	10.7	ri ri	5.3	7.3	11.5	12.7	0.7	13.4	10.1		6.8
DD 4.9	3.7	5.6	0.2	0.5	6.9	0.1	0.7	7.0	5.2	8.0	0.7	5.1	2.6	6.4	~	63	17	1.2	1.1	4		0.7	17
TD 9.6	7.3	7.1	1.1	2.5	4.0	0.7	3.7	11.8	8.9	13.5	1.7	3.5	5.7	0.1	5.3	11.1	8.3	12.7		4.4	73	۲. ۲	6.8
TS - TD - 5.8	-3.3	0.0	- 0.4	24.9	2.4	<u>~</u>	0.0	- 5.7	-	0.0		1.5	5.0	1.1	0.0	3.8	3,4	0.0	1,4	0.0	0.0 C	0.9	0.0
G D		001		6	65	15	100		55	001		24	54	48	00		72	001		35	72	60	00
%Clc		с		16	35	63	¢		<u>1</u>	c		76	46	52	0		8	0		65	28	40	0
%Cup 61	44		35					Xt	4		20					7			69				

vear 1- nd nd ÷ . f de Table 4 Partitioni

82

R. Marques et al. / Forest Ecology and Management 92 (1997) 167-197

	20-year-	old stand							40-year-c	old stand							60-year-	old stand						
	N-*HN	NO ₃ -N	s	Р	×	Ca	Mg	Ra	NH4-N	NO3-N	s	4	×	Ca	Mg	Na	N-+HN	N-£ON	s	4	×	Ca	Mg	Ra
Dormant 5	season																							
Pı	8.1	1.5	2.0	0.5	0.7	1.0	0.2	1.9	8.1	1.5	2.0	0.5	0.7	1.0	0.2	1.9	1.8	1.5	2.0	0.5	0.7	1.0	0.2	6.1
TS	1.7	2.6	5.9	0.5	11.0	3.5	1.1	2.2	2.6	5.2	7.0	0.3	6.1	5.2	1.1	3.1	3.3	6.3	6.5	0.3	5.6	5.3	1.2	4.4
aa	3.4	3.0	3.9	0.1	0.1	0.1	0.0	0.3	4.3	3.8	5.0	0.3	0.5	0.6	0.1	1.2	3.9	3.4	4.4	0.6	0.9	1.3	0.3	2.5
TD	5.2	4.5	5.9	0.5	0.8	1.2	0.3	2.2	6.1	5.3	7.0	0.7	1.2	1.7	0.4	3.1	5.6	4.9	6.5	1.0	1.7	2.3	0.5	4.4
TS - TD	- 3.5	- 1.9	0.0	- 0.1	10.2	2.4	0.9	0.0	- 3.5	- 0.1	0.0	- 0.5	4.9	3.5	0.7	0.0	- 2.3	1.4	0.0	- 0.7	3.9	2.9	0.7	0.0
\mathcal{C}^{M}			100		90	33	23	001			100		20	32	33	100		78	100		30	44	43	100
%Clc			0		92	67	11	0			0		80	68	67	0		22	0		70	56	57	0
%Cup	67	43		14					58	-		60					40			69				
Olowing :	SC45UII								6				-		4		4						0	
Ρ1	9.0	77	C.5	0.5	<u>.</u>	7.1	0.5		5.0	7.1	C.F	C.U	Ĵ.	7.1	0.5	-	5.0	7.1	3	0.0	Į.	7.1	0.5	-
TS	2.2	1.5	5.2	0.3	16.4	3.0	0.9	1.5	3.5	4.8	6.5	0.3	8.9	5.4		2.2	4.0	5.5	6.3	0.3	7.8	4.9	1.0	2.4
DD	1.5	1.0	1.7	0.2	0.5	0.8	0.1	0.4	2.6	8.1	3.0	0.5	ñ	2.2	0.4	1.1	2.4	1.7	2.8	0.6	1.5	2.5	0.4	1.3
TD	4.5	3.1	5.2	0.7	8. I	3.0	0.5	1.5	5.6	3.9	6.5	1.0	2.5	4.3	0.7	2.2	5.4	3.8	6.3		2.8	4.6	0.8	4. 4
TS - TD	- 2.3	- 1.7	0.0	- 0.4	14.6	0.0	0.4	0.0	- 2.1	0.9	0.0	- 0.8	6.4	1.2	0.4	0.0	- 1.4	1.7	0.0	- 0.8	5.1	0.3	0.3	0.0
$\mathcal{U}^{W}D$			100		=	66	54	001		81	001		29	78	65	001		68	001		35	65	74	001
%Clc			С		89		46	0		61	0		71	22	35	0		32	0		65	5	26	0
%Cup	52	53		56					38			75					26			70				

Table 6 Annual and ann	aal averag	e nutrie:	nt budį	gets of the	Douglas	-fir stands	s, and mea	an nutrien	t budgets	s of the s	tand chro	nosequei	nce (data e	xpressec	l in kg	ha ⁻¹ y	car ⁻¹)	
	20-year-c	d stand					40-year-o	old stand					60-year-o	ld stand				
	Z	s	Р	K	Ca	Mg	z	S	Р	K	Ca	Mg	z	S	Р	X	Ca	Mg
1992 / 1993																		
Pt	7.9	4.7	0.3	1.6	3.1	0.5	7.9	4.7	0.3	1.6	3.1	0.5	7.9	4.7	0.3	1.6	3.1	0.5
DD	8.0	4.8	0.2	0.9	1.8	0.3	14.1	8.5	0.3	1.7	3.4	0.5	13.0	7.9	0.6	2.9	5.7	0.9
Total input	15.9	9.6	0.5	2.5	4.9	0.8	22.0	13.3	0.7	3.3	6.5	1.0	20.9	12.6	0.9	4.4	8.7	1.4
Immob. ^a	4.7	0.5	0.4	6.0	1.7	0.5	9.2	1.0	0.7	6.9	5.6	0.8	6.2	- 0.4	0.5	- 3.0	5.8	0.4
Dr. 60	26.6	8.9	0.1	14.4	15.4	5.9	33.5	15.9	0.3	6.7	10.2	6.4	7.5	9.9	0.0	2.7	4.2	1.7
Output 60	31.3	9.4	0.5	20.4	17.1	6.4	42.7	16.9	1.1	13.6	15.8	7.2	13.6	9.5	0.5	-0.3	10.0	2.1
Dr. 120							39.7	25.7	0.1	13.9	14.7	7.2	19.8	7.2	0.0	7.0	7.9	2.8
Output 120							48.9	26.7	0.8	20.8	20.2	8.0	26.0	6.8	0.5	4.0	13.7	3.2
Budget 60 cm	- 15.5	0.2	0.0	- 17.9	- 12.3	5.6	- 20.8	- 3.7	-0.4	- 10.3	- 9.3	- 6.1	7.3	3.1	0.4	4.8	- 1.3	- 0.7
Budget 120 cm							- 27.0	- 13.4	- 0.1	- 17.5	- 13.8	7.0	-5.1	5.8	0.4	0.5	- 5.0	- 1.8
1993 / 1994																		
Pt	8.6	6.0	1.8	3.6	3.7	0.8	8.6	6.0	1.8	3.6	3.7	0.8	8.6	6.0	1.8	3.6	3.7	0.8
aa	8.5	5.9	0.2	0.3	0.3	0.1	11.4	6'L	1.1	2.3	2.4	0.5	9.6	6.7	9.1	3.3	3.4	0.7
Total input	17.1	6.11	2.0	3.9	4.0	0.9	20.0	13.9	2.9	5.9	6.1	1.3	18.2	12.7	3.4	6.9	7.1	1.5
Immob.	4.7	0.5	0.4	6.0	1.7	0.5	9.2	1.0	0.7	6.9	5.6	0.8	6.2	- 0.4	0.5	- 3.0	5.8	0.4
Dr. 60	24.2	11.8	0.5	8.3	12.4	7.1	13.9	26.4	1.2	6.9	8.7	4.1	2.4	11.5	0.5	3.0	3.1	1.2
Output 60	28.9	12.3	0.8	14.2	14.1	7.6	23.1	27.4	1,9	13.7	14.2	4.9	8.6	11.2	1.0	0.0	9.0	1.6
Dr. 120							6.3	49.9	0.7	18.8	10.9	5.9	29.0	12.9	0.6	6.6	14.0	4.7
Output 120							15.6	50.9	4.	25.7	16.5	6.7	35.2	12.5	1.1	3.6	19.8	5.1
Budget 60 cm		- 0,4	-	- 10.3	10.1	- 6.7	- 3.1	- 13.5	0.9	- 7.8	- 8.1	- 3.6	9.6	1.5	53	6.9	- 1,8	-0.1
Budget 120 cm							4.5	- 37.0	1.5	- 19.8	- 10.4	4.6	- 17.0	0.2	сі Сі	3.3	<u>1</u>	- 3.6

R. Marques et al. / Forest Ecology and Management 92 (1997) 167-197

1001 1001																		
001 / 1001 CV21 CV21 CV21 CV21 CV21 CV21 CV21 CV2	8.7	5.8	0.8	0.8	2.6	0.4	8.7	5.8	0.8	0.8	2.6	04	87	5 8	80	0.8	<i>2</i> 6	04
aa	9.2	6.2	0.2	0.2	0.5	0.1	11.2	7.5	0.6	0.6	1.9	0.4	10.6	7.0	1.I	I.I	3.4	0.5
Total input	17.9	12.0	1.0	1.0	3.1	0.5	19.9	13.3	1.4	1.4	4.5	0.8	19.3	12.8	1.9	1.9	6.0	0.9
Immob.	4.7	0.5	0.4	6.0	1.7	0.5	9.2	1.0	0.7	6.9	5.6	0.8	6.2	-0.4	0.5	- 3.0	5.8	0.4
Dr. 60	62.9	25.9	0.3	45.5	37.3	14.4	28.2	26.9	0.5	9.4	11.8	4.9	9.2	22.5	0.3	9.1	8.1	2.7
Output 60	70.6	26.3	0.6	51.4	39.0	15.0	37.5	28.0	1.2	16.3	17.4	5.7	15.4	22.2	0.9	6.1	13.9	3.1
Dr. 120							19.3	61.4	0.5	29.2	15.2	8.0	48.6	27.0	0.3	13.3	24.0	6.2
Output 120							28.5	62.5	1.2	36.1	20.8	8.8	54.8	26.6	0.8	10.3	29.9	6.6
Budget 60 cm	- 52.7	- 14.3	0.4	-50.4	- 35.9	- 14.5	- 17.6	- 14.6	0.2	-14.9	- 12.8	-4.9	4.0	-9.3	I:I	-4.1	- 7.9	-2.2
Budget 120 cm							- 8.6	- 49.1	0.3	- 34.7	- 16.2	-8.0	-35.5	-13.8	I.1	- 8.4	- 23.8	-5.7
Annual average																		
Pr	8.4	5.5	1.0	2.0	3.1	0.6	8.4	5.5	1.0	2.0	3.1	0.6	8.4	5.5	1.0	2.0	3.1	0.6
DD	8.6	5.6	0.2	0.5	0.9	0.2	12.2	8.0	0.7	1.5	2.6	0.5	11.1	7.2	1.1	2.4	4.2	0.7
Total input	17.0	11.2	1.2	2.5	4.0	0.7	20.6	13.5	1.7	3.5	5.7	1.0	19.5	12.7	2.1	4.4	7.3	1.3
Immob.	4.7	0.5	0.4	6.0	1.7	0.5	9.2	1.0	0.7	6.9	5.6	0.8	6.2	-0.4	0.5	-3.0	5.8	0.4
Dr. 60	38.9	15.5	0.3	22.7	21.7	9.1	25.2	23.1	0.7	7.7	10.2	5.1	6.3	14.7	0.3	4.9	5.1	1.9
Output 60	43.6	16.0	0.6	28.7	23.4	9.7	34.5	24.1	1.4	14.5	15.8	5.9	12.5	14.3	0.8	1.9	11.0	2.3
Dr. 120	0.0	0.0	0.0	0.0	0.0	0.0	21.8	45.6	0.4	20.6	13.6	7.0	32.5	15.7	0.3	9.0	15.3	4.6
Output 120	0.0	0.0	0.0	0.0	0.0	0.0	31.0	46.7	1.1	27.5	19.2	7.8	38.6	15.3	0.8	6.0	21.1	5.0
Budget 60 cm	-26.7	- 4.8	0.5	-26.2	- 19.4	- 8.9	- 13.8	-10.6	0.3	-11.0	- 10.1	-4.9	7.0	- 1.6	1.3	2.5	-3.7	- 1.0
Budget 120 cm							- 10.4	-33.2	0.5	- 24.0	- 13.5	-6.8	- 19.2	- 2.6	1.2	- 1.5	- 13.8	-3.7
Year and stand n	nean bud	gets																
Budget 60 cm		נ					- 11.2	-6.1	0.8	-4.2	- 6.9	-2.9						
Budget 120 cm							- 14.8	- 17.9	0.9	- 12.8	- 13.6	-5.3						

^a Immobilisation in the stern only (thinning included).

	20-year-4	old stand					40-year-(old stand					60-year-c	id stand				
	z	s	Ь	×	Ca	Mg	z	S	Р	K	Ca	Mg	N	S	Ч	K	Ca	Mg
Dormant sea.	uos																	
Pt	3.3	2.0	0.5	0.7	1.0	0.2	3.3	2.0	0.5	0.7	1.0	0.2	3.3	2.0	0.5	0.7	1.0	0.2
aa	6.4	3.9	0.1	0.1	0.1	0.0	8.1	5.0	0.3	0.5	0.6	0.1	7.2	4.4	0.6	0.9	<u>.</u> .1	0.3
Total input	9.7	5.9	0.5	0.8	1.2	0.3	11.4	7.0	0.7	1.2	1.7	0.4	10.5	6.5	1.0	1.7	2.3	0.5
Immob.	;	:			t	c t		c u T	, C	ר נ	00		4	-	ç	, ,	0	
Dr. 60	31.7	1 .1	7.0	13.9	10./	5.1	C.22	6.01	4.0	1.0	8.U		0. †	0.11	7.0	C.C	0.0	,
Output 60	31.7	11.4	0.2	13.9	16.7	7.3	22.5	15.9	0.4	5.7	8.0	4.	4.8	11.6	0.2	3.3	3.8	4.1
Dr. 120							19.5	36.7	0.3	15.9	4.11	5.8	26.9	12.1	0.2	7.3	12.2	3.7
Output 120							19.5	36.7	0.3	15.9	4.11	5.8	26.9	12.1	0.2	7.3	12.2	3.7
Budget 60	- 22.1	-5.5	0.3	- 13.1	- 15.5	- 7.0	- 11.1	- 8.9	0.3	-4.5	- 6.3	- 3.8	5.7	-5.1	0.9	- 1.7	- 1.4	-0.9
Budget 120							- 8.1	- 29.6	0.5	- 14.7	- 9.7	- 5.5	- 16.4	-5.6	0.8	- 5.7	- 9.9	- 3.2
Growing sea	uos																	
Pr	5.1	3.5	0.5	1.3	2.1	0.3	5.1	3.5	0.5	1.3	1.1	0.3	5.1	3.5	0.5	I.3	2.1	0.3
aa	2.5	1.7	0.2	0.5	0.8	0.1	4.4	3.0	0.5	1.3	2.2	0.4	4.0	2.8	0.6	1.5	2.5	0.4
Total input	7.6	5.2	0.7	1.8	3.0	0.5	9.5	6.5	1.0	2.5	4.3	0.7	9.1	6.3	I.I	2.8	4.6	0.8
Immob. ^a	4.7	0.5	0.4	6.0	1.7	0.5	9.2	1.0	0.7	6.9	5.6	0.8	6.2	- 0.4	0.5	- 3.0	5.8	0.4
Dr. 60	7.2	4.1	0.0	8.8	5.0	6.1	2.7	7.2	0.3	2.0	2.2	1.0	1.6	3.1	0.1	1.6	4.	0.4
Output 60	0.11	4.6	0.4	14.8	6.7	2.4	11.9	8.2	1.0	8.9	7.8	1.8	7.7	2.7	0.6	- 1.4	7.2	0.8
Dr. 120							2.2	9.0	0.1	4.7	2.2	1.2	5.6	3.6	0.1	1.6	3.1	0.8
Output 120							11.5	10.0	0.8	11.6	7.8	2.0	11.7	3.2	0.6	+	8.9	1.2
Budget 60	-4.3	0.7	0.3	- 13.0	- 3.7	6.1	2.4	- 1.7	0.0	-6.3	- 3.5	- 1.1	1.4	3.5	0.5	4.2	- 2.6	- 0.1
Budget 120							2.0	- 3.5	0.2	- 9.1	- 3.5	- 13	-2.6	3.0	0.5	4.2	- 1.3	0.5
			V-A-H-A-H-RA-HARTON	Non-supervised here is some supervised	the subfraction of the local sector is any series of							of the second	the second s					

	ď,
	-
	ha
	<u>.</u>
	Е.
	expressed
	(data
	stands
	of
	budgets
	nutrient
	put-output
	Ξ.
_	al

Table 7 Seasonal

86

R. Marques et al. / Forest Ecology and Management 92 (1997) 167-197

^a Immobilisation in the stem only (thinning included).

	20-year-c	dd stand					40-year-old	stand					60-year-old	l stand				
	z	s	4	х	Ca	Mg	z	s	Ч	×	Ca	Mg	z	s	Ь	×	Ca	Mg
Stem only harves Annual budget de	tting eficits (kg ha	a-1 vear-1									,							
60 cm depth	- 26.7	- 4.8	0.5	- 26.2	- 19.4	- 8.9	- 13.8	- 10.6	0.3	- 11.0	- 10.1	- 4.9	7.0	- 1.6	1.3	2.5	- 3.7	1.0
120 cm depth							- 10.4	- 33.2	0.5	- 24.0	- 13.5	~ 6.8	- 19.2	- 2.6	1.2	- 1.5	-13.8	- 3.7
Whole rotation b	udget defici-	ts (kg ha ⁻¹	<i>.</i>															
60 cm depth	- 533.2	- 96.7	10.7	- 524.3	-388.0	- 178.5	- 809.5	- 308.8	15.7	- 744.6	-590.0	- 275.9	- 670.4	-340.2	40.8	- 694.2	- 663.3	- 195.6
120 cm depth							-414.7	- 1327.6	21.5	- 959.9	- 538.4	-271.8	- 798.1	-1379.7	- 46.3	9.099 –	- 815.2	- 346.0
Annual estimated	1 weathering	fluxes to c	ompensa)	te budget d	eficits (kg h	a-1 year-	<u>(</u>											
60 cm depth				-26.2	- 19.4	- 8.9				- 18.6	- 14.8	- 6.9				- 11.6	- 11.1	- 4.9
Tut-1.																		
whole tree hards	Sung Gain the he	l																
Alliudi Judget u.	CHCIIS (NB II)	a ycar			0 6 7	-					:	1						
ov cm aeptn	- 44.1	- 0.7	- 0.7	- 52.2	- 30.8	- 10.3	- 15.7		0.1	- 11.2	- 9.8	- 5.0	3.8	- 3.5	0.7	<u></u>	- - -	- 1.4
120 cm depth							- 12.2	- 33.7	0.4	- 24.2	- 13.2	- 6.9	- 22.3	- 4.6	0.7	- 2.8	-22.0	- 4.1
Whole rotation b.	udget deficit	ts (kg ha ^{- I}	_															
60 cm depth	-883.3	- 133.4	-13.6	-643.2	- 615.2	- 205.1	- 1196.8	- 354.8	- 12.0	- 867.8	-812.1	- 304.4	-1120.3	- 425.8	2.0	- 843.6	-1048.9	- 332.0
120 cm depth							- 489.1	- 1346.2	14.7	- 968.5	- 528.2	- 275.6	-935.1	-1437.9	28.4	- 1025.4	-968.5	-357.7
Annual estimated	I weathcring	fluxes to c	ompensat	te budget de	eficits (kg h	a-1 year-1	<u>(</u>											
60 cm depth				- 32.2	- 30.8	- 10.3				- 21.7	- 20.3	- 7.6				- 14.1	- 17.5	- 5.5

stan
the
ž
age c
the
2
corresponding
lengths
rotation
sed on
þa
arc
deficits
budget
rotation
ole

60-year-old stands. SO_4^{2-} -S, TOC, Cl^- , Na⁺, and Mn²⁺ fluxes tended to be highest in the 40-year-old stand for all soil layers. NO₃⁻-N, K⁺ and Ca²⁺ tended to be highest in the 60-year-old stand, and TOC and Mn²⁺ in the 40-year-old stand at 15 cm depth. Below this depth, NO₃⁻-N, K⁺, Ca²⁺, TOC and Mn²⁺ tended to be the highest in the 40-year-old stand. Al³⁺ and Mg²⁺ were highest in the 40-year-old stand at all depths in winter and below 30 cm in autumn.

Canopy interactions were also strongly affected by stand age and seasonality (Tables 4 and 5). Uptake of NH_4^+ -N and NO_3^- -N tended to be a little higher in the 20-year-old stand than in the older stands, and the reverse was true for H₂PO₄-P. No NO₃-N uptake was observed in the 60-year-old stand and in the 40-year-old stand the uptake of nitrate was only observed during 1 year. Most of the NH_4^+ -N. NO_3^-N and $H_2PO_4^-P$ uptake occurred in spring (Fig. 4). K- and Mg-leaching were as follows: 20year-old > 40-year-old > 60-year-old stand, but Caleaching was highest in the 40-year-old stand. Values of Ca- and Mg-leaching were larger during the dormant season (autumn-winter) (Table 5). Seasonality of K^+ and TOC fluxes in TS solutions was as follows: spring > summer \ge autumn \ge winter (Fig. 4(a)Fig. 4(b)).

Canopy interactions varied annually for most elements: NH_4^+ -N and $H_2PO_4^-$ -P uptake tended to decrease from 1992/1993 to 1994/1995 but the opposite trend applied to K-, Ca- and Mg-leaching. Total deposition of all elements was larger in 1993/1994 than in the other years (Table 4).

3.4. Input-output budgets

Tables 6–8 present the results of seasonal, yearly and mean annual budgets calculated for each stand and for various rotation lengths, i.e. 20, 40 or 60 years. All calculations were made considering two criteria for output from the soil, i.e. at 60 and 120 cm depth (only 60 cm depth for the 20-year-old stand).

Mean budgets of all elements except phosphorus were negative, but weathering inputs were not calculated.

Deficit or excess of nutrients varied over the 3 years. The deficit for most elements tended to be

highest in 1994/1995 and the excess of P was highest in 1993/1994.

Annual average deficits for N, K, Ca and Mg, at 60 cm depth, varied with stand age as follows: 20 > 40 > 60. In the case of S the picture was: 40 > 20 > 60.

The most positive phosphorus budget occurred in the 60-year-old stand.

At 120 cm depth, the budgets were generally more negative than at 60 cm depth. In the 40-year-old stand, this was true for all elements except nitrogen, but in the 60-year-old stand this was the case for all elements. These two stands differed tremendously as discussed previously in the qualitative approach: cation losses in the 40-year-old stand were related to S dynamics, whereas those in the 60-year-old stand were related to N dynamics.

Seasonal budgets showed that elements were mainly lost during the dormant season. At 60 cm depth, N-deficits in the 20- and 40-year-old stands, for example, amounted respectively to 22.1 and 11.1 kg ha⁻¹ year⁻¹ in the dormant season but decreased to 4.3 and 2.4 kg ha⁻¹ year⁻¹ in the growing season. At 120 cm. N, K, Mg and S losses were reduced on average by 80% or more and Ca losses by 40% during the growing season (Table 7).

4. Discussion

4.1. Nutrient fluxes

Mean wet deposition (Pt) was compared with the RENECOFOR network data (Ulrich and Lanier, 1994). NH_4^+-N , NO_3^--N , $SO_4^{2-}-S$ and K^+ fluxes in the Beaujolais area were similar to mean values (27 sites) from this French monitoring network; Ca²⁺ and Mg²⁺ fluxes in the Beaujolais were lower. Specific comparison with the Douglas-fir site of this network (situated in the Morvan about 80 km further north), showed that rainfall input was generally lower in the Beaujolais, except for K⁺. Direct comparison between sites for total deposition (TD) was not possible but comparison of bulk throughfall indicated higher nitrogen and lower sulphur and 'base' cations in the present site. Values were very close to those from the Douglas-fir stand in Morvan for all elements, except for K^+ which was higher in the Beaujolais.

Mean total deposition to stands was close to the values observed by Mohamed (1992) in two different adult Norway spruce stands in the Vosges mountains (France); the most significant differences were the higher $SO_4^{2-}S$, Cl^- and H^+ fluxes in the Vosges than in the Beaujolais. Lamersdorf and Meyer (1993) and Matzner and Meiwes (1994), who investigated Norway spruce stands in Germany, found much higher fluxes of NH₄⁺-N, NO₃⁻-N, SO₄²⁻-S and Cl⁻ than were observed in the present study or by Mohamed (1992) (2.5-3 times higher). Atmospheric deposition of NH_4^+-N , NO_3^--N , $SO_4^{2-}-S$, Cl^- , K^+ , Ca²⁺, and Mg²⁺ in Douglas-fir stands in the Netherlands (Draaijers et al., 1988) were much higher than inputs in France or Germany. Many reasons may explain these differences, one being the presence of pollutant sources in the close vicinity. The stands from the Netherlands were close to areas where animal husbandry was common and where emissions of NH₃ were elevated. Temporal variations, as observed during the 3 year study period in the Beaujolais, or during longer studies (Matzner and Meiwes, 1994) can also affect data, as do local climatic conditions such as wind velocity, wind direction (Draaijers et al., 1988; Le Borgne, 1992) and time between rainfall events (Bastien, 1993). Moreover, even stands in similar situations, like the three stands in the present study, may differ as a result of stand development stage (Marques and Ranger, 1996).

Estimated contributions of dry deposition to total deposition of N, K, Ca, and Mg were close to values obtained by Dambrine and Prevosto (1988) for Norway spruce stands in the Vosges mountains.

Nutrient fluxes were often highest at the forest floor level and decreased with soil depth following the pattern of water flux (Fig. 4(a), Fig. 4(b), Fig.

4(c) and Fig. 4(d)). This suggests that nutrient fluxes were mainly related to mass-flow. This was confirmed by the significant correlations between fluxes of most nutrients and water flux, indicating that variation of concentrations with time was less consequential than that of water flux (Table 9). This pattern of decreasing fluxes with soil depth has been frequently observed in many forest ecosystems for both qualitative (Feller, 1977; Edmonds et al., 1991; Ranger and Nys, 1994) or quantitative (Cole and Johnson, 1977, Mohamed and Ranger, 1994) fluxes and has been attributed to the reduction of water flux, to plant uptake, and to adsorption processes in the soil's mineral or organic phases.

In the present study, the elevated nutrient fluxes at the FF level were associated with nutrient release, i.e. litter leaching and/or organic matter mineralisation.

At the P-15 level, reduction in the 'base' cation fluxes was related to plant uptake while reduction of the NH_4^+ -N flux was considered to be associated with both plant uptake and nitrification. The large flux of nitrate in the upper soil layers was interpreted as an excess in the balance between production and plant uptake. The behaviour of Al³⁺ and Si, which was similar to that of NO₃-N, was considered to be a consequence of the neutralisation of H^+ due to nitrification within the soil, by ion exchange or weathering reactions. All these aspects have been discussed in the qualitative approach of this study. It was pointed out that local mineralisation rather than nutrient transport from upper layers was the main process contributing to the quality of solutions throughout the soil profiles. Nevertheless, the behaviour of the NO_3^--N flux in the 20-year-old stand (Fig. 4(a)) suggests that NO_3^--N was transported

of stands)

Table 9											
Correlation coefficients b	between v	water a	nd elem	ent fluxes	at diffe	rent depths	in the	soil	(from	mean	data

Flux	NH ₄ -N	NO ₃ -N	SO ₄ -S	H ₂ PO ₄ -P	K	Ca	Mg	Na	Mn	Н	Al	Fe	Si	Cl	TOC
Forest floor	0.49	0.59	0.61	0.58	0.53	0.65	0.67	0.60	0.67	0.73	0.68	0.86	0.72	0.79	0.79
P-15	0.43	0.69	0.79	0.49	0.71	0.79	0.80	0.80	0.71	0.87	0.75	0.80	0.77	0.84	0.84
P-30	0.31	0.63	0.71	0.57	0.60	0.68	0.67	0.82	0.76	0.90	0.67	0.74	0.75	0.79	0.67
P-60	0.31	0.67	0.94	0.51	0.52	0.67	0.71	0.63	0.72	0.90	0.81	0.63	0.82	0.66	0.50
P-120	0.43	0.62	0.82	0.44	0.86	0.87	0.87	0.77	0.81	0.93	0.82	0.68	0.81	0.72	0.60

Forest floor, n = 109; P-15, 15 cm depth (n = 105); P-30, 30 cm depth (n = 98); P-60, 60 cm depth (n = 71); P-120, 120 cm depth (n = 60); r = 0.32 is significant at the 99% confidence level for n = 60.

from the P-15 level in autumn to the P-30 level in winter, at least in this stand. A more detailed analysis (Fig. 5) indicated that this transport from the P-15 to the P-30 level from one season to another occurred during the autumn/winter of 1994/1995 but was not observed during other periods.

At P-60, nutrient fluxes were lower than in upper soil layers but were still high for some elements, especially in the 20-year-old stand. This was due to the high soil organic matter content in the 20-year-old stand which acted as a source of nutrients to the soil solution. Consequently, the nutrient output was large. This behaviour may be a consequence of two additional factors: (i) the high soil organic matter content itself and (ii) the decomposition of this soil organic matter, which was inherited from ancient land use. The young Douglas-fir plantation probably affected the stability of the pre-existing organic components in the soil and therefore resulted in imbalance within the soil. Another hypothesis is that this imbalance resulted from the clear-cutting of the ancient forest plantation (Likens et al., 1977; Kubin, 1995). Research on the history of this plot revealed that the land was used for crops from 1824 to 1930, then planted with Norway spruce, which was logged around 1970 when the current Douglas-fir stand was planted.

At P-120, fluxes were on average lower than at P-60 but most nutrient fluxes increased sharply in autumn-winter when large water fluxes reached this layer. Many mechanisms may account for this behaviour, such as changes in plant uptake or reduction in percolation rates, which would increase the residence time of solutions at the bottom of the soil (Ugolini et al., 1977). It was hypothesised in the qualitative approach that some nutrients were produced locally but doubt remained. This quantitative study provided more information. Fluxes that occurred simultaneously at P-15 and P-120 suggest that nutrient transport from the upper to the bottom levels did occur, but the simultaneous flux reduction at 60 cm led to two hypotheses: (i) solutions were transferred directly from P-30 to P-120 along preferential channels (e.g. dead roots) or (ii) nutrients were produced locally in the deepest horizon. The location of SO₄²⁻-S flux maximum at P-120, particularly in the



Fig. 5. Monthly water and NO₃-N fluxes in the P-15 and P-30 lysimeters of the 20-year-old stand. As sampling was done every 4 weeks, there are some months where sampling occurred twice.

40-year-old stand, seems to exclude the transport hypothesis (Fig. 6(a)). Marques et al. (1996) observed a high concentration of combined sulphur compounds (probably organic compounds) in the 'capillary' solutions collected at 120 cm in the 40vear-old stand, suggesting that organic matter components were destabilised locally. Doubts still remain regarding why SO_4^{2-} -S was the dominant anion in the 40-year-old stand and NO₃-N in the 60-year-old stand (Fig. 6(a)Fig. 6(b)). It was concluded in the qualitative approach that a reduced nitrification in the 40-year-old stand was probably the reason for this specific behaviour. This could also be related to different N/S ratios in the organic pool of each stand as a consequence of differences in the former land occupation. The land of the 60-year-old stand had previously been used for crop production, since 1824 at least, whereas that of the 40-year-old stand was abandoned land in 1824, since 1870 had been afforested with Norway spruce (two generations of about 40 years) and then with Douglas-fir, more recently, in 1955.

In spite of unsatisfactory understanding of the actual processes, excess of nitrate or sulphate production over consumption led to large fluxes of 'base' cations at the P-120 level, which can be considered as unavailable for plants.

Solutions leaving the rooting zone of the 40- and 60-year-old stands contained on average 4.9 mg 1^{-1} of Al³⁺. As only total Al was measured, Al³⁺ probably represented about 2.5 mg 1^{-1} , according to the solution's pH (Driscoll, 1989). Analyses from a spring (11 monthly samplings) showed that the pH of surface waters was close to neutrality and their Al content was very low. This indicates that drainage water was neutralised somewhere in the subsoil, safeguarding the quality of surface waters (data not shown).

4.2. Input-output budgets

Atmospheric inputs into the Douglas-fir stands were among the medium range of values for forest ecosystems in France. Outputs were high and due to nutrient losses by seepage rather than to biomass immobilisation.

Annual variations in the nutrient budget were large during the 3 years of investigation. Annual

deficits tended to be correlated with the amount of rainfall: 1992/1993 < 1994/1995. Deficits were generally highest during the dormant season despite the lack of nutrient immobilisation by stands. This was related to large water fluxes, which typically belong to the cold season under temperate climates, and to drainage of elements at the bottom of the soil (Fig. 4(a), Fig. 4(b), Fig. 4(c) and Fig. 4(d)). Atmospheric inputs were totally removed by seepage; in addition, a large amount of nitrate produced within the soil was drained away and this occurred indistinctly throughout the dormant season.

Nutrient outputs during the growing season were more strongly related to stand immobilisation than to drainage, in the 40- and 60-year-old stands (Table 7). Nevertheless, the reverse occurred in the younger stand because of its specific soil dynamics as discussed before. Nitrate losses during the growing season suggest that nitrification largely exceeded forest stand requirements. Negative cation budgets were just a consequence of the nitrogen dynamics. More sulphate was lost through drainage than through immobilisation irrespective of stand age, and this was due to a relatively high atmospheric input coupled to a low immobilisation rate.

Theoretical budgets assuming forest rotation lengths of 20, 40 or 60 years were calculated for output levels at 60 and 120 cm depth. Two harvesting systems were taken into account: harvesting of stems only or whole tree harvesting (biomass and nutrient content data were presented in detail in Ranger et al., 1995).

For a 60 year rotation (stem only harvesting), at 60 cm depth, the nutrient deficit (atmospheric input – immobilisation – drainage) amounted to 670 kg N, 694 kg K, 663 kg Ca, 296 kg Mg and 346 kg S (Table 8). Balancing this budget would require an annual ecosystem N accumulation of 11 kg and an input by weathering of about 12 kg K, 11 kg Ca, 5 kg Mg and 6 kg S ha⁻¹. N- and S-sources are very difficult to estimate since they are related to the behaviour of microorganisms, therefore only internal soil fluxes of K, Ca, and Mg are discussed here. Bonneau et al. (1991) and Mohamed (1992) estimated weathering rates, in soils with a similar 'base' cation content, at 6.8×10^{-5} kg K, 2×10^{-4} kg Ca and 5×10^{-5} kg Mg ha⁻¹ year⁻¹. Applying these rates to the total amount of elements in the fine earth of the soils (Table 1) yields a weathering flux of 10.5 kg K, 0.6 kg Ca, and 1.2 kg Mg (assuming the stones did not contribute to the weathering flux). Not enough data were available to confirm weathering fluxes but

preliminary results suggest that estimated values were reasonable for K but not for Ca and Mg (Ezzaïm et al., 1996). It was thus hypothesised that the weathering flux was sufficient to conpensate for K deficits.



Fig. 6. (a) Temporal variations of nitrate, sulphate and estimated water fluxes in the plate lysimeters of the 40-year-old stand. As sampling was done every 4 weeks, there are some months where sampling occurred twice. (b) Temporal variations of nitrate, sulphate and estimated water fluxes in the plate lysimeters of the 60-year-old stand. As sampling was done every 4 weeks, there are some months where sampling occurred twice.



Fig. 6 (continued).

Conservative values for Ca and Mg weathering fluxes were adopted, i.e. 3.5 kg Ca ha⁻¹ year⁻¹ and 2.5 kg Mg ha⁻¹ year⁻¹. Even with this conservative hypothesis, deficits at the end of a 60 year rotation amounted to 450 kg Ca and 140 kg Mg. This means that the soil has lost about 50% of the Ca and 42%

of the Mg present in the initial pool of available nutrients over the last 60 years.

For a 60 year rotation, at 120 cm depth, nutrient losses amounted to 798 kg N, 991 kg K, 815 kg Ca, 346 kg Mg and 1380 kg S. Annual weathering fluxes necessary for compensating these deficits should be

	N	S	Р	К	Ca	Mg
Stem only harvesting						
Total input	1141.3	747.2	97.8	208.2	339.9	60.9
Total immobilisation	402.4	26.0	32.7	197.1	262.4	34.6
Total drainage fluxes	1135.3	1215.5	32.3	404.5	512.3	241.5
Budget for the forest rotation	- 396.4	- 494.4	32.8	- 393.4	- 434.8	-215.2
Estimated total weathering				630	210	150
Budget including weathering input				236.6	- 224.8	65.2
Whole tree harvesting						
Total input	1141.3	747.2	97.8	208.2	339.9	60.9
Total immobilisation	852.0	108.0	72.0	346.0	648.0	70.0
Total drainage fluxes	1135.3	1215.5	32.3	404.5	512.3	241.5
Budget for the forest rotation	-846.0	- 576.4	-6.5	- 542.3	-820.4	- 250.6
Estimated total weathering				630	210	150
Budget including weathering input				87.7	- 610.4	- 100.6

Table 10 Estimated nutrient budgets for a second 60-year Douglas-fir rotation at the Beaujolais site (data expressed in kg ha⁻¹).

16.5 kg K, 13.6 kg Ca and 5.8 kg Mg ha⁻¹ year⁻¹. Compared with the budget calculated at 60 cm, deficits at 120 cm depth did not increase too much. Assuming fluxes twice as large as fluxes at 60 cm depth, the deficit of Mg may be almost balanced but not that of Ca, of which the amount initially available would be depleted by 61%.

Shorter rotation lengths may deplete 'base' cations dramatically; estimated deficits (at a 60 cm depth) would amount to 26 kg K, 19 kg Ca and 9 kg Mg ha⁻¹ year⁻¹ for a 20 year rotation, and to 19 kg K, 15 kg Ca and 7 kg Mg ha⁻¹ year⁻¹ for a 40 year rotation (Table 8). These estimations agree with previous works with respect to the negative impact of short rotations on nutrient budgets (Switzer and Nelson, 1973; Kimmins, 1974, 1977; Leaf, 1979), and may be more drastic since losses due to clear-cutting and regeneration operations were not estimated (Likens et al., 1977; Dyck et al., 1994; Kubin, 1995).

Deficits were very much higher for whole tree harvesting (WTH) than for stem-only harvesting (SH) as a consequence of the huge amount of nutrients immobilised in the tree crown (Ranger et al., 1995). The cumulation of increased removal from harvesting and a shorter rotation led to very high nutrient demands. The weathering flux required for compensating the Ca-deficit in a 20 year rotation associated with WTH would be 1.6 times greater than under SH (Table 8). The nutrient budget for a new identical forest rotation (SH vs. WTH) was simulated to estimate the soil depletion during the next 60 years (Table 10). The following assumptions were made:

- 1. the impact of the early stage of development on organic matter mineralisation would be more or less stabilised and the drainage flux at 60 cm depth was estimated as the weighted average values observed today in the 40- and 60-year-old stands;
- 2. atmospheric inputs and stand immobilisation would remain similar to those of the first rotation. Deficits under SH would amount to 395 kg K, 435 kg Ca and 215 kg Mg, per harvesting cycle, representing annual losses of 6.6 kg K, 7.2 kg Ca and 3.6 kg Mg ha $^{-1}$. The weathering flux values assumed above should be able to compensate for K losses, barely compensate for Mg losses but cannot compensate for Ca losses. The Ca deficit at the end of the second Douglas-fir rotation would amount to about 225 kg. The initial available storage before the first rotation, which was estimated at 905 kg Ca (455 kg in the present storage plus 450 kg lost during the first rotation) would be reduced to 230 kg Ca (455 kg in the present storage minus 225 kg lost during the second rotation). Total losses during the 120 year period represent 75% of the initial storage.

With WTH, the imbalance of the ecosystem would be almost doubled for N, K and Ca and would even extend to P whose budget was positive during the first rotation. The weathering fluxes would barely compensate for the K and Mg deficits and in no way compensate for the Ca deficit. The present available Ca-storage of about 455 kg would be sold out. The Ca nutrition of the next generation would undoubtedly be affected.

If the budget results are confirmed after evaluation of the weathering fluxes (work in progress), severe changes would occur in the soil function whatever silvicultural scenario is adopted. The following changes in soil fertility would occur:

- 1. reduction of available Ca pool;
- desaturation of the soil exchange complex, expressed as an enhancement of Al retention and a reduction of 'base' cations (BC) on the exchange sites;
- 3. decrease of BC/Al ratios in the soil solution accompanied by the well known consequences for plant nutrition (Rost-Siebert, 1983; Huttermann and Ulrich, 1984; Boudot et al., 1995).

5. Conclusions

The pattern of fluxes obtained after 3 years of observation did not give a definitive picture for the behaviour of this Douglas-fir ecosystem. Owing to large temporal variations, monitoring should continue for several years to validate the present trends and to reduce the weight of singular climatic years.

The chronosequence approach proved to be a very useful tool to discriminate differences in the plantsoil relationships related to stand development. This was true in spite of uncertainties regarding the complete similarity of plot characteristics at time zero.

Differences in historical land usage of the plots certainly affected the dynamics of elements in the soil today. However, it is difficult to determine the extent of this influence.

Negative input-output budgets for N, S, Ca and to a lesser extent for Mg, suggest that the soil chemical dynamics were imbalanced, probably because of the impact of the present vegetation. High rates of nitrification and organic sulphur oxidation indirectly induced strong losses of nutrients out of the rooting zone. Soil acidification results more from excess nitrification throughout the soil profile than from external acidic deposition. An unsolved question is: why was sulphur oxidation the acidifying source in the 40-year-old stand and nitrification in the 60-year-old stand?

Negligible impact on surface water chemistry was determined by monitoring a spring but requires confirmation by more extensive investigations.

Stands presently had a satisfactory health status, identified from direct morphological observations and foliar analysis (Ranger et al., 1995). Nutrient supply was large enough to sustain the present high production rate, but negative budgets suggest that this ecosystem is not stable and that long-term productivity cannot be ensured if a similar trend occurs at the next rotation.

The budget deficits revealed that the present stand growth is obtained at the expense of the soil organic matter compartment, thereby promoting soil desaturation which may seriously compromise future forest growth. Extrapolation of the budget to the next rotation is merely speculative. The trend of decrease in the budget deficit with stand age suggests that the effect of the vegetation change will be reduced at the next rotation but the impact of stand development may remain.

The present results did not lead to definitive conclusions about the impact of high productive Douglas-fir stands on soil dynamics and surface water quality. Nevertheless, the present trend indicates a very high loss of 'base' cations from the soil profile, which may decrease soil fertility within a rather short time scale.

A sustainable management strategy for these Douglas-fir stands should exclude whole tree harvesting as this would quickly lead to large soil nutrient deficiencies. Silvicultural practices like remnant windrowing, remnant burning or full-tree skidding would lead to the same result and the maintenance of sustainability would require nutrient input by fertilisation.

Acknowledgements

We would like to thank the technical staff of the 'Cycles biogéochimiques' research team for assistance, Dominique Gelhaye for managing the field work, Séverine Bienaime, Louisette Gelhaye and Benoît Pollier for laboratory measurements; the forest managers, the 'Office National des Forêts', and particularly G. Poly and B. Jobard for providing facilities during investigation and sampling; Philip Butler for revising the English; and the 'Ministère de l'Agriculture, Division de l'Espace Rural et de la Forêt', for providing the financial support for this research project.

References

- Aussenac, G. and Boulangeat, C., 1980. L'interception des précipitations et évapotranspiration réelle dans des peuplements de feuillu (*Fagus silvatica* L.) et de résineux (*Pseudotsuga menziesii* (Mirb.) Franco). Ann. Sci. For., 37: 91–107.
- Bastien, J., 1993. Modélisations des concentrations et flux en éléments chimiques présents dans les pluies et pluviolessivats en fonction de facteurs météorologiques. Université Nancy I/INRA, D.E.S.S. d'Ingénierie mathématique et d'outils informatiques, Nancy, 50 pp.
- Beets, P.N., Terry, T.A. and Manz, J., 1994. Management systems for sustainable productivity. In: W.J. Dyck, D.W. Cole and N.B. Comerford (Editors), Impacts of Forest Harvesting on Long-term Site Productivity. Chapman and Hall, London, pp. 219–246.
- Bonneau, M., Dambrine, E. and Ranger, J., 1991. Apports acides et cycle des cations dans des pessières du Nord-Est: interêt de bilans saisonniers. Sci. Sol. 21: 125–145.
- Boudot, J.-P., Becquer, T., Merlet, D., Rouiller, J., Ranger, J., Dambrine, E. and Mohamed, D.A., 1995. Potential role of aluminium toxicity in nutrient deficiencies as related to forest decline: an assessment of soil solution data from the Vosges mountains. In: G. Landmann and M. Bonneau (Editors), Forest Decline and Atmospheric Deposition Effects in the French Mountains. Springer, Berlin/Heidelberg, pp. 270–285.
- Breda, N., 1994. Analyse du fonctionnement hydrique des chênes sessile (*Quercus petraea*) et pédonculé (*Quercus robur*) en conditions naturelles; effets des facteurs du millieu et de l'éclaircie. Université Nancy I/INRA, Thèse de Doctorat en Biologie Forestière, Nancy, 59 pp. + annexes.
- Bredemeier, M., 1988. Forest canopy transformations of atmospheric deposition. Water, Air, Soil Pollut., 40: 121–138.
- Cole, D.W. and Johnson, D.W., 1977. Atmospheric sulfate additions and cations leaching in a Douglas Fir ecosystem. Water Resour. Res., 13: 313–317.
- Dambrine, E. and Prevosto, B., 1988. Flux des éléments minéraux dans un écosystème forestier d'altitude soumis à la pollution atmosphérique, relation avec le dépérissement. Rapp. Sci., DEFORPA, Décembre 1988.
- Draaijers, G.P.J., Ivens, W.P.M. and Bleuten, W., 1988. Atmospheric deposition in forest edges measured by monitoring canopy throughfall. Water, Air, Soil Pollut., 42: 129–136.
- Driscoll, C.T., 1989. The chemistry of aluminum in surface waters. In: G. Sposito (Editor), The Environmental Chemistry of Aluminum. CRC Press, Boca Raton, pp. 242–273.

- Duchaufour, P. and Bonneau, M., 1959. Une methode nouvelle du dosage du phosphore assimilable dans les sols forestiers. Bull. Assoc. Française pour l'Etude du Sol, 4: 193–198.
- Dyck, W.J., Cole, D.W. and Comerdorf, N.B. (Editors), 1994. Impacts of forest harvesting on long-term site productivity. Chapman and Hall, London, 368 pp.
- Edmonds, R.L., Thomas, T.B. and Rhodes, J.J., 1991. Canopy and soil modification of precipitation chemistry in a temperate rain forest. Soil Sci. Soc. Am. J., 55: 1685–1693.
- Ezzaïm, A., Turpault, M.-P. and Ranger, J., 1996. Répartition des nutriments dans un sol brun acide développé sur tuf (Beaujolais, France). Conséquences pour l'évolution de la fertilité minérale à long terme. Annales des Sciences Forestières, in press.
- Feger, K.H., 1988. Historical changes in catchment use. In: H. Barth (Editor), Effects of land use in catchments on the acidity and ecology of natural surface waters. CEC, Air Pollut. Rep., 13: 65-74.
- Feller, M.C., 1977. Nutrient movement through western hemlock-western redcedar ecosystems in southwestern British Columbia. Ecology, 58: 1269-1283.
- Granier, A., 1987. Evaluation of transpiration in a Douglas-fir stand by means of sap flow measurements. Tree Physiol., 3: 309-320.
- Huttermann, A. and Ulrich, B., 1984. Solid phase-solution-root interactions in soils subjected to acid deposition. Phil. Trans. R. Soc. London, Ser. B, 305: 353–368.
- Kimmins, J.P., 1974. Sustained yield, timber mining, and the concept of ecological rotation; a British Columbian view. For. Chron., Feb.: 27–31.
- Kimmins, J.P., 1977. Evaluation of the consequences for future tree productivity of the loss of nutrients in whole-tree harvesting. For. Ecol. Manage., 1: 169–183.
- Kubin, E., 1995. The effect of clear cutting, waste wood collecting and site preparation on the nutrient leaching to groundwater. In: L.O. Nilsson, R.F. Hüttl and U.T. Johansson (Editors). Proc. of the CEC/IUFRO Symp. on Nutrient Uptake and Cycling in Forested Ecosystems, Halmstad, Sweden, 7-10 June 1993, Kluwer, Dordrecht, pp. 661-670.
- Lamersdorf, N. and Meyer, M., 1993. Nutrient cycling and acidification of a northwest German forest site with high atmospheric nitrogen deposition. For. Ecol. Manage., 62: 323-354.
- Leaf, A.L. (Editor), 1979. Proc. of Symp. on Impact of Intensive Harvesting on Forest Nutrient Cycling, School of Forestry. State University of New York, Syracuse, 421 pp.
- Le Borgne, A., 1992. Étude de la composition chimique des precipitations à decouvert en relation avec les types de temps et le climat de pollution. Le cas de la station d'Aubure (Vosges). Université de Paris I, Mémoire de D.E.A, de Géographie et de l'Environnement Global, Paris, 50 pp.
- Likens, G.E., Bormann, F.M., Pierce, R.S., Eaton, J.S. and Jonhson, N.M., 1977. Biogeochemistry of a Forested Ecosystem. Springer, New York, 146 pp.
- Lovett, G.M. and Lindberg, S.E., 1984. Dry deposition and canopy exchange in a mixed oak forest as determined by analysis of throughfall. J. Appl. Ecol., 21: 1013–1027.
- Marques, R. and Ranger, J., 1996. Nutrient dynamics in a chronosequence of Douglas-fir (*Pseudotsuga menziesii* (Mirb).

Franco) stands on the Beaujolais Mounts (France). 1. Qualitative approach. For. Ecol. Manage., in press.

- Marques, R., Ranger, J., Gelhaye, D., Pollier, B., Ponette, Q. and Gœdert, O., 1996. Comparison of chemical composition of soil solutions collected by zero-tension plate lysimeters with those from ceramic-cup lysimeters in a forest soil. Eur. J. Soil Sci., 47: 407-417.
- Matzner, E. and Meiwes, K.J., 1994. Long-term development of element fluxes with bulk precipitation and throughfall in two German forests. J. Environ. Qual., 23: 162–166.
- Mohamed, A.D., 1992. Rôle du facteur édaphique dans le fonctionnement biogéochimique et l'état de santé de deux pessières vosgiennes. Effet d'un amendement calci-magnésien. Université de Nancy I/INRA-CRF, Thèse Doctorat, Nancy, 206 pp.
- Mohamed, A.D. and Ranger, J., 1994. The biogeochemical cycle in a healthy and highly productive Norway spruce (*Picea abies*) ecosystem in the Vosges, France. Can. J. For. Res., 24: 839–849.
- Page, G., 1970. Some effects of conifer crops on soil properties. Comm. For. Rev., 47: 52-62.
- Ranger, J. and Nys, C., 1994. The effect of spruce (*Picea abies* Karst.) on soil development: an analytical and experimental approach. Eur. J. Soil Sci., 45: 193–204.
- Ranger, J., Marques, R., Colin-Belgrand, M. and Flammang, N., 1995. The dynamics of biomass and nutrient accumulation in a Douglas-fir (*Pseudotsuga menziesii* Franco) stand studied using a chronosequence approach. For. Ecol. Manage., 72: 167– 183.

- Rost-Siebert, K., 1983. Aluminium Toxizität und Toleranz am Keimpflanzen von Fichte und Buche. Allgem. Forstz., 5: 686–689.
- Soil Survey Staff, 1975. Soil Taxonomy. USDA Soil Conservation Service. US Government Printing Office, Washington, DC, 754 pp.
- Switzer, G.L. and Nelson, L.E., 1973. Maintien de la productivite avec des revolutions courtes. In: Colloque International sur l' Utilisation des Engrais en Forêt, FAO/IUFRO, Paris, 3–7 December 1973, pp. 355–381.
- Ugolini, F.C., Minden, R., Dawson, H. and Zachara, J., 1977. An example of soil processes in the *Abies amabilis* zone of Central Cascades, Washington. Soil Sci., 124: 291-302.
- Ulrich, B., 1983. Interaction of forest canopies with atmospheric constituents: SO₂, alkali and earth alkali cations and chloride.
 In: B. Ulrich and J. Pankrath (Editors), Effects of Accumulation of Air Pollutants in Forest Ecosystems. D. Reidel, Dordrecht, pp. 33–45.
- Ulrich, E. and Lanier, M., 1994. Depots atmospheriques et concentrations des solutions du sol (sous reseau Cataenat) Rapport scientifique sur l'année 1993. Fontainebleau, Office National des Forêts-Département des Recherches Techniques, 117 pp.
- Villette, S., 1994. Etablissement du bilan hydrique sur une chronoséquence de peuplements de Douglas (*Pseudotsuga menziesii* (Mirb.) Franco) du Beaujolais. Essai de modélisation. ENITA/INRA-CRF, Nancy, Mémoire de fin d'études, 50 pp.