



HAL
open science

Essential role for the homeoprotein vHNF1/HNF1beta in visceral endoderm differentiation.

Catherine Coffinier, Dominique Thepot Thépot, C Babinet, M Yaniv, J Barra

► To cite this version:

Catherine Coffinier, Dominique Thepot Thépot, C Babinet, M Yaniv, J Barra. Essential role for the homeoprotein vHNF1/HNF1beta in visceral endoderm differentiation.. *Development* (Cambridge, England), 1999, 126 (21), pp.4785-94. hal-02690542

HAL Id: hal-02690542

<https://hal.inrae.fr/hal-02690542>

Submitted on 1 Jun 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Essential role for the homeoprotein $vHNF1/HNF1\beta$ in visceral endoderm differentiation

Catherine Coffinier¹, Dominique Thépot^{1,*}, Charles Babinet², Moshe Yaniv^{1,‡} and Jacqueline Barra²

¹Unité des Virus Oncogènes, CNRS URA 1644, and ²Unité de Biologie du Développement, CNRS URA 1960, Institut Pasteur, 25 rue du Dr Roux, F-75724 Paris Cedex 15, France

*Present address: Laboratoire de Biologie Cellulaire et Moléculaire, Unité de Différenciation Cellulaire, INRA, F-78352 Jouy en Josas, France

‡Author for correspondence (e-mail: yaniv@pasteur.fr)

Accepted 26 July; published on WWW 6 October 1999

SUMMARY

$vHNF1/HNF1\beta$, a member of the divergent HNF1/ $vHNF1$ homeoprotein family, is expressed in polarized epithelia of several adult organs and may participate in controlling the transcription of specific genes. In addition to this late requirement, $vHNF1$ may play earlier roles during development, as it is first expressed in the visceral endoderm at the onset of gastrulation. In order to shed light on its function during embryogenesis, we have inactivated the murine gene by homologous recombination. The homozygous mutation results in embryonic lethality by day 7.5 of development and $vHNF1^{-/-}$ embryos display a disorganized visceral endoderm and a significantly reduced size. Studies of ES cell differentiation and aggregation with tetraploid morulae establish that $vHNF1$ expression is

essential for visceral endoderm differentiation, both in vitro and in vivo. Analysis of differentiation markers confirms that $vHNF1$ is part of a genetic network that directs the expression of $HNF4$ and downstream endodermal genes. Furthermore, the complementation of the mutant embryos with wild-type visceral endoderm rescues the day 7.5 lethality and reveals an additional phenotype linked to $vHNF1$ later expression. The examination of chimeric embryos suggests that $vHNF1$ expression might be cell-autonomously required in the gut for the proper morphogenesis of the embryo.

Key words: Visceral endoderm, Embryonic lethality, Tetraploid rescue, Homeoprotein, Gene knock-out.

INTRODUCTION

A surprisingly limited number of steps during mouse development are essential for the survival of the embryo (Copp, 1995). Early embryonic lethality is frequently observed when the interactions between the mother and the embryo are perturbed. For example, the inactivation of genes encoding for Mash-2 or Ets-2 results in embryonic failure due to an impaired trophoblastic function (Guillemot et al., 1994; Yamamoto et al., 1998). Differentiation of the visceral endoderm is another important step of embryonic development. The visceral endoderm plays a crucial role by supporting the metabolism and the growth of the embryo. In addition, its anterior territory has recently been shown to function as an organizer for head structures (Beddington and Robertson, 1998). A number of transcription factors are known to be specifically expressed in the visceral endoderm around gastrulation, among them $HNF3\beta$ and $HNF4\alpha$ (Duncan et al., 1994; Monaghan et al., 1993; Taraviras et al., 1994). For both $HNF3\beta$ and $HNF4\alpha$ mutations, a defect in the visceral endoderm has been suggested (Ang and Rossant, 1994; Chen et al., 1994; Weinstein et al., 1994). These mutations lead to embryonic lethality between days 8.5 and 10.5 of development. $HNF3\beta$ expression in the anterior visceral endoderm is important for

the patterning of the anteroposterior axis (Dufort et al., 1998), while $HNF4\alpha$ is involved in regulating the metabolic functions of extraembryonic visceral endoderm (Duncan et al., 1998).

Interestingly, $HNF3$ and $HNF4$ were initially characterized as liver-enriched transcription factors. Indeed, the liver and the visceral endoderm perform several equivalent functions, such as protein secretion and metabolite absorption; they also share an important number of differentiation markers (Meehan et al., 1984). Hence, it is plausible that, in addition to $HNF3$ and $HNF4$, other classes of Hepatocyte Nuclear Factors could be involved in visceral endoderm differentiation. Good candidates included the two closely related atypical homeoproteins, $HNF1$ and $vHNF1$ (also named $HNF1\alpha$ and $HNF1\beta$, respectively). These factors bind to identical DNA sequences, as either homodimers or heterodimers (Tronche and Yaniv, 1992). $vHNF1$ mRNA is detectable in the visceral endoderm from early developmental stages, while $HNF1$ is first expressed in the yolk sac and in the liver bud (Blumenfeld et al., 1991; Cereghini et al., 1992). Later on, they are expressed throughout organogenesis of the liver, the kidney and the pancreas where their expression is maintained up to the adult stage. Still, $HNF1$ and $vHNF1$ display some specificity: for instance, $vHNF1$ and not $HNF1$ is expressed in the lungs (Cereghini et al., 1992; Ott et al., 1991). This suggests the existence of specific targets for

each gene, and indeed *HNF1* inactivation has provided the first evidence for the existence of *HNF1*-specific target genes in these tissues (Pontoglio et al., 1996).

HNF1^{-/-} mice develop normally but fail to transcribe several hepatic genes, are defective in renal glucose reabsorption and suffer from type II diabetes (Pontoglio et al., 1998). Since *vHNF1* expression precedes *HNF1* synthesis, it has been suggested that *vHNF1* might compensate for *HNF1* function during development; alternatively, *HNF1* could be dispensable during embryogenesis. To address these issues, we have performed the inactivation of the *vHNF1* gene. *vHNF1*^{-/-} embryos die around day 7.5 post-coitum (E7.5) at a time where *vHNF1* is specifically expressed in the extraembryonic visceral endoderm. The defect in visceral endoderm differentiation can be partially rescued by wild-type visceral endoderm. Further development results in *vHNF1*^{-/-} embryos of a normal size but displaying severe abnormalities.

MATERIALS AND METHODS

Targeting of the *vHNF1* locus

vHNF1 genomic clones were isolated from a 129/SvJ mouse phage library. Genomic fragments of 3.3 kb and 2.2 kb flanking the first exon were introduced into the recombination vector pPNT (Tybulewicz et al., 1991). A NLS-*lacZ* cassette, described by Pontoglio et al. (1996), was introduced at the initiation codon of *vHNF1* into a *NcoI* site generated using the primers: 5'-GTCCAAAGGTACCCTAAAAGT-TTGG-3' and 5'-CTTGACACGGTCCAAGGACG-3'. CK35 ES cells (Kress et al., 1998) were electroporated with the *NotI*-linearized vector as described by Pontoglio et al. (1996). G418-resistant clones were analyzed by Southern blot after *XbaI* digestion using a *EcoRI*-*FokI* 5' external probe and a *BamHI*-*EcoRI* 3' external probe. Single-copy integration was verified using a *PstI* *neo* probe. ES cell injections into blastocysts were accomplished as previously described (Pontoglio et al., 1996).

Mouse and ES cell DNA analysis

DNA from either mouse tails or ES cells was directly precipitated with isopropanol after proteinase K treatment. A three-way PCR was performed with a sense primer hybridizing to the *vHNF1* promoter (5'-CCCAACTGTCCAGCTCTCTACC-3') and two reverse primers in the *vHNF1* first exon (5'-CTGAGCATCCGGTCCACCTCG-3') and in the *lacZ* gene (5'-GGGAAGGGCGATCGGTGCGG-3') generating respectively a wild-type 900 bp band and a mutant *vHNF1/lacZ* 800 bp band (arrows in Fig. 1A).

X-gal staining and histological analysis

Embryos were collected at different gestation times, plugs in the morning being counted as day 0.5 post-coitum (E0.5). Embryos were fixed for 10 minutes in 4% paraformaldehyde and abundantly washed in PBS. Whole-mount X-gal staining was performed overnight as described by Pontoglio et al. (1996). Genotyping of the embryos was made using the three-way PCR described above, on DNA isolated from ectoplacental cones cultured for 1 week in microwells with DMEM 10% FCS (Ang and Rossant, 1994). Historesin (Leica)-embedded embryos were thin-sectioned with a microtome and counterstained with Safranin.

Homozygous ES cells generation and embryoid body culture

vHNF1/lacZ^{-/-} ES cells were isolated from *vHNF1/lacZ*^{+/-} clones by increasing G418 selection (1.5 mg/ml) (Mortensen et al., 1992) and genotyped using PCR. Either *vHNF1*^{+/+}, *vHNF1*^{+/-} or *vHNF1*^{-/-} ES cells were mildly trypsinized and put in suspension culture in bacterial

plates (Rudnicki and McBurney, 1987). Embryoid bodies (EB) were collected after 14 days culture and stained with SJA lectin (*Sophora japonica* agglutinin, Sigma) as described by Soudais et al. (1995).

RT-PCR analysis on embryoid bodies and embryos

Total RNAs were prepared from EB according to Koopman (1993). RT-PCR analysis was performed including [³²P]dATP incorporation as described by Duncan et al. (1997). Either 1 µg from EB RNA or from one wild-type embryo was used per RT reaction; homozygous embryos were pooled as sets of four embryos. The PCR conditions and the primer sequences used for *HNF4*, *HNF1*, *TFN*, *TTR*, *AFP*, *GATA4*, *Apo A1* and *Apo AIV* were described by Duncan et al. (1997). The following oligonucleotides were used to detect the expression of *HNF3α*: (5'-GTAGACAGTAGGGGCTC-3', 5'-GGGGAATCCTT-TAAACGG-3'); *HNF3β*: (5'-GCCTGAGCCGCGCTCGGGAC-3', 5'-GGTGCAGGGTCCAGAAGGAG-3'); *HNF3γ*: (5'-GCTTCC-GGGTATGTAGCCCC-3', 5'-GCAAGGCCAGTAGGAGCC-3'); *GATA-6*: (5'-CAGCCCACGTTACGATGAACG-3', 5'-AAAATGC-AGACATAACATTCC-3'). *HPRT* primers were used to normalize the RT-PCR reaction (Wiles, 1993) and RT performed without the reverse transcriptase was used as a negative control.

Production of a *vHNF1*-specific antiserum and western blot analysis

A GST-fusion protein containing *vHNF1* amino acids 321 to 378 was used to immunize rabbits. Immune serum was purified by affinity first on a GST column then on a GST-*vHNF1* column. Purified antiserum was tested on nuclear kidney extract and in vitro *vHNF1* protein, and recognized specifically a 69 kDa band corresponding to the expected molecular weight of the *vHNF1* protein. Western blot analysis was performed with this antiserum using 30 µg of total protein extracts from 14-day-old embryoid bodies.

Tetraploid aggregation and chimeras

Tetraploid embryos were generated from F₁ B6/SJL 2-cell embryos using an electric shock to fuse the blastomers. They were cultured in vitro up to the 4-cell stage and then aggregated by pairs of two embryos with a clump of ES cells (Nagy et al., 1993). Resulting blastocysts were reimplanted the following day into the uterus of day 2.5 p.c. pseudopregnant mice. Chimeric embryos were obtained by the injection of 7-8 ES cells into wild-type blastocysts.

RESULTS

Generation of *vHNF1* mutant alleles

To produce a null allele of the *vHNF1* gene and prevent the production of any truncated protein that could interact with *HNF1*, we chose to delete the first exon of *vHNF1* that encodes the dimerization domain. Genomic DNA covering the 5' regulatory regions and the two first exons of *vHNF1* was isolated from a 129/SvJ mouse DNA library. We constructed a recombination vector in which the first exon of *vHNF1* was replaced by an NLS-*lacZ* reporter gene placed at the initiation codon (Fig. 1A). After electroporation into CK35 embryonic stem (ES) cells (Kress et al., 1998) and double selection (*neo* *tk*), seven recombinant clones were identified by Southern blot analysis, which contained a *vHNF1/lacZ* recombinant allele (Fig. 1B). All of them carried a single-copy integration at the locus as revealed by Southern blot analysis with a *neo* probe (data not shown).

ES cells homozygous for the *vHNF1* mutation were isolated from two independent *vHNF1*^{+/-} clones after culture in high G418 concentration (Mortensen et al., 1992). A large majority of the clones isolated had spontaneously replaced the wild-type




Fig. 1. Targeted inactivation of the murine *vHNF1* gene. (A) Strategy used for inactivating the *vHNF1* gene by homologous recombination in ES cells. Clone analysis was performed using the 5' and 3' probes and *Xba*I sites, two sites external to the construct and a third site introduced by the recombination. The arrows indicate the positions of the primers used for the PCR analysis of embryos and mice. X, *Xba*I; R, *Eco*RI; Pml, *Pml*I; H, *Hind*III; B, *Bam*HI. (B) Southern blot analysis of the ES clones isolated after positive-negative selection using the 5' and 3' probes. Two *vHNF1* heterozygous (+/-) and four wild-type (+/+) clones are presented. (C) PCR analysis of a litter from *vHNF1*^{+/-} intercross collected at E7.5, featuring wild-type (+/+), heterozygous (+/-) and homozygous (-/-) embryos. The arrows indicate the bands characteristic of the wild-type (wt) and the *vHNF1/lacZ* (*lacZ*) alleles. (D) Western blot analysis of total protein extracts from differentiated *vHNF1*^{+/-}, *vHNF1*^{+/-} and *vHNF1*^{-/-} ES cells using a *vHNF1*-specific antiserum. The arrow indicates the 69 kDa *vHNF1*-specific band, and the star, a non-specific band.

allele by a second *vHNF1/lacZ* allele, as confirmed using PCR and Southern blot analysis (Fig. 1C; data not shown). As *vHNF1* synthesis is induced after *in vitro* differentiation of ES cells (Abe et al., 1996), western blot analysis was performed on differentiated *vHNF1*^{+/+}, *vHNF1*^{+/-} and *vHNF1*^{-/-} ES cells to confirm that the *vHNF1/lacZ* allele was indeed a null allele. As shown in Fig. 1D, *vHNF1* protein was absent in total protein extracts from differentiated *vHNF1*^{-/-} ES cells. In contrast, a 69 kDa band corresponding to the *vHNF1* protein was detected in the *vHNF1*^{+/+} and *vHNF1*^{+/-} cell extracts.

A second construct containing three loxP sites flanking the first exon of *vHNF1* and the *neo* cassette was introduced by homologous recombination using the same strategy as for the *vHNF1/lacZ* construct. After isolation of recombinant clones, ES cells were transiently transfected with the Cre-expressing vector pIC-Cre (Gu et al., 1993) and clones where both the exon and the selection gene have been deleted, were isolated (data not shown). Such a deletion is predicted to result in a null allele as the promoter, the transcription initiation site and the first exon of *vHNF1* were all eliminated.

For both the *vHNF1/lacZ* and the *vHNF1/del* alleles, one recombinant ES clone was injected into blastocysts; the two alleles were successfully transmitted through germline and gave rise to normal and fertile heterozygous animals. As identical phenotypes were observed for both strains of mice at the homozygous state, only results concerning the *vHNF1/lacZ* strain are presented here (noted below simply as *vHNF1*^{-/-}).

vHNF1/lacZ expression during early development

Heterozygous mice carrying the *vHNF1/lacZ* allele were used

to follow the expression of *vHNF1* during embryonic development by X-gal staining of whole embryos. *vHNF1/lacZ* was first detected in the visceral endoderm at day 7.5 post-coitum (E7.5) (Fig. 2A). Notably, β -galactosidase activity was detected exclusively in the extraembryonic part of the visceral endoderm (Figs 2A, 4A); this region is formed of columnar cells and differs morphologically from the squamous epithelium surrounding the embryo. Later on, the extraembryonic visceral endoderm gives rise to the yolk sac endoderm, which also expressed *vHNF1/lacZ* (Fig. 2B). In addition to this very early expression in extraembryonic tissues, *vHNF1/lacZ* was detected in the embryo itself from E8. It was visible as early as the 2-somite stage in the neural tube and a few hours later in the invaginating gut (Fig. 2B).

The X-gal staining in the neural tube was very dynamic. It was detected in the central part of the neural folds as early as the 2-somite stage with only scattered nuclei marked (Fig. 2B). As the neural folds merge, the number of positive cells and the intensity of the signal increased very rapidly (Fig. 2C). Concomitantly to the neural tube closure, β -galactosidase expression became progressively restricted to a sharp stripe at the level of the otic vesicle (Fig. 2D) and to a broader and more posterior domain. Half a day later, only cells of the roof plate and the floor plate are still expressing *vHNF1/lacZ* in this part of the neural tube as can be seen on sections (data not shown).

vHNF1/lacZ expression starts in the gut epithelium during its invagination. A sharp anterior boundary was observed at E9.5 at the level of the pancreas and liver primordia (Fig. 2D). The expression extended posteriorly to the tip of the gut.

The expression pattern described above was confirmed by




Fig. 2. Expression pattern of *vHNF1* at early stages of development as revealed by β -galactosidase activity. (A) E7.5 embryo; (B) E8 embryo; (C) E8.5 embryo; (D) E9.5 embryo. All embryos are heterozygous. Orientation: (A) ectoplacental cone on top and embryo on bottom; (B-D) head on top, back on the left. (A) Arrowheads: boundaries between embryonic and extraembryonic visceral endoderms; (B,C) black arrows, remainder of the yolk sac; white arrowheads, neural folds; white arrows, gut. (D) Star indicates the position of the otic vesicle; white arrowhead, roof plate staining; white arrow, anterior boundary of the gut staining. Scale bars: 250 μ m on A and D, 500 μ m on B and C.

X-gal staining on chimeric embryos obtained from a second independent *vHNF1/lacZ* clone.

***vHNF1* inactivation results in early embryonic lethality**

vHNF1^{+/-} mice are normal and fertile. They were crossed together and the resulting litters were genotyped by PCR. No *vHNF1*^{-/-} mice were born (Table 1). Therefore, litters from heterozygotes were collected at different gestation times to investigate the timing of the embryonic lethality. At E6.5, *vHNF1*^{-/-} embryos were normal and not distinguishable from their littermates (data not shown). Live *vHNF1*^{-/-} embryos were not found after E7.5 (Table 1). In the litters collected later than E8.5, only resorbed tissue could be genotyped as homozygous. At E7.5, *vHNF1*^{-/-} embryos were identified by PCR (Fig. 1C). They all presented a severe growth retardation when compared to their *vHNF1*^{+/+} and *vHNF1*^{+/-} littermates (Fig. 3). Morphological analysis of *vHNF1*^{-/-} embryos revealed in all cases a disorganization of the embryonic and extraembryonic tissues, despite some variations in the severity of the phenotype (Fig. 4). On sections of E7.5 heterozygous embryos, two territories in the visceral endoderm are morphologically distinct: the extraembryonic visceral

endoderm composed of columnar epithelial cells is stained by X-gal, whereas the epithelial layer surrounding the gastrulating embryo is squamous and negative for this staining (Fig. 4A). In contrast, the *vHNF1*^{-/-} embryos present a lack of development of the embryonic tissues (Fig. 4B,C). Most of them were highly degenerated and disorganized. They comprised a mass of undifferentiated cells, some of them positive for the X-gal staining, and were surrounded by trophoblastic tissues (Fig. 4C). Some embryos were more developed and underwent cavitation. However, the embryonic tissue was reduced and presented no sign of gastrulation. No clear regionalization of the visceral endoderm was observed in these embryos (Fig. 4B). Indeed, the visceral endoderm was uniformly composed of cuboidal epithelial cells, some expressing the *lacZ* gene even at the distal pole of the embryos (Fig. 4B). This morphology was reminiscent of visceral endoderm from a E6.5 embryo. The defects in distinct visceral endoderm specialization led us to investigate the role of *vHNF1* in this differentiation pathway.

Table 1. Viability of *vHNF1*^{-/-} embryos

Age	Number of litters	Genotype of embryos			N.D.	Number of pups
		+/+	+/-	-/-		
Week 2	12	35	65	0		100
E10.5	4	5	15	(1)	2	23
E8.5	7	11	25	(10)	11	57
E7.5	6	14	23	7	10	54

PCR analysis of the litters from *vHNF1*^{+/-} intercrosses.

DNA was isolated from embryos between E7.5 and E10.5 and from 2-week-old mice (Week 2). It was analyzed by PCR using the three primers featured on Fig. 1.

In parentheses are indicated the number of resorbed tissues genotyped as *vHNF1*^{-/-}.

N.D. (not determined) marks the number of embryos or resorbed tissues that could not be genotyped.




Fig. 3. A representative litter obtained from a *vHNF1*^{+/-} intercross, stained for *vHNF1/lacZ* activity. *vHNF1*^{-/-} embryos present a severe growth retardation and disorganized tissues at E7.5. Genotypes obtained by PCR are indicated next to the embryos. *, broken *vHNF1*^{+/-} embryo. Scale bar, 300 μ m.




Fig. 4. Sections of (A) vHNF1^{+/-} and (B,C) vHNF1^{-/-} E7.5 embryos after X-gal staining. Black arrows, ectoplacental cone; black arrowheads, embryonic/extraembryonic boundary of the visceral endoderm; white arrows in B, cuboidal visceral endoderm; white arrows in C, vHNF1/*lacZ*-positive cells; t, trophoblastic tissue. Longitudinal sections. Scale bars, 120 μm.

vHNF1 inactivation affects in vitro differentiation of ES cells into visceral endoderm

ES cells differentiation provides a good model to study visceral endoderm formation. Indeed, when cultured in suspension, clumps of ES cells form complex vesicles called embryoid bodies (EB). These vesicles contain both ectoderm and visceral endoderm. Their development in culture can be considered as equivalent to the early stages of embryogenesis (Keller, 1995). As one of the genes activated during this process is vHNF1 (Abe et al., 1996), we have compared the formation of EB generated from vHNF1^{+/+}, vHNF1^{+/-} and vHNF1^{-/-} ES cells. Cells were cultured in suspension for 14 days, a time by which visceral endoderm differentiation should be completed (Abe et al., 1996). The resulting EB were analyzed for both their morphology and the expression of various differentiation markers (Fig. 5). Experiments were repeated several times with independently isolated clones. Embryoid bodies obtained from vHNF1^{-/-} ES cells were always smaller than the control (+/-) EB (Fig. 5A). They never formed expanded vesicles containing a tissue equivalent to yolk sac. However, both vHNF1^{+/-} and vHNF1^{-/-} EB contain endodermal cells as detected by SJA lectin staining (Sophora japonica agglutinin, see Soudais et al., 1995) (data not shown).




Fig. 5. vHNF1 inactivation alters the ES cell differentiation into visceral endoderm. (A) 14-day-old embryoid bodies from either vHNF1^{-/-} or vHNF1^{+/-} cells. Scale bar, 300 μm. (B) RT-PCR analysis of endodermal markers of total RNAs from vHNF1^{-/-}, vHNF1^{+/-} or vHNF1^{+/+} embryoid bodies. Genotypes are indicated on top. 0, PCR control – no DNA. HPRT primers were used as a normalizing standard for the RT samples (HPRT⁺) and as a reverse transcription control (HPRT⁻, control reaction without reverse transcriptase). TFN, Transferrin; TTR, Transthyretin; AFP, Alpha Fetoprotein; ApoAI, Apolipoprotein AI; ApoAIV, Apolipoprotein AIV; BMP-6, Bone Morphogenetic Protein 6.

To examine the expression of endodermal markers, total RNAs were prepared from EB and RT-PCR analysis performed. Primers for the HPRT gene were used as controls (Fig. 5B). Early endoderm markers like GATA-4 (Fig. 5B) and GATA-6 (not shown) were expressed in vHNF1^{-/-} EB as in controls, despite a decrease in their level. On the contrary, late endodermal markers, such as *alphafetoprotein* (AFP), *transthyretin* (TTR) and *apolipoproteins*, were not expressed in the vHNF1^{-/-} EB. Remarkably, expression of the transcription factors HNF4α and HNF1 was strongly reduced, in contrast to the expression levels of the HNF3 genes, which were approximately equivalent to those of the vHNF1^{+/+} and vHNF1^{+/-} samples. The expression of signaling molecules that are expressed at the equivalent stage of embryonic development was also examined: no variations were observed for the two secreted factors BMP-6 (Fig. 5B) and BMP-4 (not shown).

These results suggest a block of the differentiation of the vHNF1^{-/-} ES cells into visceral endoderm. GATA factor expression and positive SJA staining of vHNF1^{-/-} EB demonstrate that the first steps of visceral endoderm differentiation still occur in the absence of vHNF1. However, this process is not completed as shown by the lack of expression of the late markers.

vHNF1 is essential for in vivo differentiation of the visceral endoderm

We next analyzed the expression of several endodermal




Fig. 6. Visceral endoderm differentiation is impaired in $vHNF1^{-/-}$ embryos. RT-PCR analysis of endodermal markers was performed on E7.5 $vHNF1^{-/-}$ and $vHNF1^{+/+}$ embryos. Material for $vHNF1^{-/-}$ embryos was pooled from a set of four embryos, $vHNF1^{+/+}$ were analyzed individually. The controls and the procedure are the same as in Fig. 5.

markers in E7.5 $vHNF1^{+/+}$ and $vHNF1^{-/-}$ embryos. Due to the small size of mutant embryos, genotyping was achieved on DNA from ectoplacental cone and RT-PCR was then performed either on a wild-type embryo or a pool of four mutant embryos. As previously, *HPRT* primers were used to normalize the samples. *vHNF1* inactivation in vivo results in effects similar to those observed in vitro: both *GATA-4* and *GATA-6* are expressed despite a slight decrease in their levels; in contrast, the levels of the *HNF4* and *transthyretin* genes are strongly reduced (Fig. 6).

Both in vitro and in vivo analysis revealed a requirement for *vHNF1* expression during visceral endoderm differentiation. This defect in one of the first extraembryonic tissues could explain the growth retardation and abnormal morphology of the mutant embryos. To address this issue, rescue experiments were attempted by providing wild-type extraembryonic annexes to mutant embryos. This should also permit the examination of additional *vHNF1* functions during embryonic development.

Tetraploid aggregation partially rescues *vHNF1* inactivation

Wild-type visceral endoderm can be provided to a mutant embryo by aggregating tetraploid morulae with a clump of ES cells. The resulting embryos are composed of two distinct compartments: the extraembryonic tissue is wild type and tetraploid, whereas the embryo itself derives from the diploid ES cells (Rossant and Spence, 1998). This experiment permits to investigate the effect of wild-type visceral endoderm on the viability of $vHNF1^{-/-}$ embryos. In addition, this should allow us to examine the effect of *vHNF1* inactivation in other tissues.

Aggregation experiments were performed with two independent $vHNF1^{-/-}$ ES clones, as well as with $vHNF1^{+/+}$ cells as a control. Embryos obtained with control cells were perfectly normal at E9.5 (not shown). Embryos resulting from the aggregation of wild-type tetraploid morula with $vHNF1^{-/-}$ cells (denoted $+/+::-/-$) were viable up to E10. At E8.5 these embryos were not distinguishable from $vHNF1^{+/+}$ embryos in terms of both their size and morphology. At E9.5, the rescued embryos still presented a normal size for this embryonic stage (Fig. 7A,B). These results demonstrate that wild-type visceral endoderm can efficiently rescue both the early lethality and the growth retardation due to the *vHNF1* mutation. It also showed that $vHNF1^{-/-}$ epiblast can normally develop after implantation and undergo gastrulation. However, morphological examination of the E9.5 $+/+::-/-$ embryos revealed some interesting abnormalities. The embryos did not turn, they were rather short

and their general appearance corresponded more to a E8.5 embryo (Fig. 7C). The neural tube remained open and exhibited expanded tissue on both neural folds; often, this overgrowth was associated with several oedemas on the neural folds and on the forebrain vesicles. Despite being wild type, the endodermal layer of the yolk sac was oversized and ruffled (not shown). An enlarged allantois and a distended pericardium were also observed in a majority of $+/+::-/-$ embryos (Fig. 7C).

X-gal staining of the $+/+::-/-$ embryos revealed an additional abnormality. As it was reported for the heterozygotes, β -galactosidase activity was detected in both the gut and the neural folds. Still, *vHNF1/lacZ* expression in the neural tissue was not restricted, either laterally and ventrally, as it should be at E9.5. This expression was even expanded (Fig. 7B,C). In addition, the staining remained uniform at E9.5 all along the neural tube as if no anteroposterior regionalization of *vHNF1/lacZ* expression was accomplished in the absence of *vHNF1* protein.

$vHNF1^{-/-}$ cells contribute differentially to chimeric embryo tissues

The morphological abnormalities exhibited by the rescued $vHNF1^{-/-}$ embryos suggested a requirement for the expression of this gene during morphogenesis, in addition to its essential function in visceral endoderm. To determine if the activity of the gene was cell-autonomously required or not in embryonic tissues, the behavior of the $vHNF1^{-/-}$ ES cells in a wild-type environment was examined by injecting 7 to 8 cells into wild-type blastocysts. The resulting chimeras were collected between E9.5 and E 11.5. They all presented a normal morphology (Fig. 7D-F). After X-gal staining, *vHNF1/lacZ*-positive cells could be detected in both the neural tube and the mesonephros (Fig. 7E,F). The neural tube was closed and the mesonephros was normal.

On the contrary, $vHNF1^{-/-}$ blue cells were observed in the gut in only one out of eight chimeric embryos. In this case, the stained cells were clustered in spots and some were detected at a position more anterior than expected (not shown). In contrast, in all three chimeras obtained by the injection of $vHNF1^{+/+}$ ES cells, we observed a strong and uniform staining in the gut, similar to that observed for F1 heterozygous embryos. So, these results suggest that $vHNF1^{-/-}$ cells present a strongly reduced capacity to contribute to gut formation.

DISCUSSION

Transcription factors interact in a functional network to specify cell differentiation. We have been interested in analyzing the function of the *vHNF1* homeoprotein by inactivating the corresponding gene in the mouse. From in vivo and in vitro studies, we demonstrate that *vHNF1* is required for visceral endoderm differentiation as well as for morphogenesis and we identify potential target genes of *vHNF1* in the visceral endoderm.

Dynamic expression of *vHNF1* during early embryogenesis

The introduction of an NLS-*lacZ* reporter gene at the initiation codon of *vHNF1* has provided us with an easy visualization of

vHNF1 expression by X-gal staining. β -galactosidase activity was first detected in the extraembryonic part of the visceral endoderm at E7.5. At the 2-somite stage, it was also detected in the embryo proper, first in the neural folds then in the invaginating gut. This result corroborates previous data obtained by in situ hybridization, where vHNF1 mRNA was detected in the visceral endoderm at E6.5 (Cereghini et al., 1992). An interesting feature of vHNF1 expression at E9.5 is the sharp anterior boundary of the gut expression domain. The regionalization of the gut is established by the spatially restricted expression of several genes in the mesoderm during epithelium-mesenchyme interactions (Roberts et al., 1998). The mechanisms of endoderm patterning are less understood, but they certainly imply the differential expression of regulatory genes. Considering this model, the sharp anterior boundary observed for vHNF1/lacZ expression suggests that vHNF1 could contribute to the determination of territories in the gut in combination with other transcription factors. In particular it could play a role in the specification of the liver and pancreatic primordia. This issue will be the subject of future studies using the conditional inactivation of the gene.

The expression pattern of vHNF1, as depicted by X-gal staining, and the phenotype of the tetraploid-rescued mutant embryos strongly argue for vHNF1 playing a role in neural tube formation. This function was unexpected. Indeed, the expression of vHNF1 in the neural tube of the mouse has not been described previously except as a transient signal in the neuroepithelium of the murine embryo. However, it should be recalled that vHNF1 expression has been reported in the neural tube of two other vertebrates, *Xenopus* (Demartis et al., 1994) and zebrafish (Thisse and Thisse, 1998).

vHNF1 expression is critical for visceral endoderm differentiation

As mentioned above, vHNF1 is specifically expressed in the extraembryonic territory of the visceral endoderm at E7.5. Combined studies of the mutant embryos and ES cell differentiation have shown that visceral endoderm formation is strongly impaired in the absence of vHNF1. These results are the first evidence for an essential role for vHNF1 in controlling this process. In addition, wild-type visceral endoderm provided by tetraploid aggregation was shown to rescue the lethality and growth defect of the mutant embryos. This approach allowed us to bypass the lethality up to E9.5.

The differentiation of the visceral endoderm is a crucial event for the mammalian embryogenesis. The discovery of the organizer properties of this tissue has recently awakened much interest (Beddington and Robertson, 1998). But the visceral endoderm plays also a major role during the periimplantation development in supporting embryonic metabolism. The knowledge concerning these functions remains fragmentary. Still, it has been established that it is a multistep process in which the HNF4 factor seems to play an important role. It is known that numerous secreted proteins are produced by this tissue such as serum proteins (AFP, TTR) or apolipoproteins (Meehan et al., 1984) and that their production is dependent upon HNF4 expression (Duncan et al., 1997).

Several mutations like Smad4/Dpc4 (Sirard et al., 1998), evx1 (Spyropoulos and Capecchi, 1994), Nf2 (McClatchey et al., 1997) or GATA-6 (Morrisey et al., 1998) were characterized that result in the failure of development due to a

defect of visceral endoderm functions. A strong similarity exists between the Smad4/Dpc4, GATA-6 and vHNF1 phenotypes. These mutations lead to embryonic lethality at E7-7.5 and to a severe growth retardation. A disorganization of the visceral endoderm has been described and the lethality occurs shortly after the initiation of their expression in this tissue. Also, the mutation of each of these three genes results in a lack of HNF4 expression.

Further insights into the relationship between regulators during visceral endoderm differentiation have come from the study of ES cells. When cultured in suspension, they develop following a program analogous to that occurring in the embryo. A regulatory network involved in this process has been recently proposed by Duncan and co-workers (1998). It organizes some of the HNF genes hierarchically: for instance, HNF3 β regulates both HNF3 α and HNF4 α expression in the visceral endoderm. Morrisey and co-workers (1998) have reported similar observations concerning GATA-6 that would directly regulate HNF4 expression. Our studies allow us to place vHNF1 in this pathway. Although some endoderm is formed from the ES cells in the absence of vHNF1 as demonstrated by the expression of the GATA factors and the positive staining for SJA lectin, their differentiation remains incomplete as they lack late endodermal markers. The genes examined in the embryos were affected both in vitro and in vivo in the absence of vHNF1. These results are the first evidence for an important role of vHNF1 during visceral endoderm differentiation.

A model for visceral endoderm differentiation is shown in Fig. 8. GATA-6 and GATA-4 are required for the onset of visceral endoderm differentiation right after the specification of this tissue, but the completion of this process requires the additional expression of both HNF3 and vHNF1. As the expressions of the HNF3 and GATA genes are not affected by vHNF1 inactivation, we can propose, in accordance with the results of Duncan (1998) and Morrisey (1998), that HNF3 β , GATA-6 and vHNF1 are acting through separate pathways converging on HNF4 α (Fig. 8).

Furthermore, as the promoters for serum proteins like TTR contain HNF4- and vHNF1/HNF1-binding sites (Xanthopoulos et al., 1991), both these transcription factors could be essential for their expression. The combined action of vHNF1 and HNF4 would switch on the expression of the target genes. It is worth recalling that vHNF1 is expressed in the absence of HNF4 α (Duncan et al., 1997). As a binding site for vHNF1 is present in the HNF4 α promoter (Taraviras et al., 1994), vHNF1 could lie just upstream of HNF4. Still, the early lethality observed at E7.5 for the vHNF1^{-/-} embryos contrasts with the phenotype of the HNF4^{-/-} embryos that survive up to E10.5 and display a normal morphology (Chen et al., 1994). This confirms our conclusions that vHNF1 plays a role in the regionalization and structural specification of the extraembryonic visceral endoderm.

In addition, in embryoid bodies, HNF1 expression is not affected by HNF4 inactivation (Duncan et al., 1997), but is downregulated in the absence of vHNF1. This result suggests a direct requirement for vHNF1 in the visceral endoderm to activate HNF1 expression. This hierarchy of transcription factors differs from the network reported for the hepatic model. Indeed, in hepatocytes, HNF1 expression has been described as strictly dependent upon that of HNF4, whereas both HNF1 and HNF4 seem to be independent of vHNF1 (reviewed in




Fig. 7. *vHNF1* might play multiple roles during the embryonic morphogenesis. (A-C) Wild-type visceral endoderm is able to rescue partially the *vHNF1* mutation. (A) E9.5 *vHNF1*^{+/-} embryo. (B,C) E9.5 embryos resulting from aggregations of *vHNF1*^{+/+} tetraploid embryos with *vHNF1*^{-/-} ES cells. All three embryos were stained for β -galactosidase activity. Note the expanded neural tube staining for *vHNF1/lacZ* in the rescued embryos (arrowheads, compare A to B and C). The wild-type yolk sacs (white arrows) are negative for β -galactosidase activity. Black star in C, gut. Orientation: (A,B) dorsal view, anterior on top; (C) lateral view, anterior left. Scale bar: 600 μ m. (D-F) *vHNF1*^{-/-} cells can contribute to normal neural tube and mesonephros. Chimeras were obtained after injection of *vHNF1*^{-/-} cells into wild-type blastocysts; they were stained for β -galactosidase activity. (D) E9.5 chimera; (E) E10.5 chimera; (F) E11.5 chimera after abdominal wall removal. No blue cells were detected in the gut of these embryos. Black arrowheads, mesonephros; white and black arrows, neural tube; black square, hindbrain staining. Scale bars: (D) 200 μ m; (E) 400 μ m; (F) 100 μ m.

Tronche et al., 1994). Hence, cell differentiation would not only involve the combined expression of transcription factors, but these factors would also be hierarchically organized into different networks depending on the differentiation program.

Multiple developmental requirements for *vHNF1*

The tetraploid aggregation allowed us to provide mutant embryos with wild-type extraembryonic annexes. We were able to show that the wild-type visceral endoderm can rescue both the early lethality and the growth defect of the *vHNF1*^{-/-} epiblast. Furthermore, it leads to a normal development including gastrulation, up to E9. We can conclude that the embryonic lethality of the *vHNF1*^{-/-} embryos at E7.5 is not caused by any epiblast defect, but is the consequence of impaired visceral endoderm functions. Normal development was observed up to E9, then morphological abnormalities appeared at the two sites of *vHNF1* expression, the neural tube and the gut. The neural tube remained completely open and the gut failed to close and/or invaginate. Additionally, the embryos


failed to turn and the short appearance of their bodies suggested a possible truncation of the posterior part. These embryos looked more like E8.5 than E9.5/10 stages. This phenotype suggested a block occurring during morphogenesis.

The *vHNF1*^{+/+::-/-} embryos present striking similarities in their phenotypes with four knock-outs: *Fgf receptor1* (Deng et al., 1994; Yamaguchi et al., 1994), *α 5 integrin* (Yang et al., 1993), *N-cadherin* (Radice et al., 1997) and *fibronectin* (George et al., 1993). These mutations result, to different extents, in the association of the following defects: kinking of the neural tube, ruffling of the visceral endoderm, swelling of the pericardium and overgrowth of the allantois. These similarities suggest that signaling pathways involving secreted factors or cell adhesion might be impaired in the *vHNF1*^{+/+::-/-} embryos.

To study the effect of the *vHNF1* mutation in a wild-type environment, we produced chimeric embryos by injecting ES cells into wild-type blastocysts. Monitoring the presence of *vHNF1*^{-/-} cells by X-gal staining of the tissues normally expressing *vHNF1* should allow us to determine if the effect of the mutation is cell-autonomous or not. The resulting chimeras displayed a normal morphology at E9.5-11.5. Hence, *vHNF1*^{-/-} cells could contribute to both the neural tube and the mesonephros in a wild-type context. In contrast, a majority of the chimeras lacked blue cells in the gut, indicating a possible exclusion of the *vHNF1*^{-/-} cells from the gut or a lower ability to colonize this tissue. This analysis suggests that *vHNF1* could act through different pathways depending on the tissues where it is expressed. The absence of *vHNF1* could be deleterious for cells in the gut; but not in the neural tube where the *vHNF1* mutation would be complemented by the surrounding wild-type cells.

In conclusion, the null mutation of the murine *vHNF1* gene has allowed us to characterize three early and specific functions of *vHNF1* during embryogenesis. We have shown that *vHNF1* expression is necessary for visceral endoderm differentiation and we present evidence for the involvement of *vHNF1* during morphogenesis, both in the neural tube and in the gut.

The embryonic lethality has hampered the study of the later



¹: Morrisey et al., 1998; ²: Soudais et al., 1995; ³: Duncan et al., 1998; ⁴: Duncan et al., 1997

Fig. 8. A regulatory network of transcription factors involved in visceral endoderm differentiation.

functions of the gene in the polarized epithelia where vHNF1 is specifically expressed. To analyze these possible functions during late embryogenesis or in adult tissues, we will pursue the conditional inactivation of the gene using the Cre/lox system. This study will be of a particular interest in the pancreas as two human vHNF1 mutations have been recently characterized that are associated with a type II diabetes (Horikawa et al., 1997; Nishigori et al., 1998).

We thank C. Kress for providing the CK35 ES cells, P. Brulet and H. Le Mouellic for the lacZ cassette and the mouse genomic library, J. C. Benichou and C. Cimper for helpful advice in sectioning, A. Doyen and B. Bourachot for technical advice, M. Pontoglio and D. Lallemand for help in the production of the vHNF1 antiserum, F. Tronche for the vector containing the loxP sites. We are grateful to J. Weitzman, A. Yeivin and A. El Amraoui for critical reading of the manuscript. C. C. was an Allocataire Moniteur Normalien at the Paris XI University. This work was supported by grants from the EEC Biotechnology program, the ARC and the LNFCC.

REFERENCES

- Abe, K., Niwa, H., Iwase, K., Takiguchi, M., Mori, M., Abe, S. I., Abe, K. and Yamamura, K. I. (1996). Endoderm-specific gene expression in embryonic stem cells differentiated to embryoid bodies. *Exp. Cell Res.* **229**, 27-34.
- Ang, S. L. and Rossant, J. (1994). HNF-3 beta is essential for node and notochord formation in mouse development. *Cell* **78**, 561-574.
- Beddington, R. S. and Robertson, E. J. (1998). Anterior patterning in mouse. *Trends Genet.* **14**, 277-284.
- Blumenfeld, M., Maury, M., Chouard, T., Yaniv, M. and Condamine, H. (1991). Hepatic nuclear factor 1 (HNF1) shows a wider distribution than products of its known target genes in developing mouse. *Development* **113**, 589-599.
- Cereghini, S., Ott, M. O., Power, S. and Maury, M. (1992). Expression patterns of vHNF1 and HNF1 homeoproteins in early postimplantation embryos suggest distinct and sequential developmental roles. *Development* **116**, 783-797.
- Chen, W. S., Manova, K., Weinstein, D. C., Duncan, S. A., Plump, A. S., Prezioso, V. R., Bachvarova, R. F. and Darnell, J. E. J. (1994). Disruption of the HNF-4 gene, expressed in visceral endoderm, leads to cell death in embryonic ectoderm and impaired gastrulation of mouse embryos. *Genes Dev.* **8**, 2466-2477.
- Copp, A. J. (1995). Death before birth: clues from gene knockouts and mutations. *Trends Genet.* **11**, 87-93.
- Demartis, A., Maffei, M., Vignali, R., Barsacchi, G. and De Simone, V. (1994). Cloning and developmental expression of LFB3/HNF1 beta transcription factor in *Xenopus laevis*. *Mech. Dev.* **47**, 19-28.
- Deng, C. X., Wynshaw-Boris, A., Shen, M. M., Daugherty, C., Ornitz, D. M. and Leder, P. (1994). Murine FGFR-1 is required for early postimplantation growth and axial organization. *Genes Dev.* **8**, 3045-3057.
- Dufort, D., Schwartz, L., Harpal, K. and Rossant, J. (1998). The transcription factor HNF3beta is required in visceral endoderm for normal primitive streak morphogenesis. *Development* **125**, 3015-3025.
- Duncan, S. A., Manova, K., Chen, W. S., Hoodless, P., Weinstein, D. C., Bachvarova, R. F. and Darnell, J. E. J. (1994). Expression of transcription factor HNF-4 in the extraembryonic endoderm, gut, and nephrogenic tissue of the developing mouse embryo: HNF-4 is a marker for primary endoderm in the implanting blastocyst. *Proc. Natl. Acad. Sci. USA* **91**, 7598-7602.
- Duncan, S. A., Nagy, A. and Chan, W. (1997). Murine gastrulation requires HNF-4 regulated gene expression in the visceral endoderm: tetraploid rescue of Hnf-4(-/-) embryos. *Development* **124**, 279-287.
- Duncan, S. A., Navas, M. A., Dufort, D., Rossant, J. and Stoffel, M. (1998). Regulation of a transcription factor network required for differentiation and metabolism. *Science* **281**, 692-695.
- George, E. L., Georges-Labouesse, E. N., Patel-King, R. S., Rayburn, H. and Hynes, R. O. (1993). Defects in mesoderm, neural tube and vascular development in mouse embryos lacking fibronectin. *Development* **119**, 1079-1091.
- Gu, H., Zhou, Y. R. and Rajewsky, K. (1993). Independent control of immunoglobulin switch at individual switch regions evidenced through the Cre-loxP-mediated gene targeting. *Cell* **73**, 1155-1164.
- Guillemot, F., Nagy, A., Auerbach, A., Rossant, J. and Joyner, A. L. (1994). Essential role of Mash-2 in extraembryonic development. *Nature* **371**, 333-336.
- Horikawa, Y., Iwasaki, N., Hara, M., Furuta, H., Hinokio, Y., Cockburn, B. N., Lindner, T., Yamagata, K., Ogata, M., Tomonaga, O. et al. (1997). Mutation in hepatocyte nuclear factor-1 beta gene (TCF2) associated with MODY. *Nature Genet.* **17**, 384-385.
- Keller, G. M. (1995). In vitro differentiation of embryonic stem cells. *Curr. Opin. Cell Biol.* **7**, 862-869.
- Koopman, P. (1993). Analysis of gene expression by reverse transcriptase-polymerase chain reaction. In *Essential Developmental Biology: a Practical Approach*, (ed. C. D. Stern and P. W. H. Holland), pp. 231-242. Oxford: IRL Press Limited.
- Kress, C., Vandormael-Pournin, S., Baldacci, P., Cohen-Tannoudji, M. and Babinet, C. (1998). Nonpermissiveness for mouse embryonic stem (ES) cell derivation circumvented by a single backcross to 129/Sv strain: establishment of ES cell lines bearing the *Om*^d conditional lethal mutation. *Mammalian Genome* **9**, 998-1001.
- McClatchey, A. I., Saotome, I., Ramesh, V., Gusella, J. F. and Jacks, T. (1997). The Nf2 tumor suppressor gene product is essential for extraembryonic development immediately prior to gastrulation. *Genes Dev.* **11**, 1253-1265.
- Meehan, R. R., Barlow, D. P., Hill, R. E., Hogan, B. L. M. and Hastie, N. D. (1984). Pattern of serum protein gene expression in mouse visceral yolk sac and foetal liver. *EMBO J.* **3**, 1881-1885.
- Monaghan, A. P., Kaestner, K. H., Grau, E. and Schutz, G. (1993). Postimplantation expression patterns indicate a role for the mouse forkhead/HNF-3 alpha, beta and gamma genes in determination of the definitive endoderm, chordamesoderm and neuroectoderm. *Development* **119**, 567-578.
- Morrissey, E. E., Tang, Z., Sigrist, K., Lu, M. M., Jiang, F., Ip, H. S. and Parmacek, M. S. (1998). GATA6 regulates HNF4 and is required for differentiation of visceral endoderm in the mouse embryo. *Genes Dev.* **12**, 3579-3590.
- Mortensen, R. M., Conner, D. A., Chao, S., Geisterfer-Lowrance, A. A. and Seidman, J. G. (1992). Production of homozygous mutant ES cells with a single targeting construct. *Mol. Cell. Biol.* **12**, 2391-2395.
- Nagy, A., Rossant, J., Nagy, R., Abramow-Newerly, W. and Roder, J. C. (1993). Derivation of completely cell culture-derived mice from early-passage embryonic stem cells. *Proc. Natl. Acad. Sci. USA* **90**, 8424-8428.
- Nishigori, H., Yamada, S., Kohama, T., Tomura, H., Sho, K., Horikawa, Y., Bell, G. I., Takeuchi, T. and Takeda, J. (1998). Frameshift mutation, A263fsinsGG, in the hepatocyte nuclear factor-1beta gene associated with diabetes and renal dysfunction. *Diabetes* **47**, 1354-1355.
- Ott, M. O., Rey-Campos, J., Cereghini, S. and Yaniv, M. (1991). vHNF1 is expressed in epithelial cells of distinct embryonic origin during development and precedes HNF1 expression. *Mech. Dev.* **36**, 47-58.
- Pontoglio, M., Barra, J., Hadchouel, M., Doyen, A., Kress, C., Bach, J. P., Babinet, C. and Yaniv, M. (1996). Hepatocyte nuclear factor 1 inactivation results in hepatic dysfunction, phenylketonuria, and renal Fanconi syndrome. *Cell* **84**, 575-585.
- Pontoglio, M., Sreenan, S., Roe, M., Pugh, W., Ostrega, D., Doyen, A., Pick, A. J., Baldwin, A., Velho, G., Froguel, P. et al. (1998). Defective insulin secretion in hepatocyte nuclear factor 1alpha-deficient mice. *J. Clin. Invest.* **101**, 2215-2222.
- Radice, G. L., Rayburn, H., Matsunami, H., Knudsen, K. A., Takeichi, M. and Hynes, R. O. (1997). Developmental defects in mouse embryos lacking N-cadherin. *Dev. Biol.* **181**, 64-78.
- Roberts, D. J., Smith, D. M., Goff, D. J. and Tabin, C. J. (1998). Epithelial-mesenchymal signaling during the regionalization of the chick gut. *Development* **125**, 2791-2801.
- Rossant, J. and Spence, A. (1998). Chimeras and mosaics in mouse mutant analysis. *Trends Genet.* **14**, 358-363.
- Rudnicki, M. A. and McBurney, M. W. (1987). Cell culture methods and induction of differentiation of embryonal carcinoma cell lines. In *Teratocarcinomas and Embryonic Stem cells: a Practical Approach*, (ed. E. J. Robertson), pp. 19-49. Oxford: IRL Press Limited.
- Sirard, C., de la Pompa, J. L., Elia, A., Itie, A., Mirtsos, C., Cheung, A., Hahn, S., Wakeham, A., Schwartz, L., Kern, S. E. et al. (1998). The tumor suppressor gene Smad4/Dpc4 is required for gastrulation and later for anterior development of the mouse embryo. *Genes Dev.* **12**, 107-119.

- Soudais, C., Bielinska, M., Heikinheimo, M., MacArthur, C. A., Narita, N., Saffitz, J. E., Simon, M. C., Leiden, J. M. and Wilson, D. B.** (1995). Targeted mutagenesis of the transcription factor GATA-4 gene in mouse embryonic stem cells disrupts visceral endoderm differentiation in vitro. *Development* **121**, 3877-3888.
- Spyropoulos, D. D. and Capecchi, M. R.** (1994). Targeted disruption of the even-skipped gene, *evx1*, causes early postimplantation lethality of the mouse conceptus. *Genes Dev.* **8**, 1949-1961.
- Taraviras, S., Monaghan, A. P., Schutz, G. and Kelsey, G.** (1994). Characterization of the mouse HNF-4 gene and its expression during mouse embryogenesis. *Mech. Dev.* **48**, 67-79.
- Thisse, C. and Thisse, B.** (1998). Antivin, a novel and divergent member of the TGFbeta superfamily, negatively regulates mesoderm induction. *Development* **126**, 229-240.
- Tronche, F. and Yaniv, M.** (1992). HNF1, a homeodomain member of the hepatic transcription regulatory network. *BioEssays* **14**, 579-587.
- Tronche, F., Bach, I., Chouard, T., David-Wattine, B., Pontoglio, M., Ringeisen, F., Sourdive, D., Thépot, D. and Yaniv, M.** (1994). Hepatocyte Nuclear factor 1 (HNF1) and liver gene expression. In *Liver Gene Expression*, (ed. F. Tronche and M. Yaniv), pp. 155-181. Austin: R. G. Landes Company.
- Tybulewicz, V. L., Crawford, C. E., Jackson, P. K., Bronson, R. T. and Mulligan, R. C.** (1991). Neonatal lethality and lymphopenia in mice with a homozygous disruption of the *c-abl* proto-oncogene. *Cell* **65**, 1153-1163.
- Weinstein, D. C., Ruiz i Altaba, A., Chen, W. S., Hoodless, P., Prezioso, V. R., Jessell, T. M. and Darnell, J. E. J.** (1994). The winged-helix transcription factor HNF-3 beta is required for notochord development in the mouse embryo. *Cell* **78**, 575-588.
- Wiles, M. V.** (1993). Embryonic stem cells differentiation in vitro. In *Methods in Enzymology*, vol. 225, pp 900-918. New York: Academic Press Inc.
- Xanthopoulos, K. G., Prezioso, V. R., Chen, W. S., Sladek, F. M., Cortese, R. and Darnell, J. E. J.** (1991). The different tissue transcription patterns of genes for HNF-1, C/EBP, HNF-3, and HNF-4, protein factors that govern liver-specific transcription. *Proc. Natl. Acad. Sci. USA* **88**, 3807-3811.
- Yamaguchi, T. P., Harpal, K., Henkemeyer, M. and Rossant, J.** (1994). *fgfr-1* is required for embryonic growth and mesodermal patterning during mouse gastrulation. *Genes Dev.* **8**, 3032-3044.
- Yamamoto, H., Flannery, M. L., Kupriyanov, S., Pearce, J., McKercher, S. R., Henkel, G. W., Maki, R. A., Werb, Z. and Oshima, R. G.** (1998). Defective trophoblast function in mice with a targeted mutation of *Ets2*. *Genes Dev.* **12**, 1315-1326.
- Yang, J. T., Rayburn, H. and Hynes, R. O.** (1993). Embryonic mesodermal defects in $\alpha 5$ integrin-deficient mice. *Development* **119**, 1093-1105.