



HAL
open science

Sélection pour abaisser le potentiel glycolytique du muscle chez le porc Large White 1. Réponses directes

Pascale Le Roy, Catherine Larzul, Jean-Marcel Gogué, André Talmant, Gabriel Monin, Pierre Sellier

► To cite this version:

Pascale Le Roy, Catherine Larzul, Jean-Marcel Gogué, André Talmant, Gabriel Monin, et al.. Sélection pour abaisser le potentiel glycolytique du muscle chez le porc Large White 1. Réponses directes. *Genetics Selection Evolution, BioMed Central*, 1998, 30 (5), pp.469-480. hal-02692371

HAL Id: hal-02692371

<https://hal.inrae.fr/hal-02692371>

Submitted on 1 Jun 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Selection for reduced muscle glycolytic potential in Large White pigs.

I. Direct responses

Pascale Le Roy^a, Catherine Larzul^{a*}, Jean Gogué^b,
André Talmant^c, Gabriel Monin^c, Pierre Sellier^a

^a Station de génétique quantitative et appliquée, Institut national de la recherche agronomique, 78352 Jouy-en-Josas cedex, France

^b Institut national de la recherche agronomique, domaine de Galle, 18520 Avord, France

^c Station de recherches sur la viande, Institut national de la recherche agronomique, Theix, 63122 Saint-Genès-Champanelle, France

(Received 3 March 1998; accepted 10 September 1998)

Abstract – A six-generation selection experiment comprising a selected (S) and a control (C) line has been conducted with the objective of decreasing muscle glycolytic potential in purebred French Large White pigs. Both lines consisted of six to eight sires and about 40 dams per generation, and each dam produced two litters. The selection criterion in the line S was the *in vivo* glycolytic potential (IVGP) of the *longissimus* (ld) muscle, measured on a shot-biopsy sample removed at 75 kg live weight on boars and gilts from first-parity litters. In addition, the post mortem glycolytic potential (PMGP) of ld, *semimembranosus* (sm) and *semispinalis capitis* (sc) muscles was recorded on pigs from second-parity litters slaughtered at 100 kg live weight. Throughout the experiment, 2 981 and 454 animals were recorded for IVGP and PMGP, respectively. A consistent decrease in IVGP and, to a lesser extent, in PMGP was obtained in the line S compared with the line C. Estimates of genetic changes per generation were -0.18 , -0.11 , -0.07 and -0.09 SD units of the trait for IVGP and PMGP of ld, sm and sc muscles, respectively. The REML heritability estimates were 0.25 ± 0.02 , 0.15 ± 0.06 , 0.14 ± 0.06 and 0.17 ± 0.05 for the above four traits, respectively. The REML estimate of genetic correlation of IVGP with PMGP ld (0.87 ± 0.15) was somewhat higher than those of IVGP with PMGP of sm and sc (0.56 ± 0.14 and 0.68 ± 0.13 , respectively). It is concluded that downward selection on muscle glycolytic potential may be effective in pigs. © Inra/Elsevier, Paris

pig / muscle glycolytic potential / selection experiment / genetic parameters

* Correspondence and reprints
E-mail: larzul@toulouse.inra.fr

Résumé – Sélection pour abaisser le potentiel glycolytique du muscle chez le porc Large White. I. Réponses directes. Une expérience de sélection comportant une lignée sélectionnée (S) et une lignée témoin (C) a été conduite sur six générations en vue d'abaisser le potentiel glycolytique du muscle chez des porcs de race pure Large White français. L'une et l'autre lignée était constituée de 6–8 pères et d'environ 40 mères par génération, et chaque mère produisait deux portées. Le critère de sélection de la lignée S était le potentiel glycolytique in vivo (IVGP) du muscle *longissimus* (ld), mesuré sur un échantillon prélevé par biopsie au poids vif de 75 kg chez les mâles et femelles issus des premières portées. Le potentiel glycolytique post mortem (PMGP) des muscles ld, *semimembranosus* (sm) et *semispinalis capitis* (sc) a été mesuré chez des porcs issus des deuxièmes portées abattus au poids vif de 100 kg. Sur l'ensemble de l'expérience, 2 981 animaux ont été mesurés pour IVGP et 454 animaux pour PMGP. Un abaissement notable de IVGP et, dans une moindre mesure, de PMGP a été obtenu dans la lignée S, par rapport à la lignée C. Les évolutions génétiques par génération ont été estimées à $-0,18$, $-0,11$, $-0,07$ et $-0,09$ unité d'écart type phénotypique du caractère respectivement pour IVGP et PMGP des muscles ld, sm et sc. Les héritabilités, estimées par la méthode REML, ont été respectivement de $0,25 \pm 0,02$, $0,15 \pm 0,06$, $0,14 \pm 0,06$ et $0,17 \pm 0,05$ pour ces mêmes caractères. L'estimée REML de la corrélation génétique entre IVGP et PMGP du muscle ld ($0,87 \pm 0,15$) a été plus élevée que celles existant entre IVGP et PMGP des deux autres muscles (respectivement $0,56 \pm 0,14$ et $0,68 \pm 0,13$ pour les muscles sm et sc). Il est conclu que la sélection pour un faible potentiel glycolytique du muscle peut être efficace chez le porc. © Inra/Elsevier, Paris

porc / potentiel glycolytique musculaire / expérience de sélection / paramètres génétiques

1. INTRODUCTION

From the early 1980s, the French pig industry has become more and more interested in meat quality, particularly in yields of curing-cooking ham processing, the most important part of pig meat transformation. One of the main factors that affect technological meat quality, as defined here by the technological yield of curing-cooking ham processing, is the extent of the muscle pH fall post mortem. In France, a meat quality index was established from post mortem measurements (ultimate pH, reflectance, water holding capacity) made on ham muscles in order to predict technological yield [12]. This index was introduced into the breeding objective to maintain meat technological quality while improving growth rate, lean meat content and feed efficiency. The value of introducing a selection criterion on meat quality measured on live animals was studied earlier from a theoretical point of view [21]. However, such an approach has not been applied so far on a large scale in practical pig breeding except in very particular situations, e.g. halothane phenotype or genotype for proneness to the PSE meat condition. The principle was to choose a criterion easy to measure, heritable, genetically related to meat quality and preferably favourably linked to other production traits under selection. Such a live criterion would allow meat quality to be assessed directly on candidates for selection instead of slaughtered sibs. Experimental results showed that muscle glycolytic potential (GP), an estimator of the intra vitam muscle glycogen content, is closely related to ultimate pH and meat technological quality [14]. The glycolytic potential is based on the measurement of the main muscle

glucidic compounds susceptible to transformation into lactic acid through post mortem glycolysis, thus determining the extent of muscle pH fall. Moreover, this trait can be measured on the *longissimus* muscle of live animals by shot biopsy [27].

The question was to study the value of the in vivo glycolytic potential measurement as a selection criterion for improving technological quality of meat in a population free of the RN⁻ allele that is known to greatly affect GP [16, 17]. Thus, a selection experiment aiming at decreasing muscle glycolytic potential, measured on live animals, was carried out from 1988 onwards using pigs from a breed, the French Large White, presumably free of the HALⁿ and RN⁻ alleles [25].

2. MATERIALS AND METHODS

2.1. Experimental animals

The animals involved in the present study were purebred French Large White pigs. The base population was constituted by the progeny of a foundation breeding stock consisting of 24 unrelated AI boars and 42 sows. Male and female progeny were randomly allocated to two closed lines. One line (line S) was selected downward for muscle glycogen content as assessed by the in vivo glycolytic potential (IVGP) of the *longissimus* muscle, whereas the control line (line C) was randomly bred. Both lines were maintained contemporaneously under standardised conditions at the Inra experimental farm of Bourges-Avord.

Starting from the base population, the selection experiment was conducted over six generations from 1989 to 1996. Both lines consisted of six to eight sires and about 40 dams per generation. Each dam was planned to produce two litters at around 12 and 17.5 months of age. After weaning at 4 weeks of age, animals were reared in a post-weaning unit until 10 weeks of age and then moved to fattening pens of 10–12 animals of the same line and the same sex. During the fattening period (from 25 to 100 kg live weight), animals were given ad libitum access to a commercial standard diet (17.0 % crude protein, 1.5 % crude fat, 4.5 % crude fibre, 6.8 % ash, 0.85 % lysine, 3.091 Mcal ME/kg).

Replacement boars and gilts were chosen among first-parity litter progeny and the generation interval was about 14 months. In the control line, animals kept for breeding were randomly chosen on a within-family basis. Except in a few cases, each sire was replaced by one of his sons and each dam by one of her daughters. In the selected line, a strict mass selection was practised with a proportion selected of around 8 and 35 % for boars and gilts, respectively. In each generation, the mating scheme for both lines was set up to minimise the increase in the average coefficient of inbreeding of the resulting progeny. The average coefficients of inbreeding within line and generation were computed from the inverse of the numerator relationship matrix [9]. At the end of the experiment, the average coefficient of inbreeding reached 6–8 % in the lines C and S, respectively.

For assessing the correlated responses to selection for carcass and meat quality traits, one animal per second-parity litter, either a female or a castrated male, was randomly chosen to be slaughtered at 100 kg live weight in a series

comprising five or six individuals from each line. Animals slaughtered in a given series were from the same fattening batch. The animals fasted for 16 h before they were transported for 2 h to a commercial abattoir. Then, animals were allowed to rest for an additional 18 h before being slaughtered by electrical stunning and immediate exsanguination.

2.2. Measurements

Boars and gilts from first-parity litters were measured for the selection criterion (IVGP) at about 75 kg live weight. A small muscle sample of about 1 g was drawn by shot biopsy from the *longissimus* muscle, at 7–8 cm behind the 14th rib on the left side, and around 5 cm from the dorsal line [27]. Penetration depth was 4.5–5 cm. The biopsy was carried out in the fattening pen without any restraint to ensure minimal stress. The muscle sample was immediately trimmed of skin and fat. For the first four generations, samples were crushed in liquid nitrogen and freeze dried before being homogenised in 10 mL of 0.5 M perchloric acid. From the fourth generation onwards, muscle samples were directly homogenised in perchloric acid. Then, 0.5 mL of homogenate was taken for the simultaneous determination of muscle glycogen, glucose and glucose-6-phosphate contents using the enzymatic method of Dalrymple and Hamm [4]. The rest of the perchloric acid homogenate was centrifuged at 2 500 *g* for 10 min and the supernatant was used for lactate determination [1]. Glycolytic potential was given by the formula of Monin and Sellier [18], i.e. $GP = 2 \{[\text{glycogen}] + [\text{glucose}] + [\text{glucose-6-phosphate}]\} + [\text{lactate}]$, and expressed in micromoles equivalent lactate per g of fresh tissue.

Post mortem glycolytic potential (PMGP) was also measured on slaughtered animals from second parity-litters. Thirty minutes after slaughter, 1-g samples of the *longissimus*, *semimembranosus* and *semispinalis capitis* muscles were drawn and homogenised in 10 mL of 0.55 M perchloric acid for determination of the muscle glycolytic potential as previously described. The choice of these three muscles was based on both their anatomical location and their contractile and metabolic properties, which are known to be different [13, 15]: a white muscle from the loin (*longissimus*) or from the ham (*semimembranosus*) and a red muscle (*semispinalis capitis*).

The pigs were reared and slaughtered in compliance with the current national regulations prevailing for commercial slaughtering and animal research experimentation.

Numbers of animals measured for muscle glycolytic potential (IVGP and PMGP) are given in *table I*.

2.3. Statistical analysis

Preliminary least squares analyses were performed using the GLM procedure of SAS [22] to estimate the effect of live weight at biopsy or carcass weight as a covariate for IVGP and PMGP, respectively, and the fixed effects of sex, day of measurement nested within generation, generation, line and line \times generation combination. Annual selection differentials were calculated in each line, from differences between the average performance of animals selected as parents and the average performance of all animals measured in each line, weighted

Table I. Numbers of animals measured for muscle glycolytic potential.

Trait	Number of pigs	Number of sires	Number of litters	Body weight at measurement (kg) ^a
In vivo glycolytic potential (IVGP)	2 981	126	573	75.5 (8.2)
Post mortem glycolytic potential (PMGP)	454	92	397	101.1 (3.6)

^a In brackets, within-generation standard deviation.

by the proportion of their offspring in the individuals measured in the next generation [10]. In the line C, selection differentials were calculated to detect unintentional selection. Cumulative selection differential was calculated as the difference between the cumulative selection differential in the selected line and the cumulative selection differential in the control line to take into account the unintentional selection made in the line C. The annual selection response was measured as the difference between the average performance of animals in the line S and the average performance of animals in the line C. The realised heritability for IVGP was determined by regressing the annual selection response against the cumulative selection differential, with regression constrained to pass through the origin, because both lines were taken from the same base population [10]. The approximate standard errors for the annual phenotypic response and for the realised heritability were calculated accounting for drift variance and measurement error variance [10, 11].

For the genetic analysis, the model included live weight for IVGP or carcass weight for PMGP as a covariate, sex as a fixed effect, biopsy date (75 levels) for IVGP or slaughter series (39 levels) for PMGP and animal additive genetic values as random effects. For IVGP, litter effect was also taken into account as a random effect. Litter effect was not included in the model for PMGP because only one slaughtered animal per second-parity litter was recorded for this trait.

Variance and covariance components were estimated using a restricted maximum likelihood (REML) procedure applied to a four-trait individual animal model with missing data. All the ancestors of the recorded animals, up to the grand-parents of the base population from which the control and selected lines were derived, were taken into account for establishing the numerator relationship matrix of the animals. There were 3 701 individuals in the pedigree file. The estimation of genetic parameters was performed with version 3.2 of the VCE computer package, using a quasi-Newton algorithm with exact first derivatives to maximise the log likelihood [20]. Heritability (h^2) was computed as the additive genetic variance divided by the adjusted phenotypic variance. Approximate standard errors of variance components and genetic parameters were obtained from the inverse of an approximation of the Hessian matrix when convergence was reached [24].

Additive genetic breeding values were estimated in a four-trait analysis using the BLUP (best linear unbiased predictor) methodology applied to an individual animal model as previously described for REML analysis. The

REML-estimated genetic parameters were used in the model. The analysis was performed using the PEST computer package [8]. The response to selection was estimated from the within-generation line difference (selected-control) for average predicted breeding values (genetic response). When averaging predicted breeding values for a trait, only individuals recorded for that trait were taken into account. For simplification, the approximate standard errors for the annual S-C differences were calculated for each trait with REML-estimated parameters, considering that animal breeding values were computed in univariate analyses [26].

Coheritabilities for the selection criterion (IVGP) with the PMGPs were calculated from REML estimates. Their standard errors were approximated from the standard errors of parameters using the first order term of a Taylor expansion.

3. RESULTS

3.1. Selection differentials

Table II gives the number of animals measured and selected, as well as the selection differentials, by line, generation and sex, and the selection intensity (standardised selection differential) by line and generation. In line S, selection on females became really effective from generation 3 and selection on males became more intense from the same generation onwards. It should be pointed out that an unintended selection differential occurred in line C at generation 3 and, to a lesser extent, at generation 5. When randomly choosing control boars at generation 3, it happened that one boar showing the lowest IVGP value was kept.

3.2. Responses to selection

The line differences (selected-control) for phenotypic means and average breeding values across generations are shown in figure 1. For IVGP, the phenotypic and genetic line differences are very similar whatever the generation. A significant response was observed for IVGP from generation 3 onwards in the selected line. At the last generation, the line difference for IVGP average breeding values was $-26.4 \pm 7.3 \mu\text{mol/g}$ fresh tissue, i.e. -1.15 phenotypic standard deviation units of the trait. In line S, a decrease in PMGP was observed in the three muscles. At the last generation, the differences between the average breeding values of the selected and control lines were -14.4 ± 5.4 , -7.9 ± 4.7 , and $-8.1 \pm 3.8 \mu\text{mol/g}$ fresh tissue for the PMGP of the *longissimus*, *semimembranosus* and *semispinalis capitis* muscles, respectively. When considering the regression of line difference on generation number, the genetic responses to selection per generation were $-4.2 \mu\text{mol/g}$ for IVGP and -2.5 , -1.4 and $-1.3 \mu\text{mol/g}$ per generation for the PMGP of the *longissimus*, *semimembranosus* and *semispinalis capitis* muscles, respectively.

3.3. Heritabilities

The realised heritability (h^2) value of IVGP was 0.21 ± 0.05 and was slightly lower than the REML h^2 estimate of 0.25 (table III). Estimated h^2 of PMGP

Table II. Numbers of animals measured and selected for IVGP, and selection differentials per sex, line and generation.

Line	Generation	Females			Males			Overall selection intensity
		Number measured	Number selected	Selection differential (in $\mu\text{mol/g}$)	Number measured	Number selected	Selection differential (in $\mu\text{mol/g}$)	
Base	G0	187			190			
S	G0		29	-5.2		10	-25.0	-0.76
	G1	71	40	-5.5	71	7	-28.8	-0.95
	G2	103	42	-13.7	62	11	-28.8	-0.96
	G3	143	49	-26.1	103	6	-44.6	-1.40
	G4	157	43	-21.9	178	7	-41.5	-1.54
	G5	151	41	-19.5	137	7	-49.7	-1.48
	G6	102	-	-	120	-	-	-
C	G0		22	1.1		14	-0.1	0.02
	G1	41	40	1.9	38	8	-2.6	-0.09
	G2	111	41	-2.8	50	9	0.2	-0.08
	G3	104	47	0.4	70	8	-26.3	-0.59
	G4	154	47	-0.9	110	8	-5.8	-0.15
	G5	141	39	-10.3	135	8	-9.8	-0.32
	G6	139	-	-	113	-	-	-




Figure 1. Differences across generations between selected (S) and control (C) lines for in vivo glycolytic potential of the *longissimus* muscle (IVGP) and post mortem glycolytic potential of the *longissimus* (PMGP Ld), *semimembranosus* (PMGP sm) and *semispinalis capitis* (PMGP sc) muscles., phenotypic trends; —, genetic trends; - - -, one SD unit.

was of the same magnitude for the three muscles studied (0.14–0.17). The part of the variance of PMGP due to slaughter date was larger for the *semispinalis capitis* muscle (31 %) than for the *semimembranosus* and *longissimus* muscles (20 %). The part of the variance due to the biopsy date for IVGP of the *longissimus* muscle (26 %) was slightly larger than that due to the slaughter date for PMGP of the same muscle.

3.4. Correlations

Phenotypic and genetic correlations are given in table IV. Genetic correlations (r_A) between IVGP (*longissimus*) and PMGP decrease from the *longissimus* muscle to *semispinalis capitis* muscle and to the *semimembranosus* muscle. As could be expected, the highest genetic correlation was found between IVGP and PMGP measured in the *longissimus* muscle. These two measurements, even though they were not made under the same conditions (live weight, environment), were made on the same muscle at close locations. The lowest genetic correlation (0.56 ± 0.14) was found between IVGP and PMGP of the

Table III. Estimates of least squares means by sex, phenotypic standard deviation (σ_p), heritability (h^2), litter effect (c^2) and date of measurement effect (s^2) for in vivo (IVGP) and post mortem muscle glycolytic potential (PMGP), in $\mu\text{mol/g}$ of fresh tissue, for the *longissimus* (ld), *semimembranosus* (sm) and *semispinalis capitis* (sc) muscles.

Item		IVGP ld	PMGP ld	PMGP sm	PMGP sc
Means by sex (\pm SE)	boars	183 \pm 1			
	gilts	193 \pm 1	118 \pm 2	101 \pm 1	48 \pm 1
	castrates		113 \pm 2	100 \pm 1	45 \pm 1
σ_p		23	23	21	14
$h^2 \pm$ SE		0.25 \pm 0.02	0.15 \pm 0.06	0.14 \pm 0.06	0.17 \pm 0.05
$c^2 \pm$ SE		0.09 \pm 0.01			
$s^2 \pm$ SE		0.26 \pm 0.03	0.20 \pm 0.03	0.20 \pm 0.03	0.31 \pm 0.04

Table IV. Estimates of phenotypic correlations^a (below diagonal), genetic correlations (\pm SE) (above diagonal), and coheritabilities (\pm SE).

Trait	PMGP ld	PMGP sm	PMGP sc
In vivo glycolytic potential (IVGP)	0.87 \pm 0.15	0.56 \pm 0.14	0.68 \pm 0.13
<i>longissimus</i> (PMGP ld)		0.83 \pm 0.13	0.48 \pm 0.25
<i>semimembranosus</i> (PMGP sm)	0.65***		-0.08 \pm 0.34
<i>semispinalis capitis</i> (PMGP sc)	0.51***	0.52***	
Coheritability with IVGP	0.17 \pm 0.05	0.11 \pm 0.03	0.14 \pm 0.03

^a Level of significance: *** $P < 0.001$.

semimembranosus muscle, though the *longissimus* and *semimembranosus* muscles are of similar metabolic type. This correlation could indicate that muscles were not reacting in the same way during the slaughter procedure and/or the *semimembranosus* muscle is less homogeneous than the *longissimus* muscle.

The phenotypic correlations between PMGPs of the three muscles studied were of the same order (0.5–0.6). However, the r_A estimates for PMGP were strongly dependent on the two muscles considered. They were very low (-0.08 ± 0.34 between *semimembranosus* and *semispinalis capitis* PMGPs) to very high (0.83 ± 0.13 between *longissimus* and *semimembranosus* PMGPs), the genetic correlation between *longissimus* and *semispinalis capitis* PMGPs being intermediate (0.48 ± 0.25). Coheritabilities (table IV) were of the same order for PMGPs of the three muscles.

4. DISCUSSION

During the experiment, over all generations, we observed a significant genetic change in the in vivo glycolytic potential observed in the selected line, with a significant difference between the two lines at the 6th generation. The absence of the RN⁻ allele, known to largely increase IVGP [17], was reinforced by the absence of high IVGP values in the base population. Genetic response differed from one generation to another. The lack of response during the first two generations may be explained by the intensity of selection which was not as high as initially planned. The explanation is to be found in the number of animals measured and selected and in the way of selecting parents. In the first generations, the number of males and females recorded for IVGP was lower than in the following generations. And during these first generations, parents were not selected among all the batches because, at time of mating, animals of the last batch had not reached puberty. To avoid this problem in the next generations, generation interval was increased. Then, the response to selection was much greater in the following generations.

The estimated value of heritability for in vivo glycolytic potential of the *longissimus* muscle is much lower than values reported for the same trait by Le Roy et al. [16]. In the latter study, the RN⁻ allele was segregating in the two composite lines involved, leading to very high values of heritability (0.86–0.90). The heritability values estimated by Burlot [2] in the same lines taking into account the RN genotype effect were of medium range (0.32–0.42), and only slightly higher than the present estimated value.

The proportion of variance due to slaughter date for PMGP of the *longissimus* muscle is in agreement with that estimated by de Vries et al. [6] for ultimate pH of the same muscle (0.22). In earlier studies [3, 23], the proportion of variance due to slaughter date was found to be even larger for ultimate pH of *longissimus* and *adductor femoris* (about 0.40), but these estimates were probably inflated by other factors (e.g. genetic) than the slaughter date effect itself. Thus, slaughter date seems to affect to the same extent muscle glycogen content and ultimate pH of *longissimus* muscle. In the present study, glycolytic potential of the *semispinalis capitis* muscle was more affected by slaughtering conditions than PMGP of white muscles.

The differences between responses obtained for IVGP on the one hand and PMGP of the three muscles on the other hand could be explained by the fact that IVGP and PMGP are different traits: IVGP is a measurement of the muscle glycogen content under minimal stress conditions whereas PMGP is a measurement of muscle glycogen content in animals stressed by pre-slaughtering conditions. Moreover, the average value of IVGP is higher than the average value of PMGP (*table III*) in the *longissimus* (190 and 115 $\mu\text{mol/g}$ fresh tissue, respectively). One part of the difference between measurements could be due to difference in live weight at measurement: as shown by Dalrymple et al. [5], muscle glycogen content tended to decrease when animals grew older. However, the muscle glycolytic potential is essentially lowered by all the treatments applied before slaughter (fasting, transportation, resting time at the abattoir, etc.) known to deplete muscle glycogen content (for review, Fernandez and Tornberg [7]). The phenotypic correlation between IVGP and PMGP in the

longissimus muscle is known to be low (0.20) for animals free of the RN⁻ allele [14].

5. CONCLUSION

The significant genetic changes in the *in vivo* glycolytic potential observed in the selected line and the medium heritability value found for this trait allow it to be concluded that it is possible to introduce this trait as a selection criterion [14]. From estimated genetic correlations with post mortem glycolytic potential, it appears that selection for decreasing *in vivo* glycolytic potential in the *longissimus* muscle would result in a lesser decrease in post mortem glycolytic potential. However, the magnitude of the correlated genetic change in the latter trait depends on the metabolic type of the muscle considered. The consequences of downward selection on IVGP for meat quality and production traits are developed in subsequent papers.

ACKNOWLEDGEMENTS

This work was part of the project 'régulation du potentiel glycolytique du muscle chez le porc' and was supported by grants from the Inra-Agrobio programme initiated in 1990. Thanks are due to Pierre Vernin (SRV, Theix), Hervé Lagant (SGQA, Jouy-en-Josas) and the staff of the pig experimental unit in Bourges-Avord for their technical assistance. Comments and suggestions of one referee were appreciated.

REFERENCES

- [1] Bergmeyer H.U., in: Bourne G.H. (Ed.), *Methods of Enzymatic Analysis*, Academic Press, New York, 1974, pp. 1127–1196.
- [2] Burlot T., Étude de la variabilité génétique du potentiel glycolytique du muscle dans deux lignées synthétiques porcines, Mémoire de fin d'études, ISA de Beauvais, 1994, 43 pp.
- [3] Charpentier J., Monin G., Ollivier L., Correlations between carcass characteristics and meat quality in Large White pigs, in: *Proceedings of the 2nd International Symposium on Condition and Meat Quality of Pigs*, Pudoc, Wageningen, 1971, pp. 255–260.
- [4] Dalrymple R.H., Hamm R., A method for the extraction of glycogen and metabolites from a single muscle sample, *J. Food Technol.* 8 (1973) 439–444.
- [5] Dalrymple R.H., Kastenschmidt L.L., Cassens R.G., Glycogen and phosphorylase in developing red and white muscle, *Growth* 37 (1973) 19–34.
- [6] De Vries A.G., van der Wal P.G., Lang T., Eikelenboom G., Merks J.W.M., Genetic parameters of pork quality and production traits in Yorkshire populations, *Livest. Prod. Sci.* 40 (1994) 277–289.
- [7] Fernandez X., Tornberg E., A review of the causes of variation in muscle glycogen content and ultimate pH in pigs, *J. Muscle Foods* 2 (1991) 209–235.
- [8] Groeneveld E., Kovac M., A generalized computing procedure for setting up and solving mixed linear models, *J. Dairy Sci.* 73 (1990) 513–531.
- [9] Henderson C.R., A simple method for computing the inverse of a numerator relationship matrix in prediction of breeding values, *Biometrics* 32 (1976) 69–83.
- [10] Hill W.G., Estimation of realised heritabilities from selection experiments. II. Selection in one direction, *Biometrics* 28 (1972) 767–780.

- [11] Hill W.G., Design of quantitative genetics selection experiments, in: Robertson A. (Ed.), *Selection Experiments in Laboratory and Domestic Animals*, Commonwealth Agricultural Bureaux, Slough, UK, 1980, pp. 1–13.
- [12] Jacquet B., Sellier P., Runavot J.P., Brault D., Houix Y., Perrocheau C., Gogué J., Boulard J., Prédiction du rendement technologique de la fabrication du « jambon de Paris » à l'aide de mesures prises à l'abattoir, in: 16^{es} Journées de la recherche porcine en France, Paris, 1–3 février 1984, Institut technique du porc, Paris, 1984, pp. 49–58.
- [13] Laborde D., Talmant A., Monin G., Activités enzymatiques métaboliques et contractiles de 30 muscles du Porc. Relations avec le pH ultime atteint après la mort, *Reprod. Nutr. Dev.* 25 (1985) 619–628.
- [14] Larzul C., Le Roy P., Monin G., Sellier P., Variabilité génétique du potentiel glycolytique du muscle chez le porc, *Inra Prod. Anim.* 11 (1998) 183–197.
- [15] Lefaucheur L., Le Dividich J., Mourot J., Monin G., Ecolan P., Krauss D., Influence of environmental temperature on growth, muscle and adipose tissue metabolism, and meat quality in swine, *J. Anim. Sci.* 69 (1991) 2844–2854.
- [16] Le Roy P., Przybylski W., Burlot T., Bazin C., Lagant H., Monin G., Étude des relations entre le potentiel glycolytique du muscle et les caractères de production dans les lignées Laconie et Panshire, in: 26^{es} Journées de la recherche porcine en France, Paris, 1–3 février 1994, Institut technique du porc, Paris, 1994, pp. 311–314.
- [17] Le Roy P., Monin G., Elsen J.M., Caritez J.C., Talmant A., Lebreton B., Lefaucheur L., Mourot J., Juin H., Sellier P., Effect of the RN genotype on growth and carcass traits in pigs, in: Van Arendonk J.A.M. (Ed.), *Book Abstr. of the 47th E.A.A.P. Ann. Meeting*, Wageningen Pers, Wageningen, the Netherlands, 1996.
- [18] Monin G., Sellier P., Pork of low technological quality with a normal rate of muscle pH fall in the immediate post mortem period: the case of the Hampshire breed, *Meat Sci.* 13 (1985) 49–63.
- [19] Monin G., Mejenes-Quijano A., Talmant A., Sellier P., Influence of breed and muscle metabolic type on muscle glycolytic potential and meat pH in pigs, *Meat Sci.* 20 (1987) 149–158.
- [20] Neumaier A., Groeneveld E., Restricted maximum likelihood estimation of covariances in sparse linear models, *Genet. Sel. Evol.* 30 (1998) 3–26.
- [21] Ollivier L., Potier D., L'amélioration de la qualité de la viande porcine par la sélection, in: 7^{es} Journées de la recherche porcine en France, Paris, 19–21 février 1975, Institut technique du porc, Paris, 1975, pp. 293–302.
- [22] SAS[®], SAS[®] Inst. Inc., Cary, NC, 1985.
- [23] Schepers J., Influence of environmental and genetic factors on meat quality, *Acta Agric. Scand. suppl.* 21 (1979) 20–31.
- [24] Searle S.R., Casella G., McCulloch C.E., *Variance Components*, John Wiley & Sons, New York, NY, 1992.
- [25] Sellier P., Monin G., Genetics of pig meat quality, a review, *J. Muscle Foods* 5 (1994) 187–219.
- [26] Sorensen D.A., Kennedy B.W., Analysis of selection experiments using mixed model methodology, *J. Anim. Sci.* 63 (1986) 245–258.
- [27] Talmant A., Fernandez X., Sellier P., Monin G., Glycolytic potential in Longissimus dorsi muscle of Large White pigs as measured after in vivo sampling, in: *Proceedings of the 35th Int. Cong. Meat Sci. and Technol.*, Copenhagen, Danish Meat Research Institute, Roskilde, 1989, pp. 1129–1131.