



**HAL**  
open science

# PCR cloning and detection of point mutations in the eburicol 14 alpha-demethylase (CYP51) gene from *Erysiphe graminis* f. sp. hordei, a "recalcitrant" fungus

Christophe Délye, Lydia Bousset, Marie-France Corio-Costet

## ► To cite this version:

Christophe Délye, Lydia Bousset, Marie-France Corio-Costet. PCR cloning and detection of point mutations in the eburicol 14 alpha-demethylase (CYP51) gene from *Erysiphe graminis* f. sp. hordei, a "recalcitrant" fungus. *Current Genetics*, 1998, 34, pp.399-403. hal-02692966

**HAL Id: hal-02692966**

**<https://hal.inrae.fr/hal-02692966>**

Submitted on 1 Jun 2020

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Copyright

## PCR cloning and detection of point mutations in the eburicol 14 a-demethylase (CYP51) gene from *Erysiphe graminis* f. sp. *hordei*, a “recalcitrant” fungus

Délye C<sup>a</sup>, Bousset L<sup>b</sup>, Corio-Costet MF<sup>a</sup>

<sup>a</sup> *Unité de Recherches Intégrées sur la Vigne, INRA, Domaine de la Grande Ferrade, B. P. 81, F-33883 Villenave d’Ornon Cedex, France*

<sup>b</sup> *Laboratoire de Pathologie Végétale, INRA, F-78850 Thiverval-Grignon, France*

### Abstract

Molecular studies of some micro-organisms are hampered by the difficulty of obtaining sufficient amounts of nucleic acids. A cloning strategy based on PCR has therefore been used to clone the eburicol 14a-demethylase (*CYP51*) gene of the obligate fungus *Erysiphe graminis* f. sp. *hordei* (*Egh*) using minute amounts of genomic DNA. The *CYP51* gene encodes the enzymatic target of a major group of fungicides. Sequencing *CYP51* from different *Egh* isolates revealed the occurrence of two alleles for this gene. An allele-specific PCR assay was developed to detect each *CYP51* allele.

**Keywords:** *Erysiphe graminis*, Polymerase chain reaction, Cytochrome P450, Fungicide

### Introduction

Genome sequencing programs, as well as cloning and sequencing of particular genes, has facilitated a number of studies concerning the cell biochemistry and physiology, taxonomy, phylogeny and/or genetic diversity of numerous micro-organisms. However, the vast majority of known sequences have been obtained by screening genomic libraries. This time-consuming approach can be used on micro-organisms such as fungi capable of saprophytic growth, in which it is easy to obtain large amounts of pure fungal material. It has, however, only rarely been employed for “recalcitrant” species such as slow-growing or obligate biotrophic phytopathogenic fungi (Sherwood and Somerville 1990; Justesen et al. 1996; Délye et al. 1997a), mostly because of the difficulty of obtaining adequate amounts of nucleic acids. This indeed is the major bottleneck for molecular and biochemical studies of many phytopathogenic fungi of major economic importance, such as rusts, downy mildews and powdery mildews. The availability in the literature of numerous sequences coding for genes of taxonomic or metabolic importance now makes it possible to clone a variety of genes from “recalcitrant” micro-organisms species, without having any sequence data from these species. The purpose of the present work was to clone the single-copy gene (*CYP51*) encoding cytochrome P450 eburicol 14a-demethylase from the major cereal pathogen *Erysiphe graminis* DC. ex M rat f. sp. *hordei* Em. Marchal (*Egh*), a haploid, obligate biotrophic fungus. *CYP51* is a key enzyme of the sterol biosynthesis pathway (for a review see Yoshida 1993) which is the enzymatic target for a major class of antifungals called DeMethylase Inhibitors (DMIs) (Gadher et al. 1983). Although *Egh* has developed field resistance towards DMIs for many years (Walmsey-Woodward et al. 1979), the molecular basis of this resistance is still unknown.

To clone the *Egh CYP51* gene, we successfully used a PCR cloning procedure consisting of the three following steps: (1) production of a fragment of the gene using PCR with degenerated primers, (2) digestion of 100–200 ng of genomic DNA and ligation into a plasmid vector, and (3) cloning of the remainder of the gene sequence using two sets of primer pairs. In each of these pairs, one primer is based on the plasmid sequence, and the other is based on the sequence of the gene fragment obtained in (1).

### Materials and methods

#### *Fungal material.*

A total of seven single-spore *Egh* isolates has been used for this work. Isolates AL1, Tr2, JEH11 and Ge3 are reference isolates sensitive to DMIs (Limpert 1987). They were kindly provided by Dr. E. Limpert (Swiss Institute of Technology, Zürich, Switzerland). Isolates GV1-22 and AP2-19 were collected in 1995 in Châlons en Champagne (France) and in Amiens (France), respectively. Isolate 92-18 was collected in 1992 in Grignon (France). The collection of these three isolates has been done in DMI-treated areas before the beginning of the

spraying program. Isolates were mass-produced by blowing conidia on detached leaves from the susceptible barley (*Hordeum vulgare*) cultivar Igri, which were then placed on a 0.4% water agar medium, to which 30 mg/l of benzimidazole was added, in 9-cm-diameter Petri dishes. Inoculated leaves were incubated for 14 days at 16°C under 24 h/day illumination (10 mE/m<sup>2</sup>/s). Conidia were then dislodged onto a glass slide and collected in a microcentrifuge tube using a razor blade. Conidia were kept at –20°C and freeze-dried before nucleic-acid extraction. *DNA extraction and PCR assays.* For PCR-based experiments, DNA was extracted from about 5 mg dry weight of conidia as described by Délye et al. (1995). PCR conditions were also as described (Délye et al. 1997 b). Amplified fragments were visualised on 1% (w/v) agarose gels run in 0.5 × tris-borate EDTA buffer and stained with ethidium bromide (0.4 mg/ml gel).

#### *Obtaining a fragment of the Egh CYP51 gene.*

The three published *CYP51* sequences from filamentous fungi were aligned. These sequences are from *Penicillium italicum* (Van Nistelrooy et al. 1996), *Ustilago maydis* (Hargreave and Keon 1996) and *Uncinula necator* (Délye et al. 1997a). Degenerate primers D-CR1 (5'-TAYGGIGAYRTITTYWSITT, sense strand, degeneracy = 64) and D-CR4 (5'-ATCATCATIYSIGCDATYTC, antisense strand, degeneracy=24), which correspond to filamentous fungi conserved *CYP51* aminoacid sequences Y-G-D-(I,V)-F-(T,S)-F and E-I-A-(H,G)-M-M-I, respectively, were used to amplify a fragment with an expected size of about 700 bp. The annealing temperature was 50 °C and the final primer concentration was 2.0 mM for primer D-CR1 and 0.6 mM for primer D-CR4. Cloning and sequencing procedures were as described (Délye et al. 1997 a).

*Obtaining the complete Egh CYP51 sequence.* DNA was extracted from 15 mg dry weight of conidia from isolate AL1 using a CTAB protocol (Murray and Thompson 1980). Approximately 200 ng of DNA were digested to completion with *Xba*I (MBI Fermentas) following the manufacturer's instructions, purified using the Cleanmix purification kit (Talent) and ligated into the *Xba*I site of the pBS+ plasmid vector (Stratagene). Ligation was performed overnight at 15°C using T4 DNA ligase (MBI Fermentas) in a total volume of 10 ml. Primer pairs PBS-U (5¢-GCCAGTGAATTGTAATACGACTCACTATAGG)/ C51-700R (5'-TCGACCTAGTATATCCGTCGTCTTCTTACCC) and PBS-R (5'-CCATGATTACGCCAAGCTCGAAATTAACCC)/ C51-700 (5'-TGCGTCTTCTTACAAAGATGGCAGCCCG) were used to amplify DNA fragments encompassing sequences located upstream of (primers PBS-U and C51-700R) and downstream from (primers PBS-R and C51-700) the sequence obtained using the degenerate primers D-CR1 and D-CR4. Primers PBS-U and PBS-R targeted sequences flanking the polylinker site of the pBS+ plasmid vector. Primers C51-700 and C51-700R targeted sequences located on the D-CR1/D-CR4 *CYP51* fragment. Primers were used at a final concentration of 0.1 mM each. Amplifications were performed on 1/10 dilutions of the ligation mix. The cycling program consisted of 37 cycles with 30 s denaturation at 94°C and 2 min annealing and extension at 72°C. Primers C51Egh (5'-CCGTCCTTATCGCAAGATTTG) and C51EghR (5'-CATAGTAGCCTGTAATCTAAGC), targeting sequences located 10 bp upstream of the ATG initiation codon and 66 bp downstream from the final TGA codon respectively, were used at a final concentration of 0.2 mM each with an annealing temperature of 60°C to amplify a 1791-bp fragment encompassing the whole *CYP51* sequence from the *Egh* isolates. *CYP51 cDNA cloning.* Reverse-transcription PCR, followed by cloning and sequencing, was used to confirm that the putative introns found within the *Egh CYP51* sequence were readily excised from transcripts. Total RNA was extracted from 5 mg dry weight of conidia from isolate AL1 using the RNable™ reagent (Eurobio). Reverse transcription was performed using a first-strand cDNA synthesis kit (Pharmacia). A cDNA fragment was amplified from 1/10 dilutions of the reverse transcription mix using primers C51Egh and C51EghR. Cloning and sequencing procedures were as above.

#### *Detection of CYP51 alleles: allele-specific PCR.*

Primers MUT-T (5'-AATTAGGACAGTCAA) and MUT-A (5'-AATTAGGACAGTCAT) were designed specifically for the priming of *Egh CYP51* sequences exhibiting respectively a T or a A at nucleotide 458, considering that a 3¢ mismatch does not prime in a PCR reaction under specific annealing temperatures (Sommer and Tautz 1989). Each of these primers was used in PCR amplifications, together with primer C51Egh, to amplify a 503-bp fragment. Both primers in each primer pair were used at a final concentration of

0.1 mM. The annealing temperature was 47°C for primer pair C51Egh/MUT-T and 49 °C for primer pair C51Egh/MUT-A.

## Results

### Cloning the *Egh CYP51* gene


Amplifications with primers D-CR1 and D-CR4 yielded a single major PCR product of the expected size (data not shown). This 714-bp fragment potentially encoded a 220 amino-acid polypeptide interrupted by a putative 52-bp intron. This polypeptide displayed 76.4% identity and 80.9% similarity with the corresponding region of *U. necator* CYP51.

Amplification of *Egh* *Xba*I-digested DNA ligated into the pBS+ plasmid vector using primer pairs PBS-U/C51-700R and PBS-R/C51-700 yielded a single major PCR product of approximately 1100 and 1400 bp, respectively (data not shown). The 1100-bp fragment encompassed a partial open reading frame of 357 bp including a putative 51-bp intron and the first 60 bp of the D-CR1/D-CR4 fragment. The 1400-bp fragment contained a partial open reading frame of 681 bp terminated by a TGA codon and including the last 58 bp of the D-CR1/D-CR4 fragment. The nucleotide sequence of the *Egh CYP51* gene, obtained after joining the overlapping sequences included in the 1100-bp fragment, the 714-bp D-CR1/D-CR4 fragment and the 1400-bp fragment was 1672-bp long.

### Cloning the *Egh CYP51* cDNA

The *Egh CYP51* gene was interrupted by two putative introns at nucleotides 247–297 and 496–547, respectively. A comparison of the size of the PCR fragments amplified with primers C51Egh and C51EghR from genomic DNA or cDNA (Fig. 1), as well as the sequencing of the cDNA obtained from isolate AL1, revealed that the two putative introns were readily excised. The remainder of the cDNA coding sequence was identical to that of the genomic DNA.

**Figure 1** PCR products obtained by amplification of cDNA (tracks 1–7) and genomic DNA (tracks 8–14) from 1 mg dry weight of conidia from *Egh* isolates using primers C51Egh and C51EghR. Tracks: L molecular-weight marker (1-kb DNA ladder, Gibco-BRL); H H<sub>2</sub>O negative control (no DNA); 1 and 8 isolate GV1-22; 2 and 9 isolate AP2-19; 3 and 10 isolate 92-18; 4 and 11 isolate AL1; 5 and 12 isolate *Ge3*; 6 and 13 isolate *Tr2*; 7 and 14 isolate *JEH11*



### Analysis of the *Egh CYP51* sequence

The inferred 522 amino-acid protein encoded by the 1569-bp coding sequence of the 1672-bp *Egh CYP51* gene was compared to the known complete CYP51 sequences. The strongest homology (72.2% identity and 77.8% similarity) was with *U. necator* CYP51. This is much more than the 40% homology required for two cytochrome P450 genes to belong to the same family (Nelson et al. 1993). The weakest homology (28.4% identity and 42.1% similarity) was with *Sorghum bicolor* CYP51 (Bak et al. 1997). Alignment of the predicted amino-acid sequence of CYP51 from *Egh* with those of the filamentous fungus *U. necator*, the yeast *Saccharomyces cerevisiae* (Kalb et al. 1987), the human *Homo sapiens* (Strömstedt et al. 1996) and the plant *S. bicolor* is given in Fig. 2. The alignments highlight the conserved domains CR1–CR6. Domains CR1–CR4

are believed to be involved in substrate specificity (Aoyama et al. 1996). Domains CR5 (k-helix) and CR6 (heme-binding domain) are hallmarks of cytochrome P450 (Gotoh 1993).

The two introns in the *Egh CYP51* gene interrupted nucleotide sequences encoding domain CR1 and a region located eight amino-acids downstream from CR2 (Fig. 2). The position of the two introns in the *Egh CYP51* gene is exactly the same as that of the two introns in the *U. necator CYP51* gene and of the two first introns in the *P. italicum CYP51* gene. These are the only other known fungal *CYP51* sequences for which introns have been identified to-date. The *Egh CYP51* sequence will appear in Genbank under the accession number AF052515.

**Figure 2** Alignment of predicted *CYP51* amino-acid sequences from *Egh* (Genbank: AF052515) with representative *CYP51* sequences from the fungal (*U. necator*, Genbank: U72657, *S. cerevisiae*, Genbank: M18109), animal (*H. sapiens*, Genbank: U23942) and plant (*S. bicolor*, Genbank: U74319) kingdoms. The alignment was generated using the GCG program pileup (Devereux et al. 1984). Gaps have been introduced to maximise the alignment (gap weight: 12.0, gap length weight: 4.0). Identical residues are marked by an asterisk. CR1–CR6 regions are in bold, and underlined in the *Egh* sequence. The position of introns is arrowed. Numbers on the right refer to amino-acid positions in *Egh CYP51*. Identity and similarity percentages are given at the left bottom of the figure


<i>E. graminis</i>	~~~~~	~~~~~	MGISES	FMPYLPQLL	QLGFGIALAS	GILSLLLLL	FLNVLKQLLF	46
<i>U. necator</i>	~~~~~	~~~~~	MYIADI	LSDLLTQOTT	RYGWIFMVT	IAFSIILLAV	GLNVLSQLLF	
<i>S. cerevisiae</i>	~~~~~	~~~~~	M SATKSI	VGGEA	LE . YVNI	GL SHFLALPLAQ	RISLIIIPF	IYINWQLLY
<i>H. sapiens</i>	MAAAAGMLL	GLLQAGGSVL	QGAMEKVTGG	NLLSMLLIAC	APTLSLVYLI	RLAAGHLVQL		
<i>S. bicolor</i>	~~~~~	~~~~~	MDLA	DIPQQQLMA	GLALVVATVI	FLKLLSFRS		
				intron ↓	CR1			105
<i>E. graminis</i>	K.NPNEPPIV	FHWIPIIGST	ISYGMNPYKF	<b><u>FHESQAKYGN</u></b>	<b><u>IFTF</u></b>	ILLGKK	TTVYLGRQGN	
<i>U. necator</i>	R.RPYEPPVV	FHWFPPIIGST	ISYGLDPYKF	<b><u>YFDCRAKYGD</u></b>	<b><u>IFTF</u></b>	ILLGKK	TTVYLGQIGN	
<i>S. cerevisiae</i>	SLRRDRPPLV	PYWIPWVGS	VVYGMKPYEF	<b><u>FEECQKKYGD</u></b>	<b><u>IFSF</u></b>	VLLGRV	MTVYLGPKGH	
<i>H. sapiens</i>	PAGVKSPPYI	PSPIPFLGHA	IAPGKSPIEF	<b><u>LENAYEKYGP</u></b>	<b><u>VFSF</u></b>	IMVGT	FTVYLLGSDAA	
<i>S. bicolor</i>	GGGKRLPPT	IPGAPVVGGL	VKPMRGPIMP	<b><u>IREQYAAALGS</u></b>	<b><u>VFTV</u></b>	PIITRR	ITFLIGPEVS	
	*	*	*	*	*	*	*	*
				intron ↓	CR2			165
<i>E. graminis</i>	NFILNGKLRD	VNAEEIYTVL	<b><u>TFPVFGTDVV</u></b>	<b><u>YDCPN</u></b>	SKLME	QKKFMKAALT	TEAFRSYVPI	
<i>U. necator</i>	NFILNGKLRD	VNAEEIYTVL	<b><u>TFPVFGTDVV</u></b>	<b><u>YDCPN</u></b>	SKLME	QKKFMKALT	IEAFHSYVTI	
<i>S. cerevisiae</i>	EFVFNARLAD	VSAAEAAYHL	<b><u>TFPVFGKQVI</u></b>	<b><u>YDCPN</u></b>	SRIME	QKKFVKGALT	KEAFKSYVPL	
<i>H. sapiens</i>	ALLFNKSNED	LNAEDVYSRL	<b><u>TFPVFGKQVA</u></b>	<b><u>YDVNP</u></b>	VFVLE	QKKMLKSLGN	IAHFQKHVSI	
<i>S. bicolor</i>	AHFFKGENAE	MSQQEVY .RF	<b><u>NVPTFGPGVV</u></b>	<b><u>FDVDYSVRQE</u></b>	QRPFPTEALR	ANKLRSYVDQ		
	*	*	*	*	*	*	*	*
								222
<i>E. graminis</i>	IQNEVKSFIE	KCDDFR . . . K	SKGIINIDAV	MAEITIIYAS	HTLQGEVRD	RFDSSI	<b><u>AVLY</u></b>	
<i>U. necator</i>	IQNEVEAYIN	NCVSFQ . . . G	ESGTVNISKV	MAEITIIYAS	HALQGEVRE	RFDSSI	<b><u>FAALY</u></b>	
<i>S. cerevisiae</i>	IAEVEYKYFR	DSKNFRLNER	TGTTIDVMVT	QPEMTIFTAS	RSLLGKEMRA	KLDTDF	<b><u>FAYLY</u></b>	
<i>H. sapiens</i>	IEKETKEYFE	S . . . . . WG	ESGEKNVFEA	LSELIIILTAS	HCLHGKEIRS	QLNEKVAQLY		
<i>S. bicolor</i>	MVAEAEYFS	K . . . . . WG	ESGTVDLKYE	LEHLIILTAS	RCLLGEVRE	KLFDDV	<b><u>SALF</u></b>	
	*	*	*	*	*	*	*	*
								281
<i>E. graminis</i>	<b><u>HDLDMGFTPI</u></b>	NFMLH . WAPL	PHNRARDHAQ	RTVAKIYMEI	INSRRTQKET	DDSNLDMWQ		
<i>U. necator</i>	<b><u>HDLDMGFTPI</u></b>	NFTFY . WAPL	PWNRARDHAQ	RTVARTYMIN	IQARREEKRS	GENKHDMWE		
<i>S. cerevisiae</i>	<b><u>SDDLKGFPTPI</u></b>	NFVFP . NLPL	EHYRKRDRHAQ	KAISGTYMSL	IKERR . KNN	DIQDRDLIDS		
<i>H. sapiens</i>	<b><u>ADLDGGFSHA</u></b>	AWLLPGWLP	PSFRRRDRAH	REIKDIFYKA	IQKRRQSQEK	IDD . . . ILQ		
<i>S. bicolor</i>	<b><u>HDLDNQIQPI</u></b>	SVLFP . YLPI	PAHKRRDKAR	ARLAEIFATI	IKSRKASQGS	EED . . . MLQ		
	***	*	*	*	*	*	*	*
								340
<i>E. graminis</i>	LMR . SSKYDG	TPVPDKETAH	<b><u>MMIALLMAGO</u></b>	<b><u>HSSSSSTWI</u></b>	MLWLAARPDI	TEELYQEQLR		
<i>U. necator</i>	LMR . STYKDG	TPVPDKETAH	<b><u>MMIALLMAGO</u></b>	<b><u>HSSSSSTWI</u></b>	MLWLAARPDI	MEELYEEQLR		
<i>S. cerevisiae</i>	LMKNSYKDG	VKMDDQETAN	<b><u>LLIGVLLMGGQ</u></b>	<b><u>HTSAA</u></b>	TSAWI	LLHLAERPDV	QQELYEEQMR	
<i>H. sapiens</i>	TLLDATYKDG	RPLDDQEVAG	<b><u>MLIGLLLAGQ</u></b>	<b><u>HTSSST</u></b>	SAMM	GFPLARDKTL	QKKCYLEQKT	
<i>S. bicolor</i>	CFIDSKYKNG	RPTTEGEVTS	<b><u>LLIALPAGQ</u></b>	<b><u>HTSSIT</u></b>	TWT	GAYMLRFKQY	FAEAVEEQKD	
	**	*	*	*	*	*	*	*
								395
<i>E. graminis</i>	LLGSE . LPP	LKYEDLSKLS	LHQNVLKEVL	<b><u>RLH</u></b>	APIHSIL	RKVKNMPV .	..PGTSYVIP	
<i>U. necator</i>	IFGSEKFPFP	LQYEDLSKQLQ	LHQNVLKEVL	<b><u>RLH</u></b>	APIHSIM	RKVKNPMIV .	..PGTKYVIP	
<i>S. cerevisiae</i>	VLDGKK . E	LTYPDLLQEMP	LLNQTIKETL	<b><u>RMHHP</u></b>	PLHSLF	RKVKMDMHV .	..PNTSYVIP	
<i>H. sapiens</i>	VCG . ENLPP	LYTDYQLKDLN	LLDRCIRETLL	<b><u>RLRPP</u></b>	IMIMM	RMARTPQIVA	...GYTIP	
<i>S. bicolor</i>	VMK . RHGDK	IDHDILAEMD	VLYRCIKRAL	<b><u>RLHPP</u></b>	LIMLL	RQSHSDFTVT	TKEGKEYDIP	
	*	*	*	*	*	*	*	*
								455
<i>E. graminis</i>	KTHSLLAAPG	WTSRDASYFP	NPLKWDPHRW	DTGSGGVIGT	DMEDEKPDYG	YGLISTGAAS		
<i>U. necator</i>	TSHVLISSPG	CTSQDATFFP	DPLKWDPHRW	DIGSGKVLGN	DAVDEKYDYG	YGLTSTGASS		
<i>S. cerevisiae</i>	AGYHVLVSPG	YTHLRDEYFP	NAHQFNIRHW	NKDSA . . SS	YSVGEVEVDY	FGAISKGVSS		
<i>H. sapiens</i>	PGHQVCVSP	VNQRLLKDSW	ERLDFNPDRY	..LQDNP . . . . .	ASGEK			
<i>S. bicolor</i>	KGHIVATSPS	FANRLPHIYK	NPDSYDPRF	GPGREED . . . . .	KAAGAF			
	*	*	*	*	*	*	*	*
								515
<i>E. graminis</i>	<b><u>PYLFPFAGRRH</u></b>	<b><u>RCIGEQFATV</u></b>	QLVVTIMATMV	RSFKFHNLDG	RNSVAETDYS	SMFSRPMQPA		
<i>U. necator</i>	<b><u>PYLFPFAGRRH</u></b>	<b><u>RCIGEQFATL</u></b>	QLVVTIMATMV	RFRFRNIDG	KQGVVKTIDS	SLFSPMLGTP		
<i>S. cerevisiae</i>	<b><u>PYLFPFAGRRH</u></b>	<b><u>RCIGEHFAYC</u></b>	QLGVLMISIFI	RTLKWHYPEG	K . TVPPDPFT	SMVTLPTGPA		
<i>H. sapiens</i>	<b><u>AYVFPFAGRRH</u></b>	<b><u>RCIGENFAYV</u></b>	QIKTIWSTML	RLYEPDLIDG	...YFPTVNY	TTMIHTPMP		
<i>S. bicolor</i>	<b><u>SYISFPFAGRRH</u></b>	<b><u>GCLGEPFAYL</u></b>	QIKAIWTHLL	RNFEPFLVSP	...FPENDWN	AMVVGIKGEV		
	*	*	*	*	*	*	*	*
								522
<i>E. graminis</i>	TIWEKR---	----		Identity		Similarity		
<i>U. necator</i>	LIGWEKR---	----		72.2%		77.8%		
<i>S. cerevisiae</i>	KLIWEKRNP	QKI~		47.0%		56.5%		
<i>H. sapiens</i>	VIRY . KRRSK-	----		37.7%		46.4%		
<i>S. bicolor</i>	MVNY . KRRKLV	VDN		28.4%		42.1%		

### Variability of the *Egh CYP51* gene and allele-specific PCR amplifications

The *CYP51* gene was cloned and sequenced from *Egh* isolate GV1-22 using primers C51Egh and C51EghR. Comparison with the sequence from isolate AL1 revealed a single A-to-T change at nucleotide 458, resulting in the presence of a phenylalanine residue at position 136 in *CYP51* of isolate GV1-22 instead of the tyrosine residue in *CYP51* of isolate AL1. The amino acid at position 136 is located within the highly conserved CR2 domain, which is presumably involved in substrate recognition (Aoyama et al. 1996).

Allele-specific PCR amplifications using primer pairs C51Egh/MUT-T and C51Egh/MUT-A revealed that isolates AP2-19 and 92-18 exhibited a T at nucleotide position 458, whereas isolates Tr2, Ge3 and JEH11 displayed a A at the same position (Fig. 3). Amplifications were reproducibly obtained using DNA extracted from less than 1 mg dry weight of conidia.

**Figure 3** PCR products obtained by allele-specific amplification of DNA from 1 mg dry weight of conidia from *Egh* isolates using primer pairs C51Egh/MUT-T (tracks 1–7) and C51Egh/MUT-A (tracks 8–14). Tracks: L molecular-weight marker (1-kb DNA ladder, Gibco-BRL); H H<sub>2</sub>O negative control (no DNA); 1 and 8 isolate GV1-22; 2 and 9 isolate AP2-19; 3 and 10 isolate 92-18; 4 and 11 isolate AL1; 5 and 12 isolate Ge3; 6 and 13 isolate Tr2; 7 and 14 isolate JEH11



## Discussion

### Cloning the *Egh CYP51* gene using PCR

By using degenerate primers designed on the basis of the highly conserved amino-acid sequences of *CYP51* from the only three known filamentous fungi, we were able to amplify a portion of the *Egh CYP51* gene. The remainder of the *Egh CYP51* sequence was also obtained by PCR from less than 200 ng of *Egh* genomic DNA, without the need to establish and screen a genomic library in bacteria. We used only 50 ml of the 100 ml of the *Xba*I-digested DNA ligation mix to achieve this result. The PCR cloning strategy may thus be considered as very “DNA-sparing”. PCR primers C51Egh and C51EghR, derived from *Egh CYP51* flanking sequences, enabled us to verify that the two putative introns identified within this gene were readily excised. Successful amplification of a DNA fragment encompassing *CYP51* was obtained from DNA extracted from less than 1 mg dry weight of *Egh* conidia (data not shown). Successful amplification of a cDNA fragment encompassing *CYP51* was obtained from RNA extracted from 1–2 mg dry weight of *Egh* conidia. Cloning and sequencing *CYP51* from a variety of *Egh* isolates is thus a task that does not require massive production of conidia from this fungus. The availability of the *Egh CYP51* gene now paves the way for molecular and biochemical studies of resistance to DMIs in this fungus.

### PCR detection of a point mutation

Two *CYP51* alleles were identified in *Egh*. The DMI-sensitive reference isolate AL1 exhibited a TAT (tyrosine) codon at position 136. Isolate GV1-22, which was collected from DMI-treated fields, exhibited a TTT (phenylalanine) codon at position 136. The occurrence of a phenylalanine residue at this position has

been reported for a laboratory mutant of *P. italicum* (De Waard 1996), for field isolates of *U. necator* (Délye et al. 1997b), and for clinical isolates of *Candida albicans* (Sanglard et al. 1998) that were all highly resistant to DMIs. DMI-sensitive isolates from these three fungi all exhibited a tyrosine residue instead of a phenylalanine residue at this position.

A PCR assay was developed for specific detection of each *Egh CYP51* allele. Allele-specific PCR was preferred to PCR-RFLP because of (1) the omission of the digestion step and (2) the lack of a suitable restriction site at nucleotide 458. We found that *CYP51* sequences from DMI-sensitive reference isolates AL1, Tr2, Ge3 and JEH11 all exhibited a A at nucleotide 458. Isolates GV1-22, AP2-19 and 92-18, which were collected from DMI-treated fields, all exhibited a T at that position. It is thus possible that, as found in *P. italicum*, *U. necator* and *C. albicans*, substitution of a phenylalanine residue for a tyrosine residue may cause resistance to DMIs in *Egh*. Allele-specific PCR will enable us to follow the distribution of the two identified *CYP51* alleles in field populations of *Egh* using only minute amounts of fungal material. Should a correlation between the presence of the phenylalanine/tyrosine at position 136 and resistance to DMIs be established, then allele-specific PCR may be used to monitor resistance of *Egh* to such compounds in the field.

The PCR gene-cloning strategy we used has proven efficient for the cloning of a gene from an organism in which obtaining significant amounts of DNA is a long and difficult procedure. This strategy may thus be recommended for other similar systems, provided that (1) a few hundred ng of DNA can be obtained from the organism in question, and (2) a few sequences of the investigated DNA exist in the literature and display conserved regions. For microorganisms such as fungi or bacteria living exclusively inside their host tissues, a PCR cloning strategy targeting genes that are either not present in the host genome, or are not conserved between the host and the micro-organism, can be considered.

## References

- Aoyama Y, Noshiro M, Gotoh O, Imaoka S, Funae Y, Kurosawa N, Horiuchi T, Yoshida Y (1996) Sterol 14-demethylase P450 (P45014DM) is one of the most ancient and conserved P450 species. *J Biochem* 119:926–933
- Bak S, Kahn RA, Olsen CE, Halkier BA (1997) Cloning and expression in *Escherichia coli* of the obtusifoliol 14a-demethylase of *Sorghum bicolor* (L.) Moench, a cytochrome P450 orthologous to the sterol 14a-demethylases (CYP51) from fungi and mammals. *Plant J* 11:191–201
- Délye C, Corio-Costet M-F, Laigret F (1995) A RAPD assay for strain typing of the biotrophic grape powdery mildew fungus *Uncinula necator* using DNA extracted from the mycelium. *Exp Mycol* 19:234–237
- Délye C, Laigret F, Corio-Costet M-F (1997 a) Cloning and sequence analysis of the eburicol 14a-demethylase gene of the obligate biotrophic grape powdery mildew fungus. *Gene* 195:29–33
- Délye C, Laigret F, Corio-Costet M-F (1997b) A mutation in the 14a-demethylase gene of *Uncinula necator* that correlates with resistance to a sterol biosynthesis inhibitor. *Appl Environ Microbiol* 63:2966–2970
- Devereux J, Haeberli P, Smithies O (1984) A comprehensive set of sequence analysis programs for the VAX. *Nucleic Acids Res* 12:387–395
- De Waard MA (1996) Molecular genetics of resistance in fungi to azole fungicides. In: Brown TM (ed) *Molecular genetics and evolution of pesticide resistance*. ACS symposium series 645. American Chemical Society, Washington, District of Columbia, pp 62–71
- Gadher PE, Mercer I, Baldwin BC, Wiggins TE (1983) A comparison of the potency of some fungicides as inhibitors of sterol 14-demethylation. *Pestic Biochem Physiol* 19:1–10
- Gotoh O (1993) Evolution and differentiation of P-450 genes. In: Omura T, Ishimura Y, Fujii-Kuriyama Y (eds) *Cytochromes P-450*, 2nd edn. Kodansha, Tokyo and VCH, Weinheim, pp 255–272
- Hargreave JA, Keon JPR (1996) Isolation of an *Ustilago maydis* ERG11 gene and its expression in a mutant deficient in sterol 14a-demethylase activity. *FEMS Microbiol Lett* 139:203–207
- Justesen A, Somerville S, Christiansen S, Giese H (1996) Isolation and characterization of two novel genes expressed in germinating conidia of the obligate biotroph *Erysiphe graminis* f. sp. *hordei*. *Gene* 170:131–135

- Kalb VF, Woods CW, Turi TG, Dey CR, Sutter TR, Loper JC (1987) Primary structure of the P450 lanosterol demethylase gene from *Saccharomyces cerevisiae*. *DNA* 6:529–537
- Limpert E (1987) Frequencies of virulence and fungicide resistance in the European barley powdery mildew population in 1985. *J Phytopathol* 119:298–311
- Murray HG, Thompson WF (1980) Rapid isolation of high-molecular-weight plant DNA. *Nucleic Acids Res* 8:4321–4325
- Nelson DR, Kamataki T, Waxman DJ, Guengerich FP, Estabrook RW, Feyereisen, R, Gonzalez FJ, Coon MJ, Gunsalus IC, Gotoh O, Okuda K, Nebert DW (1993) The P450 superfamily: update on new sequences, gene mapping, accession numbers, early trivial names of enzymes and nomenclature. *DNA Cell Biol* 12:1–51
- Sanglard D, Ischer F, Koymans L, Bille J (1998) Amino-acid substitutions in the cytochrome P450 lanosterol 14a-demethylase (CYP51A1) from azole-resistant *Candida albicans* clinical isolates contribute to resistance to antifungal agents. *Antimicrobial Agents Chemother* 42:241–253
- Sherwood JE, Somerville SC (1990) Sequence of the *Erysiphe graminis* f. sp. *hordei* gene encoding  $\beta$ -tubulin. *Nucleic Acids Res* 18:1052
- Sommer R, Tautz D (1989) Minimal homology requirements for PCR primers. *Nucleic Acids Res* 17:6749
- Strömstedt M, Rozman D, Waterman MR (1996) The ubiquitously expressed human CYP51 encodes lanosterol 14a-demethylase, a cytochrome P450 whose expression is regulated by oxysterols. *Arch Biochem Biophys* 329:73–81
- Van Nistelrooy JGM, Van den Brink JM, Van Kan JAL, Van Gorcom RFM, De Waard MA (1996) Isolation and molecular characterization of the gene encoding eburicol 14a-demethylase (CYP51) from *Penicillium italicum*. *Mol Gen Genet* 250:725–733
- Walmsey-Woodward DJ, Laws FA, Whittington WJ (1979) Studies on the tolerance of *Erysiphe graminis* f. sp. *hordei* to systemic fungicides. *Ann Plant Biol* 92:199–209
- Yoshida Y (1993) Lanosterol 14a-demethylase (cytochrome P450<sub>14DM</sub>). In: Schenkman H, Grein K (eds) *Cytochromes P450*. Springer Verlag, Berlin, pp 627–639