



HAL
open science

The influence of soil nature on polyphenols in walnut tissues. A possible explanation of differences in the expression of walnut blight

P. Radix, Catherine Bastien, C. Jay-Allemand, Gérard Charlot, F. Seigle-Murandi

► To cite this version:

P. Radix, Catherine Bastien, C. Jay-Allemand, Gérard Charlot, F. Seigle-Murandi. The influence of soil nature on polyphenols in walnut tissues. A possible explanation of differences in the expression of walnut blight. *Agronomie*, 1998, 18 (10), pp.627-637. hal-02698308

HAL Id: hal-02698308

<https://hal.inrae.fr/hal-02698308v1>

Submitted on 1 Jun 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

The influence of soil nature on polyphenols in walnut tissues. A possible explanation of differences in the expression of walnut blight

Pascal Radix^{a,c}, Catherine Bastien^b, Christian Jay-Allemand^b,
Gérard Charlot^c, Françoise Seigle-Murandi^{a*}

^aLaboratoire de botanique, cryptogamie, biologie cellulaire et génétique, groupe pour l'étude du devenir des xénobiotiques dans l'environnement (GEDEXE), UFR de pharmacie, université J.-Fourier (Grenoble I), BP 138, 38243 Meylan cedex, France

^bStation d'amélioration des arbres forestiers, Inra, centre de recherche d'Orléans, 45160 Olivet, France

^cCTIFL, « Les Colombières », 38160 Chatte, France

(Received 11 August 1997; accepted 20 October 1998)

Abstract – Walnut blight is the major disease in the walnut orchards of Dauphiné (France). The causal bacterium *Xanthomonas campestris* pv. *juglandis* can infect leaves, catkins, buds and nuts and can induce important crop losses. Field observations have shown that the nature of the soil seems to play a prominent role in crop losses. The hypothesis that soil may modify the amount of polyphenols in walnut tissues and modify the natural resistance of fruits to necrosis has been assumed. We have shown that the nature of the soil, especially its permeability, induces variations in polyphenols in walnut tissues. Differences of susceptibility to walnut blight between the two cultivars studied, Franquette and Parisienne, may also be explained by different behaviours in relation to polyphenols. Moreover, the massive use of copper treatments against walnut blight may induce an imbalance in polyphenol contents and could explain crop losses despite repeated copper sprays. (© Inra/Elsevier, Paris.)

Juglans / walnut blight / polyphenols / soil

Résumé – Influence de la nature du sol sur la composition en polyphénols des tissus de noix. Une explication possible des différences observées dans l'expression de la bactériose du noyer. La bactériose est la principale maladie des noyers dans les vergers du Dauphiné (France). La bactérie responsable de cette maladie, *Xanthomonas campestris* pv. *juglandis*, infecte les feuilles, les chatons, les bourgeons, les fruits et peut causer d'importantes pertes de récolte. Des observations de terrain ont montré que la nature du sol semblait influencer l'importance des pertes de récolte. L'hypothèse que la nature du sol puisse modifier la composition en polyphénols des tissus et ainsi modifier la résistance naturelle des fruits à la nécrose a été faite. Nous avons montré que la nature du sol, spécialement sa perméabilité, induit des

Communicated by Gérard Guyot (Avignon)

*Correspondence and reprints

Tel: (33) (0)4 76 04 10 04; fax: (33) (0)4 76 04 10 05; e-mail: francoise.seigle.murandi@ujf-grenoble.fr

variations dans la teneur en polyphénols des tissus. De même, les différences de sensibilité entre Franquette et Parisienne, les deux variétés étudiées, pourraient être expliquées par des différences de teneur en polyphénols. De plus, l'emploi massif de traitements cuivriques peut provoquer un déséquilibre dans la composition polyphénolique des tissus et pourrait expliquer dans certains cas, le manque d'efficacité de ces traitements. (© Inra/Elsevier, Paris.)

Juglans / bactériose du noyer / polyphénols / sol

1. INTRODUCTION

Walnut (*Juglans* sp.) is an economically important tree crop in many countries around the world, such as Australia, Chile, China, France, India, Iran, Italy, Spain, USA, etc. Its production has been increasing steadily over the past decade, especially in the United States and in China [25]. Sixty percent of world exportations were by the USA, only 25 % by China and 7 % by both India and France [22]. Walnut blight is the major disease in the orchards of Dauphiné (SE France); it was already mentioned by Garavel in 1959 [8]. The causal bacterium *Xanthomonas campestris* pv. *juglandis* can infect leaves, catkins, buds and nuts [19]. Crop losses, due to necrosis of the fruit, can reach 60–80 %. Necrosis of the fruit constitutes the only real problem of this disease. At the beginning of the maturation of the fruit, harmful necrosis usually appears around the stigma as an oily dark brown spot. These young infected nuts generally fail to grow and, then, fall a few days later. Late season necrosis can stain the shell and make a part of the harvest unsellable. Up to now, the only way to reduce the damages caused by this disease has been to use periodical copper treatments although they are not always effective. Concern for environmental side-effects has restricted the use of copper-based treatments. Moreover, strains resistant to copper have been found [4, 9] and copper accumulation in the soil may disturb the metabolism of the trees [24].

Numerous field observations have shown that beyond the infection of walnut orchards by *Xanthomonas campestris* pv. *juglandis*, the nature of the soil and other factors such as the cultivar seem to play a prominent role in crop losses [6]. Very often, in spite of repeated copper treatments (eight sprays per year at 2.5 kg ha⁻¹ Cu²⁺ each), it

has been observed that orchards (especially for the Franquette cultivar) located on shallow, permeable and acidic soils are more damaged than orchards located on deep, poorly permeable and basic soils.

The wounding responses in plant tissues are often related to phenolic metabolism, and these responses have a general role in protecting the injured organ from infection [1, 16, 20]. The hypothesis that soil may modify the phenolic content of tissues and increase or decrease the natural resistance of fruit to blight has been submitted.

Naphtoquinones such as juglone (5-hydroxy-1,4 naphtoquinone) are major compounds of walnut tissues (especially of husk tissues) [2, 3, 17]. Hedin et al. [12, 13] established that juglone and related compounds accounted for the resistance of pecan (*Carya illinoensis*) to *Fusicladium effusum*; cultivars less affected by *Fusicladium effusum* generally had a higher level of juglone [12, 13].

Moreover, these compounds are known for their toxicity to micro-organisms. Juglone is a depressant agent which immobilises and tranquillises fish, Daphnia, frogs and mammals [26]. Juglone inhibited *Lemna minor* growth, chlorophyll content and photosynthesis at treatments between 10 and 40 mM [14]. The hypothesis that phenolic compounds could be used as stress detectors and could induce a change in the susceptibility of the fruit to this disease has been submitted. The first aim of this study was to reveal any possible influence of soil factors on the amount of phenolic compounds in husk tissues. In the second place, it was to show through biochemical disturbances that shallow, permeable, acidic soils, induce higher occurrences of necrosis.

The study was conducted in two parts. A first study aimed to find out which soil factors may influence the amount of phenolic compounds in

husk tissues. A screening study, made to select the most pertinent factors, employed a fractional factorial design.

A second study, adapted from the results of the first experimental design, was conducted for a more acute analysis aiming to describe and explain soil effects on necrosis through variations in polyphenolic compounds. A complete factorial design was used.

2. MATERIALS AND METHODS

2.1. Experimental design 1 (fractional factorial design 2⁸⁻⁴)

In this exploratory experiment, 8 different factors of variation have been studied for their potential effect on the quantitative variations of some phenolic compounds. The effect of the nature of the soil was evaluated through three factors with two levels each (according to our experimental design requirements), corresponding to two physical characteristics, the depth of the soil (Dp) and its permeability (Pm) and to one chemical characteristic, the pH (pH).

A possible cultivar effect (Cv) was taken into account by using two different varieties: Franquette which is the most common cultivar found in Dauphiné (with a medium susceptibility to walnut blight) and Parisienne, a cultivar that has been grown in Dauphiné for a longer time and is rather resistant to walnut blight.

In order to determine the possible role of copper treatment on walnut metabolism (Cu), the absence or the presence of such a treatment was considered as the fifth factor. Copper sprays are usually performed from blossom to late growing, especially during rainy periods. No treatment is applied during fall and winter.

Nuts were collected from two different trees (Tree) in each selected orchard. Since preliminary analysis had proved that variability between fruits is negligible compared to variability between zones of tissue sampled on one fruit, only two factors of variation were studied at the fruit level. The first one was related to the presence or the absence of necrosis (Nec). The second one distinguished two parts in the tissues sampled on the nut, the apical zone and the parietal zone (Ti).

In order to combine a reduced size of the experiment with the analysis of numerous factors of variation, a fractional factorial design 2⁸⁻⁴ was finally chosen [5, 10]. By this means, the study was limited to 32 tissue samples collected from only 16 different orchards (*table I*). Finally, data analysis tried to show any influences (statistically significant) of independent variables (factors)

Table I. Characteristics of the orchards studied, in relation to the fractional factorial design 2⁸⁻⁴.

Orchards	Depth	Soil permeability	pH	Cultivar	Copper treatment
A	deep	poor	basic	Franquette	absence
B	deep	poor	basic	Franquette	presence
C	deep	poor	basic	Parisienne	absence
D	deep	poor	basic	Parisienne	presence
E	deep	permeable	acidic	Franquette	absence
F	deep	permeable	acidic	Franquette	presence
G	deep	permeable	acidic	Parisienne	absence
H	deep	permeable	acidic	Parisienne	presence
I	shallow	poor	acidic	Franquette	absence
J	shallow	poor	acidic	Franquette	presence
K	shallow	poor	acidic	Parisienne	absence
L	shallow	poor	acidic	Parisienne	presence
M	shallow	permeable	basic	Franquette	absence
N	shallow	permeable	basic	Franquette	presence
O	shallow	permeable	basic	Parisienne	absence
P	shallow	permeable	basic	Parisienne	presence

on dependent variables, i.e. the concentration of polyphenols in walnut husk tissues.

Independent variables (factors) were:

- Cv, *Juglans regia* cultivars: Franquette (Fr) or Parisienne (Pa);
- Cu, copper treatment: absence or presence;
- Dp, soil depth: deep or shallow;
- Pm, soil permeability: permeable or poorly permeable;
- pH, soil pH: acidic or basic;
- Nec, necrosis: absence or presence;
- Ti, tissue: apex (1) or parietal (2);
- Tree, tree: tree n_1 1 or tree n_1 2;

Dependent variables were:

- JUG, juglone concentration;
- HJG, hydrojugloneglucoside concentration;
- MYR, myricitrin concentration;
- QUE, quercitrin concentration;
- CAF, caffeic acid concentration;
- HCA, hydroxycinnamic acid concentration (partially identified);
- FLV₁, flavonol 1 concentration (partially identified);
- FLV₂, flavonol 2 concentration (partially identified);
- FLV₃, flavonol 3 concentration (partially identified);

Sample collection:

Samples were collected in each orchard from two different trees, at four different times (June 2, June 17, June 26, July 7 1993) corresponding to four different phenological stages. Orchards differed in at least one of these three factors: soils, *Juglans regia* cultivars (Franquette or Parisienne) and copper treatment (absence or presence).

Only nuts were collected (10 nuts per sample) and immediately immersed in situ in liquid nitrogen (only a few seconds elapsed between picking and freezing). Then, samples were freeze-dried and stored under vacuum in darkness.

2.2. Experimental design 2 (complete factorial design 2⁴)

After the results of the first experimental design, we selected the most relevant factors in order to constitute a complete factorial design 2⁴. Only sound nuts were selected and analysed in this part.

Independent variables (factors) were:

- Cv, *Juglans regia* cultivars: Franquette (Fr) or Parisienne (Pa);
- Cu, copper treatment: absence or presence;

- Pm, soil permeability: permeable (1) or poorly permeable (2);
- Ti, tissue: apex (1) or parietal (2).

Dependent variables were:

- JUG, juglone concentration;
- HJG, hydrojugloneglucoside concentration;
- MYR, myricitrin concentration;
- QUE, quercitrin concentration;
- CAF, caffeic acid concentration;
- HCA, hydroxycinnamic acid concentration (partially identified);
- NAPH, naphthoquinone concentration (partially identified);
- FLAV, flavane concentration (partially identified).

Minor partially identified flavonol 1, 2, 3 were ruled out in this experimental design according to the results obtained in the first analysis. On the other hand, two other compounds (belonging to naphthoquinone and flavane families) were included because high concentrations were found during the exploratory experiment. They were implicated in interactions between cultivar, copper treatment and permeability as well.

2.3. Analysis of soluble phenolic compounds

Phenolic compounds were extracted from husk tissues and quantified according to a method described by Jay-Allemand [15]. About 20 mg of dry tissue was cut off from the freeze-dried fruit, and put in a 2-mL Eppendorf tube. Then 2 mL of acetone/water (4/1, v/v) mixture with 25 mg L⁻¹ of 6-methoxyflavone as standard, were poured in the Eppendorf tube and was homogenised by ultrasound for 30 min. Centrifugation at 12 000 rd min⁻¹ (11 250 g) lasted 20 min. Then 500 µL of supernatant were removed and dried under vacuum by means of the Speed-Vac system. Finally, the dry residue was diluted with 1 mL methanol to constitute the final sample for HPLC quantification. Samples were kept at 4 °C throughout the extraction phase.

Analyses were performed by HPLC Multisolvant Delivery System Waters (600E) with Photodiode Array Detector (Waters 991), Sample Processor (Waters Ultrawist 715), Powermate SX Plus Computer NEC. Conditions for phenolic analysis were: Column SFCC Hypersil C18 (5 µm, 25 cm); a linear gradient elution using solvent A acetonitrile/methanol (50/50, v/v), solvent B water (1 % acetic acid) was programmed from 15 % to 40 % A for 20 min, then from 40 % to 60 % A for 5 min, then from 60 % to 100 % A for 5 min, then 100 % A for 10 min, then from 100 % to 15 % A for 5 min, with a flow rate of 1 mL min⁻¹. Detection was performed at 340 nm and scanning from 220 to 450 nm.

Compounds were identified by their retention time and their spectrum in comparison with standards. Peak area was calculated in relation to an internal standard (6-methoxyflavone). Juglone, hydrojugloneglucoside, myricitrin and quercitrin were quantified as absolute values. Values were expressed in mg of compound/g dry weight (DW) of husk tissues.

2.4. Data analysis

Data were analysed according to the factorial design requirements with the S System software.

The effects of the different factors of variation on the amount of each phenolic compound were tested through an analysis of variance (ANOVA). Multivariate analysis for each date of collection were conducted through the principal component analysis (PCA).


3. RESULTS

The preliminary results (not presented) obtained from the first experimental design have mainly


shown that phenolic variations were especially due to the state of maturation of the fruit, the cultivar, the tissue and soil permeability. The highest variations of polyphenols content in husk tissues were observed during the early stages of development of fruits when the most necroses generally occur. For these reasons, only the early stages of development (sampling times 1 and 2) were further studied.

3.1. Correlations between polyphenol contents in walnut husk tissues

The principal component analysis showed that most of the variations (86 and 94 % according to the sampling times 1 and 2, respectively) among phenolic compounds were explained by the first two principal components (*figure 1*). Moreover, these two principal components were mainly represented by juglone (principal component 1) and hydroxycinnamic acid (principal component 2). This



TIME 1



TIME 2

JUG concentration of juglone
 HJG concentration of hydrojugloneglucoside
 QUE concentration of quercitrin
 HCA concentration of hydroxycinnamic acid
 CAF concentration of caffeic acid
 FLAV concentration of flavane
 NAPH concentration of naphthoquinone

Figure 1. Correlation between polyphenol contents in walnut husk tissues. Principal component analysis was performed with the second experimental design (complete factorial design) data. Correlation circles for the two sampling times (June 2, June 17) are shown.

means that all the variations between the 7 phenolic compounds studied are well described by only the juglone and the hydroxycinnamic acid components. In particular, at sampling time 2 (June 17), the first two principal components (axes 1, 2) explained almost all the variations (94 %), and juglone alone explained almost 71 % of the variations. This suggests that regulations between polyphenolic compounds exist as has been previously demonstrated [11, 18].

3.2. Influence of the factors of variations (soil permeability, copper treatment, cultivar, tissue) on polyphenols contents in walnut husk tissues

Analysis of variance was performed in order to find out which factors were the most significant for the explanation of the variations of polyphenol content among walnut husk tissues (*table II*). Concentrations of juglone, hydrojugloneglucoside,

Table II. Influence of the 8 factors of variation on polyphenol contents in walnut husk tissues at the first two sampling times. ANOVA was performed with 8 variables (polyphenol contents) and 4 factors of variation (cultivar, soil permeability, copper treatment, tissue). Values indicate *P*-values of the principal effects and interactions. Only significant effects at a 10 % level ($\text{Pr}(F) < 10\%$) are considered.

	JUG	HJG	MYR	Polyphenols		CAF	FLAV	NAPH
				QUER	HCA			
Sampling time 1								
Copper treatment								
Tissue	0.001	0.030			0.003		0.054	0.006
Interactions								
Cv.Cu.Pm	0	0		0.019	0.001	0.091		
Cv.Ti.Pm								0.058
Sampling time 2								
Permeability								0.068
Copper treatment	0.011							
Tissue					0		0.001	
Interactions								
Cv.Cu						0.001	0.001	
Cv.Pm	0.003	0.039					0.069	
Cv.Ti						0.048		
Cu.Pm						0.065		
Ti.Pm	0.050							

Cv.Cu = interaction Cultivar × Copper treatment

Cv.Pm = interaction Cultivar × Soil permeability

Cv.Ti = interaction Cultivar × Tissue

Cu.Pm = interaction Copper treatment × Soil permeability

Ti.Pm = interaction Tissue × Soil permeability

Cv.Cu.Pm = interaction Cultivar × Copper treatment × Soil permeability

Cv.Ti.Pm = interaction Cultivar × Tissue × Soil permeability

hydroxycinnamic acid and flavane were significantly different between apical and parietal tissues during the early stage of development of the fruit (sampling time 1). Naphthoquinone concentration was modified by copper treatment as well. Two weeks later (sampling time 2), hydroxycinnamic acid and flavane concentrations were significantly different between apical and parietal tissues and juglone concentration was modified by copper treatment.

3.3. Interactions between factors

The interactions between factors play an essential role in the global understanding of the problem. A description of the effects of these interactions between factors of variation on polyphenol contents is presented in *figure 2*. The cultivar factor is involved in most of the interactions. This indicates that Franquette and Parisienne cultivars have different responses in polyphenols contents either in presence of copper treatment or with regard to the permeability of the soil.

3.4. Influence of copper treatment, soil permeability and tissue factors on polyphenol contents in walnut husk tissues

Table III indicates adjusted means for each significant factor. These means were calculated with consideration for the interactions. We notice that hydrojugloneglucoside constitutes more than 20 % of the husk tissues dry weight. For both sampling times, apical husk tissues have higher levels of juglone, hydrojugloneglucoside, hydroxycinnamic acid and flavane than parietal tissues. Copper treatment globally induced a reduction in juglone content.

4. DISCUSSION

Results shown in *table II* indicate that the 'tissue' factor is the main effect in the explanation of the


variations of polyphenols. Numerous field observations have revealed that necrosis appeared almost always at the apex of the fruit. The significant difference in composition of the two tissues (apical, parietal) may be an explanation for this observation.

Table III shows that at sampling time 2 (when blight damages are most important), the concentration of juglone decreased in the presence of copper in the soils and we have already established a relationship between high levels of copper in soils and the importance of crop losses [21]. It was established by Hedin et al. [13] that juglone could account for the resistance of the fruit to blight damages: the higher the level of juglone, the higher the resistance. Even if copper did not appear as a prominent factor, it played a significant role in some interactions with other factors such as necrosis, cultivar and soil depth (*figure 2*). It was established in previous studies [7, 24] that copper accumulation in the soil may disturb the metabolism of trees, and we found in some orchards a copper content more than fifteen times the average. These results seem to indicate that copper treatments not only have a poor efficiency but induce metabolic disturbance in some cases.

4.1. Effects of interactions between factors on polyphenol contents in husk tissues

Field observations have revealed that orchards set on permeable and rather acidic soils and the Franquette cultivar are always more damaged than other orchards, in spite of copper treatment. An increased synthesis of cinnamic acid derivatives, such as caffeic acid, flavonoids, etc., and their accumulation in plant tissues are stimulated by wounding or stress [23]. This may be explained by some interactions among the factors described in this study (*figure 2*).

The juglone content in walnut husk tissues was influenced by both the cultivar and the soil's permeability (*figure 2a*). We have found an important difference of behaviour between the two types of soil, mainly for cv. Franquette (10 mg g⁻¹ juglone in



Perm 1 : poorly permeable
 Perm 2 : permeable
 Cu 1 : presence of copper
 Cu 2 : absence of copper
 Fr : cv. Franquette
 Pa : cv. Parisienne
 Ti 1 : apex
 Ti 2 : parietal
 Pr(F) : risk α

Cjug : concentration of juglone
 Chjg : concentration of hydrojugloneglucoside
 Ccaf : concentration of caffeic acid
 Cflav : concentration of flavane

Figure 2. Influence of cultivar, soil permeability, tissue and copper treatment on polyphenol contents in walnut husk tissues. Description of the significant interactions (Pr(F) < 10 %) between factors of variation.

Table III. Influence of copper treatment, soil permeability and tissue on polyphenol contents in walnut husk tissues. Values are adjusted means of polyphenol contents and are expressed in mg g⁻¹ dry weight. Only significant effects at a 10 % level (Pr(F) < 10 %) are presented.

		JUG	HJG	Polyphenols HCA	FLAV	NAPH
Sampling time 1						
Copper treatment	absence					5.47 ± 0.2
	presence					2.48 ± 0.2
Tissue	apex	19.0 ± 0.4	282 ± 5.2	18.6 ± 0.3	1.6 ± 0.2	
	parietal	15.2 ± 0.4	216 ± 5.2	15.5 ± 0.3	0.8 ± 0.2	
Sampling time 2						
Permeability	permeable					4.3 ± 0.3
	poorly permeable					5.3 ± 0.3
Copper treatment	absence	18.0 ± 1.1				
	presence	12.1 ± 1.1				
Tissue	apex			15.9 ± 0.5	2.9 ± 0.17	
	parietal			10.1 ± 0.5	1.6 ± 0.17	

poorly permeable soils / 22 mg g⁻¹ in permeable soils). The higher susceptibility of Franquette to necrosis may be explained by the fact that the permeability of the soil strongly affects the level of juglone in Franquette husk tissues whereas it does not in Parisienne. Moreover, soil permeability influences juglone contents in parietal tissues but very little in apical tissues (*figure 2b*).

The hydrojugloneglucoside content in walnut husk tissues was also influenced by both the cultivar and the soil's permeability (*figure 2c*). Concentrations of hydrojugloneglucoside in Parisienne fruit (500 mg g⁻¹ DW) were higher than in Franquette fruits (350 mg g⁻¹ DW) for trees located in permeable soils. This could also explain why Franquette is more sensitive to necrosis than Parisienne, especially in permeable soils.

The presence or the absence of copper treatment induced a reverse behaviour in the two cultivars as far as the amount of caffeic acid in the husk tissues of their fruit was concerned (*figure 2d*). In permeable soils, Franquette (9 mg g⁻¹ DW) has a higher content of caffeic acid than Parisienne (7.2 mg g⁻¹

DW); in poorly permeable soils, the reverse situation is found.

Differences in caffeic acid contents between apical tissues (10 mg g⁻¹ DW) and parietal tissues (6.9 mg g⁻¹ DW) are higher for the Parisienne cultivar (*figure 2e*).

On the other hand, the presence of copper treatment influences caffeic acid contents between permeable (9 mg g⁻¹ DW) and poorly permeable (7 mg g⁻¹ DW) soils. Without any copper treatment, the caffeic acid content in husk tissues is roughly the same whatever the soil permeability (*figure 2f*). This increased level of caffeic acid may indicate a stress due to copper. In this way, flavane content is more influenced by presence of copper treatment in the Parisienne cultivar than in the Franquette cultivar (*figure 2g*). Likewise, flavane content was higher for the Franquette cultivar (8.9 mg g⁻¹ DW) than for the Parisienne cultivar (7 mg g⁻¹ DW) in poorly permeable soils, and roughly the same in permeable soils (*figure 2h*).

All of these effects indicated that the nature of the soil certainly modified polyphenol content in walnut tissues. As polyphenols are involved in necrosis pathways as well as resistance, we can hypothesise that the nature of the soil influences the expression of walnut blight through biochemical mechanisms involving polyphenols.

5. CONCLUSION

Even if some questions are not clarified yet, this study has shown that the nature of the soil, especially its permeability, induces variations of polyphenol contents in walnut husk tissues. Likewise, differences observed in apical tissues may explain why it is the apex of the fruit which is chiefly damaged by necrosis. Differences in sensitivity to walnut blight between the two cultivars studied, Franquette and Parisienne, may also be explained by different behaviours in relation to polyphenols. The use of copper for the control of walnut blight is more and more questioned, the accumulation of copper is alarming in many orchards, and this study has shown that copper in the soil induces disturbances in polyphenols.

REFERENCES

- [1] Agrios G.N., Plant Pathology, Academic Press, San Diego, 1997.
- [2] Binder R.G., Benson M.E., Flath R.A., Eight 1,4-naphthoquinones from *Juglans*, *Phytochemistry* 28 (1989) 2799–2801.
- [3] Borazjani A., Graves C.H., Hedin P.A., Occurrence of juglone in various tissues of pecan and related species, *Phytopathology* 75 (1985) 1419–1421.
- [4] Brault T., Étude de la résistance au cuivre chez *Xanthomonas campestris* pv. *juglandis* sur noyer, DEA, Université de Rennes, Inra, Angers, 1990.
- [5] Chapouille P., Planification et analyse des expériences, Masson, Paris, 1973.
- [6] Charlot G., Penet C., Compte rendu des travaux de recherche et expérimentations entrepris à la station de Chatte en 1988, Rapport CTIFL 1988, Station de Chatte (1988) 122–123.
- [7] Delas J., La toxicité du cuivre accumulé dans les sols, *Agrochimica* 7 (1963) 258–287.
- [8] Garavel L., La culture du noyer, Baillière et fils, Paris, 1959.
- [9] Gardan L., Brault T., Germain E., Copper resistance of *Xanthomonas campestris* pv. *juglandis* in French walnut orchards and its association with conjugative plasmids, *Acta Hort.* 311 (1993) 117–124.
- [10] Goupy J., La méthode des plans d'expérience, Dunod, Paris, 1988.
- [11] Harborne J.B., Plant phenolics, in: Bell E.A., Charlwood B.V. (Eds.), *Encyclopedia of Plant Physiology, Secondary Plant Products*, vol. 8, Springer Verlag, Berlin, 1980, pp. 329–402.
- [12] Hedin P.A., Langhans V.E., Graves C.H., Identification of juglone in pecan as a possible factor of resistance to *Fusicladium effusum*, *J. Agric. Food Chem.* 27 (1979) 92–94.
- [13] Hedin P.A., Collum D.H., Langhans V.E., Graves C.H., Distribution of juglone and related compounds in pecan and their effect on *Fusicladium effusum*, *J. Agric. Food Chem.* 28 (1980) 340–342.
- [14] Heijl A.M., Einhellig F.A., Rasmusen J.A., Effect of juglone on growth, photosynthesis and respiration, *J. Chem. Ecol.* 19 (1993) 559–568.
- [15] Jay-Allemand C., Les marqueurs biochimiques de la juvénilité chez le noyer (*Juglans nigra* et *Juglans nigra* × *Juglans regia*), Ph.D. thesis, Université des sciences et techniques du Languedoc, Montpellier, 1985.
- [16] Lagrimini L.M., Vaughn J., Erb W.A., Miller S.A., Peroxidase overproduction in tomato: wound-induced polyphenol deposition and disease resistance, *Hortscience* 28 (1993) 218–221.
- [17] Lee K.C., Campbell R.W., Nature and occurrence of juglone in *Juglans nigra* L., *Hort. Sci.* 4 (1969) 297–298.
- [18] Leistner E., Biosynthesis of plant quinones, in: Stumpf P.K., Conn E.E. (Eds.), *The Biochemistry of the Plants: A Comprehensive Treatise, Secondary Plant Products*, vol. 7, Academic Press, New York, 1981, pp. 403–423.
- [19] Mulrean E.N., Schroth M.N., Bacterial blight on Persian walnuts, *California Agric.* (1981) 11–13.
- [20] Nicholson R.L., Hammerschmidt R., Phenolic compounds and their role in disease resistance, *Annu. Rev. Phytopathol.* 30 (1992) 369–389.

[21] Radix P., Seigle-Murandi F., Hazards of copper accumulation in soils of walnut orchards, *Fresenius Environ. Bull.* 2 (1993) 565–569.

[22] Rastoin F., *La noix, son avenir. Les tendances de la consommation européenne*, Colloque : La noix, réussir sa qualité, Creysse, 17.09.1993, eds CTIF, Paris.

[23] Rhodes J.M., Wooltorton L.S.C., The biosynthesis of phenolic compounds in wounded plant storage tissues, in: Kahl G. (Ed.), *Biochemistry of Wounded Plant*

Tissues, Springer-Verlag, Berlin, New York, 1978, pp. 248–276.

[24] Salgare S.A., Asthana S., Copper an environmental pollutant, *Environ. Ser.* 3 (1990) 101–119.

[25] Siebert J.B., Economic and policy considerations in marketing walnuts, *Acta Hortic.* 311 (1993) 249–257.

[26] Westfall B.A., Russell R.L., Auyong T.K., Depressant agent from walnut hulls, *Science* 134 (1961) 1517.