



**HAL**  
open science

# **Listeria monocytogenes Scott A: cell surface charge, hydrophobicity, and electron donor and acceptor characteristics under different environmental growth conditions**

Romain Briandet, Thierry Meylheuc, Catherine Maher, Marie-Noëlle Bellon-Fontaine

► **To cite this version:**

Romain Briandet, Thierry Meylheuc, Catherine Maher, Marie-Noëlle Bellon-Fontaine. *Listeria monocytogenes* Scott A: cell surface charge, hydrophobicity, and electron donor and acceptor characteristics under different environmental growth conditions. *Applied and Environmental Microbiology*, 1999, 65 (12), pp.5328-5333. hal-02698421

**HAL Id: hal-02698421**

**<https://hal.inrae.fr/hal-02698421v1>**

Submitted on 1 Jun 2020

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

# *Listeria monocytogenes* Scott A: Cell Surface Charge, Hydrophobicity, and Electron Donor and Acceptor Characteristics under Different Environmental Growth Conditions

ROMAIN BRIANDET,<sup>1,2\*</sup> THIERRY MEYLHEUC,<sup>1</sup> CATHERINE MAHER,<sup>3</sup>  
AND MARIE NOËLLE BELLON-FONTAINE<sup>1</sup>

*Institut National de la Recherche Agronomique, Unité de Recherche en Bioadhésion et Hygiène des Matériaux, Massy,<sup>1</sup>  
and SKW Biosystems, Cultures & Enzymes Department, La Ferté sous Jouarre,<sup>2</sup> France, and University College,  
Cork, Ireland<sup>3</sup>*

Received 9 November 1998/Accepted 15 September 1999

**We determined the variations in the surface physicochemical properties of *Listeria monocytogenes* Scott A cells that occurred under various environmental conditions. The surface charges, the hydrophobicities, and the electron donor and acceptor characteristics of *L. monocytogenes* Scott A cells were compared after the organism was grown in different growth media and at different temperatures; to do this, we used microelectrophoresis and the microbial adhesion to solvents method. Supplementing the growth media with glucose or lactic acid affected the electrical, hydrophobic, and electron donor and acceptor properties of the cells, whereas the growth temperature (37, 20, 15, or 8°C) primarily affected the electrical and electron donor and acceptor properties. The nonlinear effects of the growth temperature on the physicochemical properties of the cells were similar for cells cultivated in two different growth media, but bacteria cultivated in Trypticase soy broth supplemented with 6 g of yeast extract per liter (TSYE) were slightly more hydrophobic than cells cultivated in brain heart infusion medium ( $P < 0.05$ ). Adhesion experiments conducted with *L. monocytogenes* Scott A cells cultivated in TSYE at 37, 20, 15, and 8°C and then suspended in a sodium chloride solution ( $1.5 \times 10^{-1}$  or  $1.5 \times 10^{-3}$  M NaCl) confirmed that the cell surface charge and the electron donor and acceptor properties of the cells had an influence on their attachment to stainless steel.**

*Listeria monocytogenes* is a gram-positive human pathogen that is responsible for serious infections in immunocompromised individuals and pregnant women (17). *L. monocytogenes* has been implicated in several food-borne disease outbreaks. This organism is found not only in raw food but also on working surfaces in food-processing plants (40, 41). Microorganisms attached to a surface are an important potential source of contamination for any food material coming into contact with that surface (27). Bacterial attachment to an inert surface results from complex physicochemical interactions among the cell, the surface, and the liquid phase (24), which are caused by the cell surface charge (15), the hydrophobicity (45), and electron acceptor and donor properties (47).

Many workers have described the effects of various environmental parameters on *L. monocytogenes* growth (1, 7, 13), survival (31, 32, 34), pathogenicity (14, 44), and adhesion (2, 24). The most frequently studied parameters are growth temperature (12, 39), acidification of the growth media with organic acids (20, 43), and changes in the water activity of the growth medium caused by adding NaCl (29, 33). It has also been shown that results may be dependent on the growth medium (16, 21, 25). For *L. monocytogenes* growth, the following two media have been used most frequently previously: brain heart infusion (BHI) broth (4, 30) and Trypticase soy broth supplemented with 6 g of yeast extract per liter (TSYE) (5, 22, 26).

Limited data concerning the effects of environmental and cultivation parameters on the physicochemical properties of *L.*

*monocytogenes* have been published previously. Hence, the aim of this study was to determine the surface electrical properties, hydrophobicities, and electron donor and acceptor properties of *L. monocytogenes* Scott A cells under different growth conditions.

## MATERIALS AND METHODS

**Bacterial strains, growth conditions, and preparation of microbial suspension.** *L. monocytogenes* Scott A (a human isolate obtained from the 1983 Massachusetts milk outbreak [18]) was provided by Bongrain Research and Development Centre (SOREDAB, La Boissière Ecole, France) and was stored at  $-80^{\circ}\text{C}$ .

The following four growth media were used for experiments carried out at  $37^{\circ}\text{C}$ : BHI medium (Oxoid, Dardilly, France), TSYE (Biomérieux, Marcy l'Etoile, France), TSYE supplemented with 7.5 g of glucose per liter (Prolabo, Fontenay-sous-Bois, France), and TSYE supplemented with 1 N lactic acid (Prolabo) to obtain a pH of 6.0. The effects of growth temperatures (37, 20, 15, and  $8^{\circ}\text{C}$ ) were studied by using TSYE and BHI medium. For all of the experiments, frozen cells were subcultured twice in the same medium and at the same growth temperature as the final culture. For the final culture, 1 ml of the second subculture was inoculated into 200 ml of fresh medium, and the preparation was incubated at the appropriate temperature until the stationary stage was reached. The stationary stage was reached after 15, 24, 30, and 192 h for cells grown at 37, 20, 15, and  $8^{\circ}\text{C}$ , respectively.

Cells were harvested by centrifugation for 10 min at  $7,000 \times g$  at  $4^{\circ}\text{C}$  and were washed twice with and resuspended in the relevant suspending liquid ( $1.5 \times 10^{-1}$  or  $1.5 \times 10^{-3}$  M NaCl).

**MATS.** Microbial adhesion to solvents (MATS) is based on comparing microbial cell affinity to a polar solvent and microbial cell affinity to a nonpolar solvent (8). The polar solvent can be an electron acceptor or an electron donor, but both solvents must have similar van der Waals surface tension components. The following pairs of solvents, as described by Bellon-Fontaine et al. (8), were used: chloroform, an electron acceptor solvent, and hexadecane, a nonpolar solvent; and ethyl acetate, a strong electron donor solvent, and decane, a nonpolar solvent. Due to the surface tension properties of these solvents, differences between the results obtained with chloroform and hexadecane and the results obtained with ethyl acetate and decane indicated that there were electron donor-

\* Corresponding author. Mailing address: INRA UBHM, 25 Avenue République, 91300 Massy, France. Phone: 33 1 69 53 64 00. Fax: 33 1 60 13 36 01. E-mail: briandet@massy.inra.fr.

TABLE 1. Affinities of *L. monocytogenes* Scott A cells for the four solvents used in the MATS analysis after growth in either BHI medium or TSYE at 37, 20, 15, and 8°C and after growth at 37°C in TSYE supplemented with glucose or lactic acid<sup>a</sup>

Medium	Temp (°C)	% Affinity for:			
		Chloroform	Hexadecane	Ethyl acetate	Decane
TSYE + glucose	37	90 ± 1	70 ± 6	56 ± 3	71 ± 9
TSYE + lactic acid	37	96 ± 1	85 ± 7	33 ± 3	96 ± 8
TSYE	37	91 ± 7	46 ± 11	17 ± 9	51 ± 11
	20	94 ± 6	48 ± 10	17 ± 2	64 ± 6
	15	91 ± 3	48 ± 10	7 ± 3	66 ± 7
	8	94 ± 2	53 ± 8	14 ± 4	67 ± 11
BHI	37	89 ± 3	41 ± 11	24 ± 5	44 ± 10
	20	92 ± 3	48 ± 7	13 ± 5	47 ± 5
	15	92 ± 4	39 ± 11	13 ± 2	64 ± 3
	8	93 ± 4	44 ± 4	17 ± 1	55 ± 6

<sup>a</sup> Affinity values are means ± standard deviations ( $n = 9$ ).

electron acceptor interactions at the bacterial cell surface and revealed hydrophobic and hydrophilic properties. A microbial suspension containing approximately  $10^8$  CFU in 2.4 ml of  $1.5 \times 10^{-1}$  M NaCl was vortexed for 60 s with 0.4 ml of a solvent. This high concentration of electrolyte was used to avoid charge interference by a masking cell charge, because some solvent droplets, especially hexadecane, become negatively charged in aqueous suspensions (19). The mixture was allowed to stand for 15 min to ensure that the two phases were completely separated before a sample (1 ml) was carefully removed from the aqueous phase and the optical density at 400 nm was determined. The percentage of cells present in each solvent was subsequently calculated by using the equation: % Affinity =  $100 \times [1 - (A/A_0)]$ , where  $A_0$  is the optical density at 400 nm of the bacterial suspension before mixing and  $A$  is the absorbance after mixing. Each experiment was performed in triplicate by using three independently prepared cultures.

**Electrophoretic mobility.** To measure electrophoretic mobility, bacteria were suspended in  $1.5 \times 10^{-3}$  M sodium chloride at a concentration of  $10^7$  CFU · ml<sup>-1</sup>. The pH of the suspension was adjusted to pH 2 to 7 by adding nitric acid (HNO<sub>3</sub>) or potassium hydroxide (KOH). Electrophoretic mobility was measured by using a 50-V electric field and a Laser Zetameter (Zétaphoremètre II; Société d'Étude Physico-Chimiques, Limours, France). The results were based on an automated video analysis of about 200 particles for each measurement. Each experiment was performed in duplicate by using two independently prepared cultures. The typical standard deviation for the electrophoretic mobility mean was 0.25 μm/V/cm/s.

**Microbial adhesion to AISI 304 stainless steel. (i) Solid surface and cleaning treatment.** The solid support selected for this study was AISI 304 stainless steel (Goodfellow, Cambridge Science Park, United Kingdom). Before physicochemical characterization and adhesion assays were begun, the steel was cut into rectangular chips (3 by 1 cm) and cleaned by soaking for 10 min at 50°C in a 2% (vol/vol) solution of the commercial surfactant RBS 35 (Société des Traitements Chimiques de Surface, Lambertsart, France), rinsing for 10 min in Milli-Q water (Millipore, Saint-Quentin en Yvelines, France) at 50°C, and rinsing five times in 500 ml of Milli-Q water at room temperature.

**(ii) Determination of the physicochemical properties of the solid surface.** The Lifshitz-van der Waals ( $\gamma^{LW}$ ), electron donor ( $\gamma^+$ ), and electron acceptor ( $\gamma^-$ ) surface tension components of stainless steel ( $S$ ) were determined by measuring contact angles by using the approach proposed by van Oss et al. (46). In this approach, in which spreading pressure is ignored, the contact angles ( $\theta$ ), measured with three pure liquids ( $L$ ), can be expressed as:

$$\cos\theta = -1 + 2(\gamma_s^{LW}\gamma_L^{LW})^{1/2}/\gamma_L + 2(\gamma_s^+\gamma_L^-)^{1/2}/\gamma_L + 2(\gamma_s^-\gamma_L^+)^{1/2}/\gamma_L$$

The three pure liquids used were Milli-Q water (Millipore), formamide (Sigma, Saint-Quentin Fallanier, France), and diiodomethane (Sigma).

**(iii) Adhesion experiments.** Adhesion assays were performed by using microbial cells cultured at 37, 20, 15, or 8°C in TSYE and resuspended in  $1.5 \times 10^{-1}$  or  $1.5 \times 10^{-3}$  M sodium chloride as described below. Thirty milliliters of a bacterial suspension containing approximately  $10^8$  CFU · ml<sup>-1</sup> was incubated in a petri dish (diameter, 10 cm) containing 3-cm<sup>2</sup> stainless steel chips for 3 h at the temperature at which the bacteria were grown (37, 20, 15, or 8°C). The stainless steel chips were then rinsed to remove the nonadhering bacteria by pouring 30 ml of Milli-Q water onto the chips three times. The stainless steel chips were immersed in a test tube containing 10 ml of sterile  $1.5 \times 10^{-1}$  or  $1.5 \times 10^{-3}$  M NaCl. Bacterial cells were detached from the inert support by using a sonication bath (Ultrasonik) for 2 min at 40 kHz and 35°C. CFUs were counted by using the serial dilution technique and the bacterial suspension obtained after sonication. Counts were determined on TSYE agar (Biomérieux), and the preparations were incubated for 24 h at 37°C. Each experiment was performed in triplicate by using two independently grown cultures. We assessed the viability of the bacteria in the

two suspending liquids and at the four temperatures analyzed by counting suspended cells on TSYE agar at the beginning and end of the adhesion period.

**Statistical analysis.** A principal-component analysis was performed by using STATITCF software (Institut Technique des Céréales et des Fourrages, Paris, France), and a two-way analysis of variance was performed with Statgraphics 6.0 (Manugistics, Rockville, Md.).

## RESULTS

**Influence of growth conditions (medium and temperature) on *L. monocytogenes* Scott A surface physicochemical properties. (i) Influence of growth medium on cell surface physicochemical properties.** The MATS results obtained for *L. monocytogenes* Scott A grown at 37°C in BHI, TSYE, TSYE supplemented with glucose, and TSYE supplemented with lactic acid are shown in Table 1. Regardless of the medium used, the affinity of *L. monocytogenes* Scott A was always higher with chloroform (an electron acceptor solvent) than with hexadecane (a nonpolar solvent). The differences in bacterial affinity between these two solvents were due to Lewis acid-base interactions (i.e., electron donor-electron acceptor interactions resulting from the electron donor nature of the bacteria). The electron donor nature was maximized for cells cultivated in BHI medium or TSYE. Likewise, bacterial affinity was lower with ethyl acetate (a strongly electron-donating solvent) than with decane, indicating that the electron-accepting nature of *L. monocytogenes* Scott A grown in either medium was weak. Affinity to ethyl acetate was maximized with cells cultivated in the glucose-supplemented medium; this culture was also the culture with the lowest pH in the stationary phase (pH 4.1 and 4.7 for culture media supplemented with glucose and lactic acid, respectively; pH 5.1 for cells grown in TSYE).

Figure 1 shows the electrophoretic mobilities of *L. monocytogenes* Scott A at pH 2 to 7 when it was grown in BHI medium, TSYE, TSYE supplemented with glucose, or TSYE supplemented with lactic acid. All of the cells were highly negatively charged, and so an isoelectric point could not be determined in the pH range studied. Conversely, cells grown in TSYE or BHI medium had very similar electrical characteristics (high mobility at a neutral pH and low mobility at an acidic pH) compared with cells cultivated in medium supplemented with glucose or lactic acid.

**(ii) Influence of growth temperature on cell surface physicochemical properties.** Table 1 shows the affinities with the four solvents used in the MATS method of the bacteria grown in TSYE or BHI medium at 37, 20, 15, and 8°C. Cells cultivated in TSYE exhibited slightly greater affinity with nonpolar solvents than cells cultivated in BHI medium exhibited ( $P < 0.05$ ), and the affinity with ethyl acetate was greatest at 37°C




FIG. 1. Electrophoretic mobilities of *L. monocytogenes* Scott A cells suspended in  $1.5 \times 10^{-3}$  M NaCl at pH 2 to 7 after growth at 37°C in BHI, TSYE, TSYE supplemented with glucose, or TSYE supplemented with lactic acid.

and least at 15°C ( $P < 0.05$ ). Moreover, the growth temperature had a significant effect ( $P < 0.05$ ) on the electrophoretic mobility of the bacteria, as shown in Fig. 2. Our results indicated that the nonlinear effects of the growth temperature on the electrophoretic mobility of microbial cells were the same for both growth media. Except for the most acidic pH (pH 2.0), electrophoretic mobility was greatest for cells grown at 15 or 20°C and lowest for cells cultivated at 8°C. The electrophoretic mobility of the bacteria was between  $-0.5$  and  $-2 \mu\text{m}^2/\text{s}/\text{V}/\text{cm}$  at pH 2.0. The isoelectric point of microbial cells could not be determined at any of the growth temperatures used in this study. Data were analyzed by using the data compression step of principal-component analysis, which removes the redundancy in an original data set so that only the first few principal-component scores are needed to describe most of the information in the original data set (11). The results of a principal-component analysis of the combined physicochemical properties of *L. monocytogenes* Scott A grown in TSYE or BHI medium at all temperatures are shown in Fig. 3. The first principal-component score, which accounts for 43% of the total variation in the original data set, is positively correlated with a high negative cell wall charge. This principal-component score discriminates among the samples by relating to the growth temperature. The second principal-component score represents 29% of the total variation in the original data set, and it is positively correlated with cell affinity for nonpolar solvents (hexadecane and decane) and negatively correlated with cell affinity for ethyl acetate. This principal-component score mainly distinguishes between cells grown in BHI and cells grown in TSYE.

**Adhesion of *L. monocytogenes* Scott A grown in TSYE at different temperatures.** The numbers of culturable bacteria adhering to stainless steel under the eight conditions studied here are shown in Table 2. At the four growth temperatures, bacterial cells bound more effectively to the stainless steel when they were suspended in  $1.5 \times 10^{-1}$  M NaCl than when they were suspended in  $1.5 \times 10^{-3}$  M NaCl ( $P < 0.05$ ). Furthermore, adhesion was greatest for bacterial cells cultivated at




FIG. 2. Electrophoretic mobilities of *L. monocytogenes* Scott A cells suspended in  $1.5 \times 10^{-3}$  M NaCl at pH 2 to 7 after growth at four temperatures in BHI medium (A) or TSYE (B).

37°C and lowest for cells cultivated at 15°C ( $P < 0.05$ ), and the number of attached bacteria decreased between 20 and 15°C. A graph of the affinity of cells for ethyl acetate as determined by the MATS method versus the number of adherent cells in the presence of  $1.5 \times 10^{-3}$  M NaCl at the four growth temperatures used is shown in Fig. 4. The linear coefficients of correlation between bacterial affinity for ethyl acetate and bacterial adhesion to stainless steel were 0.93 and 0.80 for adhesion assays performed in the presence of  $1.5 \times 10^{-3}$  and  $1.5 \times 10^{-1}$  M NaCl, respectively.

Differences in adherence were not due to a loss of cell viability during the adhesion experiments, because after 3 h of incubation at 37, 20, 15, or 8°C in the presence of either  $1.5 \times 10^{-1}$  or  $1.5 \times 10^{-3}$  M NaCl, the viable cell counts were not affected ( $P < 0.05$ ).

The Lifshitz-van der Waals, electron donor, and electron acceptor components of the surface free energy of the stainless steel chips, as determined by contact angle measurements, were 39, 13, and  $0.3 \text{ mJ} \cdot \text{m}^{-2}$ , respectively.




FIG. 3. Principal-component analysis of combined physicochemical properties (affinity for the solvents used with the MATS method and electrophoretic mobility of bacteria) of *L. monocytogenes* Scott A grown at different temperatures in TSYE (●) and BHI medium (□). The first principal component (principal component 1) was positively correlated with a high negative charge of bacterial cells. The second principal component (principal component 2) was positively correlated with bacterial cell affinity for nonpolar solvents and negatively correlated with cell affinity for ethyl acetate.

## DISCUSSION

The aim of this study was to investigate the influence of growth conditions (medium and temperature) on the physicochemical surface properties of *L. monocytogenes* Scott A. Microbial cell surfaces were found to be hydrophilic when bacteria were grown at 37°C in BHI medium or TSYE. However, the hydrophilicity decreased when microbial cells were cultivated in TSYE supplemented with either glucose or lactic acid. These findings are consistent with results obtained by Mafu et al. (28), who used the hydrophobic interaction chromatography technique and found that the hydrophobicity of *L. monocytogenes* Scott A increased as the pH decreased.

Furthermore, bacterial cells cultivated in BHI medium and bacterial cells cultivated in TSYE exhibited the same physicochemical differences when they were cultivated at different temperatures, and cells grown in media supplemented with glucose or lactic acid exhibited greater affinity with the electron donor solvent (ethyl acetate), which indicated that more electron acceptor groups were present on the cell wall. The presence of bound acidic compounds on the cell walls of the bacteria could explain this phenomenon. The low pH of a culture grown in a glucose-enriched medium in the stationary phase was undoubtedly related to the metabolism of glucose to lactic or acetic acid (38).

Under all of the growth conditions studied, *L. monocytogenes* cells had a high negative charge, and the electrophoretic mobilities at pH 7 were greater than the value ( $-2.07 \mu\text{m/V}$ )

TABLE 2. Numbers of adherent *L. monocytogenes* Scott A cells on stainless steel chips as determined by adhesion assays performed in the presence of  $1.5 \times 10^{-1}$  or  $1.5 \times 10^{-3}$  M NaCl after growth in TSYE at four different temperatures<sup>a</sup>

NaCl concn (M)	Log no. of adherent cells after growth at:			
	37°C	20°C	15°C	8°C
$1.5 \times 10^{-1}$	$6.4 \pm 1.2$	$5.6 \pm 0.9$	$4.1 \pm 0.8$	$4.4 \pm 0.2$
$1.5 \times 10^{-3}$	$5.2 \pm 1.0$	$4.9 \pm 0.9$	$3.0 \pm 0.3$	$3.9 \pm 0.6$

<sup>a</sup> The values are means  $\pm$  standard deviations ( $n = 6$ ).




FIG. 4. Percentage of bacteria bound to ethyl acetate, a solvent used with the MATS method, versus the number of cells that adhered to stainless steel in the presence of  $1.5 \times 10^{-3}$  M NaCl.

cm/s) obtained for the same strain by Mafu et al. (27). However, the storage conditions (4°C), growth medium (Trypticase soy broth supplemented with 1% yeast extract), and suspending medium (sodium phosphate buffer) were different in the study of Mafu et al.

Bacterial charge is attributed to cell wall constituents (36) (e.g., phosphate, carboxylate groups, and proteins). Progressive reductions in the electrophoretic mobility of cells as the pH decreases are due to gradual increases in the protonation of several chemical groups. Our failure to determine the isoelectric points in the pH range which we used (pH 2 to 7) indicates that compounds with very low  $pK_a$  were present on the cell surfaces. Considering the composition of the *L. monocytogenes* cell wall (42), the characteristics described above might be related to the low  $pK_a$  ( $pK_a < 2.1$ ) of the phosphate groups in phosphodiester bridges (R-O-HPO<sub>2</sub>-O-R/R-O-PO<sub>2</sub><sup>-</sup>-O-R) of cell wall teichoic acids (37).

The electrophoretic mobility of cells cultivated in either one of the supplemented media seemed to be less pH dependent, and the decrease in mobility was more subtle when the pH of the microbial suspension was reduced below pH 4. Supplementing the growth medium with glucose or lactic acid also reduced the electrophoretic mobility of the bacteria at pH values above 4 through direct neutralization of the negatively charged groups present on cell walls linked to a decrease in the pH of the medium.

The growth temperature had a significant nonlinear effect on the electrophoretic mobility of *L. monocytogenes* Scott A cells grown in either TSYE or BHI medium. In particular, changes in electrophoretic mobility were observed between 15 and 8°C; at these temperatures the values decreased dramatically and were less pH dependent, and almost no changes were observed after the microbial suspension was acidified. This difference in activity at low pH values could have been due to differences in the nature of the negative charge of the cells. The rapid reduction in electrophoretic mobility around pH 4 may indicate that protein- or peptidoglycan-associated COOH/COO<sup>-</sup> ( $4 < pK_a < 5$ ) was present on the cell walls (37). Hence, cells grown at 8°C may have had fewer carboxyl groups than cells grown at the other temperatures. Differences observed at this growth

temperature may also have been related to a glycolipid that was isolated from *L. monocytogenes* only after growth at a low temperature (23).

The high negative charge of bacterial cells at 15 and 20°C may have been linked to the presence of temperature-dependent production of flagella by *L. monocytogenes*. Many flagella were observed on cells grown at 20°C, whereas at 37°C very few flagella were observed (35). The rich protein (and protein-associated COOH/COO<sup>-</sup>) content of flagella could explain the specific electric properties of cells cultivated at 15 or 20°C. The differences in physicochemical properties of cells grown in different media and at different temperatures may also have reflected the synthesis of acclimation proteins (6) due to acidic or thermal stresses during bacterial growth, which could have led to changes in cell wall composition.

Unlike van der Waals and Lewis acid-base interactions, electrostatic interactions are inhibited in a high-ionic-strength suspending liquid. The greatest adhesion of microbial cells in a high-ionic-strength suspending liquid (1.5 × 10<sup>-1</sup> M NaCl) indicated that there was electrostatic repulsion between the negatively charged bacteria and the stainless steel. These results are consistent with previously published data which indicated that the isoelectric point of stainless steel was around pH 4 and varied slightly depending on the surface finish and the cleaning treatments (9). The *L. monocytogenes* Scott A adhesion data correlated best with cell affinity for ethyl acetate, which indicates the importance of Lewis acid-base interactions for cell adhesion to stainless steel. This correlation is consistent with the physicochemical characteristics of ethyl acetate and stainless steel, both of which have strong electron donor surface properties and weak electron acceptor characteristics. Hence, our data suggest that both electrical and Lewis acid-base interactions are involved in *L. monocytogenes* Scott A adhesion to stainless steel. The influence of lactic acid in the growth medium on the adhesion of *L. monocytogenes* to stainless steel has been described previously (10). Variations in cell hydrophobicity caused by lactic acid led to significant variations in the adherent bacteria, implying that van der Waals interactions play a role in cell adhesion. Therefore, the growth conditions can influence the various components of microbial cell surface physicochemical properties (electrostatic, Lewis acid-base, and van der Waals interactions) and, consequently, the adhesion of cells to surfaces. Recently, Andersson et al. found that there is a relationship among *Bacillus cereus* spore adhesion to epithelial cells, surface hydrophobicity, and the virulence of strains (3). Adhesion assays performed with different surfaces that are representative of plants or epithelial cells and with *L. monocytogenes* strains whose pathogenicity is known are needed to improve our understanding of the relationships between physicochemical properties of microbial cells and pathogenicity.

#### ACKNOWLEDGMENTS

This work was supported by the UNIR (Ultraprope Nutrition Industrie Recherche) program, which involves food companies, the French Ministry of Agriculture, and the French Ministry of Research.

We thank Michael Dever and Dipak Sarker for revising the English in the manuscript.

#### REFERENCES

- Ahamad, N., and E. H. Marth. 1989. Behavior of *Listeria monocytogenes* at 7, 13, 21, and 35°C in tryptose broth acidified with acetic, citric, or lactic acid. *J. Food Prot.* **52**:688–695.
- Al-Makhlafi, H. 1994. Influence of preadsorbed milk proteins on adhesion of *Listeria monocytogenes* to hydrophobic and hydrophilic silica surfaces. *Appl. Environ. Microbiol.* **60**:3560–3565.
- Andersson, A., P. E. Granum, and U. Rönnér. 1998. The adhesion of *Bacillus cereus* spores to epithelial cells might be an additional virulence mechanism. *Int. J. Food Microbiol.* **39**:93–99.
- Baloga, A. O., and S. K. Harlander. 1991. Comparison of methods for discrimination between strains of *Listeria monocytogenes* from epidemiological surveys. *Appl. Environ. Microbiol.* **57**:2324–2331.
- Barbosa, W. B., L. Cabedo, H. J. Wederquist, J. N. Sofos, and G. R. Schmidt. 1994. Growth variation among species and strains of *Listeria* in culture broth. *J. Food Prot.* **57**:765–775.
- Bayles, D. O., B. A. Annous, and B. J. Wilkinson. 1996. Cold stress proteins induced in *Listeria monocytogenes* in response to temperature downshock and growth at low temperatures. *Appl. Environ. Microbiol.* **62**:1116–1119.
- Begot, C., I. Lebert, and A. Lebert. 1997. Variability of the response of 66 *Listeria monocytogenes* and *Listeria innocua* strains to different growth conditions. *Food Microbiol.* **14**:403–412.
- Bellon-Fontaine, M. N., J. Rault, and C. J. van Oss. 1996. Microbial adhesion to solvents: a novel method to determine the electron-donor/electron-acceptor or Lewis acid-base properties of microbial cells. *Colloids Surfaces B Biointerfaces* **7**:47–53.
- Boulangé-Petermann, L., A. Doren, B. Baroux, and M. N. Bellon-Fontaine. 1995. Zeta potential measurements on passive metals. *J. Colloid Interface Sci.* **171**:179–186.
- Briand, R., V. Leriche, B. Carpentier, and M. N. Bellon-Fontaine. 1999. Effects of the growth conditions on the surface hydrophobicity of *Listeria monocytogenes* cells and their adhesion to stainless steel. *J. Food Prot.* **62**:994–998.
- Briand, R., E. K. Kemsley, and R. H. Wilson. 1996. Approaches to adulteration detection in instant coffees using infrared spectroscopy and chemometrics. *J. Sci. Food Agric.* **71**:359–366.
- Buchanan, R. L., H. G. Stahl, and R. C. Whiting. 1989. Effects and interactions of temperature, pH, atmosphere, sodium chloride, and sodium nitrite on the growth of *Listeria monocytogenes*. *J. Food Prot.* **52**:844–851.
- Buchanan, R. L., and J. G. Phillips. 1990. Response surface model for predicting the effects of temperature, pH, sodium chloride content, sodium nitrite concentration and atmosphere on the growth of *Listeria monocytogenes*. *J. Food Prot.* **53**:370–376.
- Buncic, S., and S. M. Avery. 1996. Relationship between variations in pathogenicity and lag phase at 37°C of *Listeria monocytogenes* previously stored at 4°C. *Lett. Appl. Microbiol.* **23**:18–22.
- Dickson, J. S., and M. Koochmaria. 1989. Cell surface charge characteristics and their relationship to bacterial attachment to meat surfaces. *Appl. Environ. Microbiol.* **55**:832–836.
- Duffy, G., and J. J. Sheridan. 1997. The effect of temperature, pH and medium in a surface adhesion immunofluorescent technique for detection of *Listeria monocytogenes*. *J. Appl. Microbiol.* **83**:95–101.
- Farber, J. M., and P. I. Peterkin. 1991. *Listeria monocytogenes*, a food-borne pathogen. *Microbiol. Rev.* **55**:476–511.
- Fleming, J. M., S. L. Cochi, K. L. MacDonald, J. Brondum, P. S. Hayes, B. D. Plikaytis, M. B. Holmes, A. Audurier, C. V. Broome, and A. L. Reingold. 1985. Pasteurized milk as a vehicle of infection in an outbreak of listeriosis. *N. Engl. J. Med.* **312**:404–407.
- Geertsema-Doornbush, G. L., H. C. Van der Mei, and H. J. Busscher. 1993. Microbial cell surface hydrophobicity: the involvement of electrostatic interactions in microbial adhesion to hydrocarbons (MATH). *J. Microbiol. Methods* **18**:61–68.
- George, S. M., B. M. Lund, and T. F. Brocklehurst. 1988. The effect of pH and temperature on initiation of growth of *Listeria monocytogenes*. *Lett. Appl. Microbiol.* **6**:153–156.
- Horska, E., J. Pokorný, and M. Labajova. 1993. Changes of surface charge and hydrophobicity of the outer bacterial membrane depending on the cultivation medium. *Biologia (Bratislava)* **48**:343–347.
- Ita, P. S., and R. W. Hutkins. 1991. Intracellular pH and survival of *Listeria monocytogenes* Scott A in tryptic soy broth containing acetic, lactic, citric, and hydrochloric acids. *J. Food Prot.* **54**:15–19.
- Jones, C. E., G. Shama, D. Jones, I. S. Roberts, and P. W. Andrew. 1997. Physiological and biochemical studies on psychrotolerance in *Listeria monocytogenes*. *J. Appl. Microbiol.* **83**:31–35.
- Kim, K. Y., and J. F. Frank. 1994. Effect of growth nutrients on attachment of *Listeria monocytogenes* to stainless steel. *J. Food Prot.* **57**:720–726.
- Kim, K. Y., and J. F. Frank. 1995. Effect of nutrients on biofilm formation by *Listeria monocytogenes* on stainless steel. *J. Food Prot.* **58**:24–28.
- Kouassi, Y., and L. A. Shelef. 1996. Metabolic activities of *Listeria monocytogenes* in the presence of sodium propionate, acetate, lactate and citrate. *J. Appl. Bacteriol.* **81**:147–153.
- Mafu, A. A., D. Roy, J. Goulet, and P. Magny. 1990. Attachment of *Listeria monocytogenes* to stainless steel, glass, polypropylene, and rubber surfaces after short contact times. *J. Food Prot.* **53**:742–746.
- Mafu, A. A., D. Roy, J. Goulet, and L. Savoie. 1991. Characterization of physicochemical forces involved in adhesion of *Listeria monocytogenes* to surfaces. *Appl. Environ. Microbiol.* **57**:1969–1973.
- McClure, P. J., A. L. Beaumont, J. P. Sutherland, and T. A. Roberts. 1996. Predictive modelling of growth of *Listeria monocytogenes*. The effects on

- growth of NaCl, pH, storage temperature and NaNO<sub>2</sub>. *Int. J. Food Microbiol.* **34**:221–232.
30. **McLauchlin, J., A. M. Tidley, and A. G. Taylor.** 1989. The use of monoclonal antibodies against *Listeria monocytogenes* in a direct immunofluorescence technique for the rapid presumptive identification and direct demonstration of *Listeria* in food. *Acta Microbiol. Hung.* **36**:467–471.
  31. **Oh, D.-H., and D. L. Marshall.** 1993. Influence of temperature, pH, and glycerol monolaurate on growth and survival of *Listeria monocytogenes*. *J. Food Prot.* **56**:744–749.
  32. **Palumbo, S. A., and A. C. Williams.** 1990. Effect of temperature, relative humidity, and suspending menstrue on the resistance of *Listeria monocytogenes* to drying. *J. Food Prot.* **53**:377–381.
  33. **Papageorgiou, D. K., and E. H. Marth.** 1989. Behaviour of *Listeria monocytogenes* at 4 and 22°C in whey and skim milk containing 6 or 12% sodium chloride. *J. Food Prot.* **52**:625–630.
  34. **Patchett, R. A., N. Watson, P. S. Fernandez, and R. G. Kroll.** 1996. The effect of temperature and growth rate on the susceptibility of *Listeria monocytogenes* to environmental stress conditions. *Appl. Microbiol.* **22**:121–124.
  35. **Peel, M., W. Donachie, and A. Shaw.** 1988. Temperature-dependent expression of flagella of *Listeria monocytogenes* studied by electron microscopy, SDS-PAGE and Western blotting. *J. Gen. Microbiol.* **134**:2171–2178.
  36. **Pelletier, C., C. Bouley, C. Cayvela, S. Boutier, P. Bourlioux, and M. N. Bellon-Fontaine.** 1997. Cell surface characteristics of *Lactobacillus casei* subsp. *casei*, *Lactobacillus paracasei* subsp. *paracasei*, and *Lactobacillus rhamnosus* strains. *Appl. Environ. Microbiol.* **63**:1725–1731.
  37. **Rijnaarts, H. H. M., W. Norbe, J. Lyklema, and A. J. B. Zehnder.** 1995. The isoelectric point of bacteria as an indicator for the presence of cell surface polymers that inhibit adhesion. *Colloids Surf. B Biointerfaces* **4**:191–197.
  38. **Romick, T. L., and H. P. Fleming.** 1998. Acetoin production as an indicator of growth and metabolic inhibition of *Listeria monocytogenes*. *J. Appl. Microbiol.* **84**:18–24.
  39. **Rosenow, E. M., and E. H. Marth.** 1987. Growth of *Listeria monocytogenes* in skim, whole and chocolate milk, and in whipping cream during incubation at 4, 8, 13, 21 and 35°C. *J. Food Prot.* **50**:452–459.
  40. **Roy, B., H. W. Ackermann, S. Pandian, G. Picard, and G. Goulet.** 1993. Biological inactivation of adhering *Listeria monocytogenes* by listeria phages and a quaternary ammonium compound. *Appl. Environ. Microbiol.* **59**:2914–2917.
  41. **Sasahara, K. C., and E. A. Zottola.** 1993. Biofilm formation by *Listeria monocytogenes* utilizes a primary colonizing microorganism in flowing systems. *J. Food Prot.* **56**:1022–1028.
  42. **Seeliger, H. P. R., and D. Jones.** 1986. *Listeria*, p. 1235–1244. In P. H. A. Sneath, N. S. Mair, M. E. Sharpe, and J. G. Holt (ed.), *Bergey's manual of systematic bacteriology*, vol. 2. Williams & Wilkins, Baltimore, Md.
  43. **Sorrells, K. M., D. C. Enigl, and J. R. Hatfield.** 1989. Effect of pH, acidulant, time, and temperature on the growth and survival of *Listeria monocytogenes*. *J. Food Prot.* **52**:571–573.
  44. **Stephens, J. C., I. S. Roberts, D. Jones, and P. W. Andrew.** 1991. Effect of growth temperature on virulence of strains of *Listeria monocytogenes* in the mouse: evidence for a dose dependence. *J. Appl. Bacteriol.* **70**:239–244.
  45. **van Loosdrecht, M. C. M., J. Lyklema, W. Norde, G. Schraa, and A. J. B. Zehnder.** 1987. The role of bacterial cell wall hydrophobicity in adhesion. *Appl. Environ. Microbiol.* **53**:1893–1897.
  46. **van Oss, C. J., M. K. Chaudhury, and R. J. Good.** 1988. Interfacial Lifshitz-van der Waals and polar interactions in macroscopic systems. *Chem. Rev.* **88**:927–941.
  47. **van Oss, C. J.** 1993. Acid-base interfacial interactions in aqueous media. *Colloids Surf. A Physicochem. Eng. Aspects* **78**:1–49.