



**HAL**  
open science

## **Cyclin F regulates the nuclear localization of cyclin B1 through a cyclin-cyclin interaction**

Monica Kong, Elizabeth A Barnes, Vincent Ollendorff, Daniel J. Donoghue

► **To cite this version:**

Monica Kong, Elizabeth A Barnes, Vincent Ollendorff, Daniel J. Donoghue. Cyclin F regulates the nuclear localization of cyclin B1 through a cyclin-cyclin interaction. *EMBO Journal*, 2000, 19 (6), pp.1378-1388. 10.1093/emboj/19.6.1378 . hal-02698812

**HAL Id: hal-02698812**

**<https://hal.inrae.fr/hal-02698812>**

Submitted on 1 Jun 2020

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# Cyclin F regulates the nuclear localization of cyclin B1 through a cyclin–cyclin interaction

Monica Kong, Elizabeth A. Barnes,  
Vincent Ollendorff<sup>1</sup> and Daniel J. Donoghue<sup>2</sup>

Department of Chemistry and Biochemistry, Center for Molecular Genetics, University of California, San Diego, La Jolla, CA 92093-0367, USA

<sup>1</sup>Present address: INSERM U119, 27 Boulevard Lei Roure, 13009 Marseille, France

<sup>2</sup>Corresponding author  
e-mail: ddonoghue@ucsd.edu

**The key regulator of G<sub>2</sub>–M transition of the cell cycle is M-phase promoting factor (MPF), a complex composed of cdc2 and a B-type cyclin. Cyclin B1 nuclear localization involves phosphorylation within a region called the cytoplasmic retention signal, which also contains a nuclear export signal. The mechanism of MPF nuclear localization remains unclear since it contains no functional nuclear localization signal (NLS). We exploited the yeast two-hybrid screen to find protein(s) potentially mediating localization of cyclin B1 and identified a novel interaction between cyclin B1 and cyclin F. We found that cdc2, cyclin B1 and cyclin F form a complex that exhibits histone H1 kinase activity. Cyclin B1 and cyclin F also colocalize through immunofluorescence studies. Additionally, deletion analysis revealed that each putative NLS of cyclin F is functional. Taken together, the data suggest that the NLS regions of cyclin F regulate cyclin B1 localization to the nucleus. The interaction between cyclin B1 and cyclin F represents the first example of direct cyclin–cyclin binding, and elucidates a novel mechanism that regulates MPF localization and function.**

**Keywords:** cyclin B1/cyclin F/cytoplasmic retention signal/M-phase promoting factor/nuclear localization signal

## Introduction

The onset of mitosis is regulated by the activation of M-phase promoting factor (MPF) in both mitotic and meiotic cells (Draetta and Beach, 1988; Westendorf *et al.*, 1989; Nurse, 1990; Hunt, 1991). MPF consists of two subunits, the serine/threonine kinase cdc2 and either cyclin B1 or cyclin B2 (Labbe *et al.*, 1989; Draetta *et al.*, 1989; Gautier *et al.*, 1990). Cyclins were first identified as proteins whose levels oscillate during the embryonic cell cycle in invertebrate eggs (Evans *et al.*, 1983). Cyclin B1 protein begins to accumulate during late S and G<sub>2</sub> phases, peaks at late G<sub>2</sub>–M phase, begins to degrade at the start of metaphase, and is nearly completely degraded at the onset of anaphase (Clute and Pines, 1999). During interphase, cyclin B1 associates with its catalytic partner

cdc2 through a conserved region of ~150 amino acids known as the cyclin box (Minshull *et al.*, 1989; Pines and Hunter, 1989; Westendorf *et al.*, 1989; Nugent *et al.*, 1991; Kobayashi *et al.*, 1992). Most cyclins interact with a cyclin-dependent kinase (cdk) through the cyclin box, and to date, cyclins have not been found to interact directly with one another.

In late G<sub>2</sub> phase, post-translational modification of an inactive cdc2–cyclin B1 complex occurs, involving both phosphorylation and dephosphorylation of tyrosine and threonine residues of cdc2 (Solomon *et al.*, 1990; Krek and Nigg, 1991). cdc2 kinase activity is negatively regulated by phosphorylation at Thr14 by the Myt1 kinase and at Tyr15 by the Wee1/Mik1 or Myt1 protein kinases (Russell and Nurse, 1987; Gould and Nurse, 1989; Parker *et al.*, 1992; Kornbluth *et al.*, 1994; Mueller *et al.*, 1995), and positively regulated by phosphorylation at Thr161 by cdc2-activating kinase (Gould *et al.*, 1991; Solomon *et al.*, 1992; Fesquet *et al.*, 1993; Poon *et al.*, 1993). Dephosphorylation of Thr14 and Tyr15 is controlled by the cdc25 phosphatase (Gautier *et al.*, 1991; Millar *et al.*, 1991; Honda *et al.*, 1993), and in conjunction with Thr161 phosphorylation, is required for full MPF activity.

B-type cyclins also undergo phosphorylation in a variety of systems such as sea urchin eggs, yeast, starfish oocytes and human cells (Booher *et al.*, 1989; Meijer *et al.*, 1989; Pines and Hunter, 1989; Pondaven *et al.*, 1990). Previous studies in our laboratory have shown that phosphorylation of cyclin B1 regulates its nuclear localization (Li *et al.*, 1995, 1997). There are five sites of phosphorylation in *Xenopus* cyclin B1 that have been identified: Ser2, Ser94, Ser96, Ser101 and Ser113 (Izumi and Maller, 1991; Li *et al.*, 1995). Four of these sites are located within the cytoplasmic retention signal (CRS) domain, a region consisting of residues 78–127 in *Xenopus* cyclin B1 that was initially believed to contain information necessary for retention of the protein in the cytoplasm (Pines and Hunter, 1994). Mutation of these Ser residues to Ala abrogates the ability of cyclin B1 to translocate to the nucleus and to mature oocytes. Incorporation of a nuclear localization signal (NLS) into the Ala mutant restores biological activity, suggesting that phosphorylation of cyclin B1 within the CRS is important for nuclear translocation (Li *et al.*, 1997). In support of this hypothesis, mutation of these Ser residues to Glu results in localization of cyclin B1 to the nucleus and an acceleration of oocyte maturation (Li *et al.*, 1997). Recently, several studies have shown that the CRS also contains a functional nuclear export signal (NES) that causes cyclin B1 to be exported from the nucleus to the cytoplasm during interphase by interaction with CRM1, and that this NES is inactivated by phosphorylation at one or more sites near the NES (Hagting *et al.*, 1998; Toyoshima *et al.*, 1998; Yang *et al.*, 1998).

While it is clear that cyclin B1 is continuously exported


from the nucleus during interphase, the mechanism of nuclear import of cyclin B1 is unclear and remains an intriguing issue. There are no consensus NLSs in either subunit of MPF, suggesting that an alternative mechanism of nuclear import may exist for MPF. Potentially, the CRS may itself be mediating nuclear localization, especially when in its phosphorylated state, which results in inactivation of the NES. However, previous studies in our laboratory have indicated that the phosphorylated CRS domain does not function as an NLS. Fusion of CRS<sup>WT</sup>, CRS<sup>Ala</sup> or CRS<sup>Glu</sup> onto the N-terminus of pyruvate kinase, which is often used as a reporter for nuclear localization since its size excludes the possibility of passive diffusion, did not result in nuclear localization as discussed in Li *et al.* (1997). In addition, when the CRS of cyclin B1 was appended to cyclin A, the nuclear cyclin A protein was retained in the cytoplasm, although the phosphorylation state of the CRS was not determined (Pines and Hunter, 1994). It has also been suggested that a bipartite NLS may be formed when a cyclin binds to a cdk (Pines and Hunter, 1994). Another possibility may be that cyclins associate with transcription factors or other nuclear proteins as shown in the E2F interaction with cyclin A (Mudryj *et al.*, 1991), or that cyclin B1 may enter the nucleus simply by association with cdc2, which has been speculated to contain a weak histidine-containing putative NLS (Boulikas, 1996). More recently, importin- $\beta$  has been shown to interact directly with a truncated form of cyclin B1, but it was inconclusive how full-length cyclin B1 enters the nucleus (Moore *et al.*, 1999). cdc2-cyclin B1 may also 'piggyback' into the nucleus by binding a third protein containing an NLS, as suggested by Maridor *et al.* (1993).

Using the CRS domain of *Xenopus* cyclin B1, we employed a yeast two-hybrid screen of a mouse embryonic library to identify proteins that may be involved in the nuclear localization of cyclin B1. Interestingly, we isolated mouse cyclin F. Cyclin F is the largest known protein of the cyclin family with a predicted mol. wt of 87 kDa, although it migrates as a 100–110 kDa protein (Bai *et al.*, 1994). Examination of the amino acid sequence of cyclin F indicates the presence of a cyclin box, F-box, PEST sequences and two putative NLS regions. Cyclin F protein accumulates in interphase, reaches maximal levels at G<sub>2</sub>-M phase, and decreases at mitosis just prior to the destruction of cyclin B1 (Bai *et al.*, 1994). Cyclins often complex to a cdk and act as the regulatory binding partner; however, neither cdc2 nor cdk2 has been found to interact with cyclin F (Bai *et al.*, 1994), although cyclin F has been shown to interact with Skp1, a protein involved in proteolysis (Bai *et al.*, 1996). The exact role of cyclin F in cell cycle regulation remains to be clearly defined. In this report, we describe the first known direct interaction between two cyclin proteins, show that the two putative NLS regions of cyclin F are functional, and propose a model by which cyclin F regulates the nuclear localization of cyclin B1.

## Results


### Isolation of cyclin F from a two-hybrid screen

Previous work from our laboratory has shown that nuclear localization of cyclin B1 is regulated by phosphorylation



**Fig. 1.** Schematic representation of *Xenopus* cyclin B1-derived fusion proteins. The single and double CRS derivatives of cyclin B1 (aa 78–127) contained either all Ala or Glu mutations at residues 94, 96, 101 and 113. The NLS-B1 derivatives contained a nuclear localization signal appended to the N-terminus of cyclin B1. The CRS-B1 derivatives contained an additional CRS region appended to full-length cyclin B1.

in the CRS region (Li *et al.*, 1997). Since cyclin B1 lacks a functional nuclear localization signal, another factor may be involved in the translocation of cyclin B1 to the nucleus. To investigate which protein(s) may be binding to the CRS region of cyclin B1 and affecting its localization, a yeast two-hybrid screen was employed. Full-length cyclin B1 could not be used as bait due to its toxicity in yeast. As previous experience suggested that a bait of at least 100 amino acids would be preferable for a successful two-hybrid screen (our unpublished data), another CRS region was appended to extend the length of each bait. With this size constraint in mind, we created two bait constructs of *Xenopus* cyclin B1: CRS<sup>Ala</sup>-CRS<sup>Ala</sup>, which is composed of two tandem CRS domains [amino acids (aa) 78–127] with each of the Ser residues mutated to Ala; and CRS<sup>Glu</sup>-CRS<sup>Glu</sup>, with the Ser residues mutated to Glu (Figure 1). The CRS<sup>Ala</sup>-CRS<sup>Ala</sup> construct prevents any phosphorylation in the CRS domain while the CRS<sup>Glu</sup>-CRS<sup>Glu</sup> construct mimics the phosphorylated state. The wild-type construct was not used in the screen as the phosphorylation state at the four phosphorylation sites within the CRS would be unpredictable. These two bait proteins were screened against a mouse embryonic cDNA library, resulting in 114 double transformants that grew on selective media after 3–4 days at 30°C (see Materials and methods). False positives were eliminated by screening against LexA-lamin, which was used to detect interactions specific to the bait (Vojtek and Hollenberg, 1995). After



**Fig. 2.** Human cyclin B1 and cyclin F constructs. (A) The CRS region of human cyclin B1 was synthesized with the Ser residues at 126, 128, 133 and 147 all mutated to either Ala or Glu. (B) Myristylated human cyclin F (top),  $\Delta$ NLS1/2-cyclin F (middle) and cyclin F (bottom) are shown. The 188 aa region of mouse cyclin F that interacted with the CRS of cyclin B1 corresponding to human cyclin F is indicated at the bottom.


**Table I.** Summary of two-hybrid interactions

LexA fusion	pVP16	Mouse cyclin F (aa 245–432)
CRS <sup>Ala</sup>	–	+
CRS <sup>Glu</sup>	–	+
CRS <sup>Ala</sup> -CRS <sup>Ala</sup>	–	+
CRS <sup>Glu</sup> -CRS <sup>Glu</sup>	–	+
hCRS <sup>Ala</sup>	–	+
hCRS <sup>Glu</sup>	–	+
hCRS <sup>Ala</sup> -CRS <sup>Ala</sup>	–	+
hCRS <sup>Glu</sup> -CRS <sup>Glu</sup>	–	+
Lamin	nd <sup>a</sup>	–

<sup>a</sup>nd, not determined.

further characterization, 10 clones were identified and confirmed as valid interactions.

One of the clones identified was a 188 aa segment (aa 245–432) of mouse cyclin F, which interacted with both CRS<sup>Ala</sup>-CRS<sup>Ala</sup> and CRS<sup>Glu</sup>-CRS<sup>Glu</sup> baits. This segment encompasses the cyclin box and flanking regions of cyclin F (Figure 2B). To examine whether mouse cyclin F could interact with other species of cyclin B1, human CRS and CRS-CRS constructs were also made of each Ala and Glu mutant. Each of these hCRS<sup>Ala</sup>, hCRS<sup>Glu</sup>, hCRS<sup>Ala</sup>-CRS<sup>Ala</sup> and hCRS<sup>Glu</sup>-CRS<sup>Glu</sup> derivatives (Figure 2A) interacted with the fragment of mouse cyclin F isolated from the two-hybrid screen (Table I). In addition, we obtained a full-length human cyclin F clone, which was tested for interaction with each of the *Xenopus* and human CRS and CRS-CRS mutants in the two-hybrid system; however, the protein proved to be toxic to the L40 yeast strain.




**Fig. 3.** Expression of cyclin F RNA during *Xenopus* development. Total RNA isolated from stage VI oocytes or staged embryos was isolated and analyzed by RT-PCR. RNA (200 ng) was assayed for either the presence of cyclin F (upper panel) or, as an internal control, *c-src* RNA (lower panel).

### The potential for cyclin B1 and cyclin F interaction

Both cyclin B1 and cyclin F proteins accumulate at G<sub>2</sub> phase, thus providing an opportunity for interaction. Previously, it had been shown that cyclin F mRNA expression is present in various human tissues (Bai *et al.*, 1994). We also wanted to determine whether cyclin F mRNA expression was present in *Xenopus* early embryonic development. Using quantitative RT-PCR, we examined cyclin F RNA expression in Stage VI oocytes, eggs and early embryos from different developmental stages after fertilization. As shown in Figure 3 (upper panel), the cyclin F RNA transcription levels remain constant to the end of blastulation (stage 9) and begin to decrease at the beginning of gastrulation (stages 10–11). *c-src* RNA expression was used as a positive control for RNA recovery by RT-PCR analysis (Figure 3, lower panel), since the level of *c-src* transcript remains constant throughout oogenesis and early embryogenesis (Rempel *et al.*, 1995). These data indicate that cyclin F mRNA is present during meiotic maturation of oocytes and during early developmental stages and are consistent with the presence of cyclin F protein; however, in the absence of an antiserum against *Xenopus* cyclin F, we have not established whether this mRNA is translated.

### Cyclin B1 and cyclin F interact in vitro and in vivo

To test further for an interaction between cyclin B1 and cyclin F, a glutathione *S*-transferase (GST) fusion protein of human cyclin F was constructed and bacterially expressed. Cyclin B1 constructs, which contain a second CRS domain appended to the N-terminus of full-length *Xenopus* cyclin B1 to mimic the baits used in the two-hybrid screen (Figure 1), were *in vitro* translated. The Ser residues of each CRS domain were mutated to alanine (CRS<sup>Ala</sup>-B1<sup>Ala</sup>), glutamic acid (CRS<sup>Glu</sup>-B1<sup>Glu</sup>) or unaltered (CRS<sup>WT</sup>-B1<sup>WT</sup>). Each *Xenopus* cyclin B1 construct was epitope-tagged to allow specific recognition by the monoclonal antibody (mAb) P5D4 (Kreis and Lodish, 1986). Incubation of GST-cyclin F with *in vitro* translated CRS<sup>Ala</sup>-B1<sup>Ala</sup>, CRS<sup>Glu</sup>-B1<sup>Glu</sup> or CRS<sup>WT</sup>-B1<sup>WT</sup> showed that cyclin F bound to all three derivatives (Figure 4A, lanes 7, 8 and 9). As a negative control, each *in vitro* translated product was incubated with GST protein alone (Figure 4A, lanes 4, 5 and 6). Figure 4B confirms the presence of GST protein (lanes 4, 5 and 6) and GST-cyclin F (lanes 7, 8 and 9).




**Fig. 4.** *In vitro* binding of GST-cyclin F and *Xenopus* cyclin B1 fusion proteins. (A) *Xenopus* cyclin B1 derivatives were translated in reticulocyte lysates (lanes 1, 2 and 3) and then incubated with GST protein alone (lanes 4, 5 and 6) or with bacterially expressed GST-cyclin F (lanes 7, 8 and 9) bound to glutathione-agarose beads. The samples were analyzed by 7.5% SDS-PAGE and immunoblotted with mAb P5D4 directed against the epitope-tag VSV-G on each cyclin B1 derivative. (B) The membrane was re-probed with  $\alpha$ -GST antibody, indicating the presence of GST protein (lanes 4, 5 and 6) and GST-cyclin F (lanes 7, 8 and 9). Molecular mass markers in kilodaltons are indicated.

The interaction between cyclin F and cyclin B1 was also tested *in vivo* by cotransfecting either CRS<sup>Ala</sup>-B1<sup>Ala</sup>, CRS<sup>Glu</sup>-B1<sup>Glu</sup> or CRS<sup>WT</sup>-B1<sup>WT</sup> with cyclin F into COS-1 cells. Cell lysates were subjected to immunoprecipitation with cyclin F antibody, analyzed by SDS-PAGE, and immunoblotted with mAb P5D4 that recognized the epitope-tag of cyclin B1. All three cyclin B1 derivatives coimmunoprecipitated with cyclin F, confirming the interaction between cyclin B1 and cyclin F *in vivo* (data not shown). Similarly, immunoprecipitation with mAb P5D4 followed by immunoblotting with cyclin F antibody also demonstrated association between cyclin B1 and cyclin F (data not shown).

#### ***cdc2*, cyclin B1 and cyclin F form a complex that exhibits histone H1 kinase activity**

Since cyclin B1 is often complexed to *cdc2*, we were interested in determining whether *cdc2* associated with the cyclin B1-cyclin F complex. Cyclin F was transfected into 293T cells, and lysates were then immunoprecipitated with  $\alpha$ -*cdc2* antibody. Immunoblot analysis demonstrated the presence of cyclin F in the  $\alpha$ -*cdc2* immunoprecipitate (Figure 5, upper panel, lane 4). Immunoblotting with cyclin B1 antibody indicated that cyclin B1 was present in each sample (Figure 5, lower panel). These data indicate that *cdc2*, cyclin B1 and overexpressed cyclin F are able to form a complex *in vivo*.

We then wished to determine whether we could detect an interaction between endogenous cyclin B1 and endogenous cyclin F. Using HeLa cells synchronized by double-thymidine block to G<sub>2</sub>-M phase, we immunoprecipitated



**Fig. 5.** *cdc2*, cyclin B1 and cyclin F form a complex. Cyclin F was transfected into 293T cells and then immunoprecipitated with  $\alpha$ -*cdc2* antibody. The samples were analyzed by 10% SDS-PAGE followed by immunoblotting with  $\alpha$ -cyclin F antibody (upper panel) and  $\alpha$ -cyclin B1 (lower panel). The lower band in the immunoprecipitated samples in lanes 3 and 4 of the lower panel represents the IgG band from the  $\alpha$ -*cdc2* antibody. Molecular mass markers are indicated.


with  $\alpha$ -cyclin B1 antibody followed by immunoblotting with  $\alpha$ -cyclin F antibody to show that cyclin F binds to cyclin B1 (Figure 6A, upper panel, lane 2). Immunoblotting with  $\alpha$ -cyclin B1 confirmed the presence of cyclin B1 (Figure 6A, lower panel, lane 2). Addition of the blocking peptide to the  $\alpha$ -cyclin B1 antiserum blocked the recovery of cyclin F in the cyclin B1 immunoprecipitate (Figure 6A, upper panel, lane 3). The lysate alone served as a positive control for the presence of cyclin B1 and cyclin F (Figure 6A, lane 1).

Next we investigated the interaction of endogenous cyclin F with endogenous MPF and whether this complex exhibited histone H1 kinase activity. Synchronized HeLa cells were harvested at G<sub>2</sub>-M phase and immunoprecipitated with  $\alpha$ -*cdc2* antibody. Figure 6B shows the presence of cyclin F (upper panel, lane 2), cyclin B1 (middle panel, lane 2) and *cdc2* (lower panel, lane 2). The lysate alone was a positive control for the presence of all three proteins in Figure 6B, lane 1 of each panel. Thus, *cdc2*, cyclin B1 and cyclin F exist in a complex endogenously. To determine if cyclin F interacts with an active MPF complex, we performed a histone H1 kinase assay. HeLa cells were lysed, immunoprecipitated with  $\alpha$ -cyclin F antibody, and then assayed for histone H1 kinase activity. As shown in Figure 6C, the  $\alpha$ -cyclin F immunoprecipitate displayed kinase activity (lane 1), which was eliminated by inclusion of a blocking peptide to the  $\alpha$ -cyclin F antibody (lane 2). The  $\alpha$ -*cdc2* immunoprecipitate served as a positive control (Figure 6C, lane 3). These results demonstrate that cyclin F can bind to the active MPF complex with retention of kinase activity, although we have not eliminated the possibility that the histone H1 kinase activity may reflect cyclin A-*cdc2* complexes as well as cyclin B1-*cdc2*.

#### ***Cyclin B1 and cyclin F colocalize in immunofluorescence studies***

Cyclin F has been shown to localize predominantly to the nucleus in non-synchronized cells (Bai *et al.*, 1994), whereas cyclin B1 has been shown to re-localize from the






**Fig. 7.** (A) Individual localization of CRS-cyclin B1 and cyclin F in COS-1 cells synchronized in G<sub>2</sub>-M phase. Panels a and c are CRS<sup>Ala</sup>-B1<sup>Ala</sup> and CRS<sup>Glu</sup>-B1<sup>Glu</sup>, respectively, fixed and stained with mAb P5D4 (green). Panel e shows cyclin F fixed and stained with  $\alpha$ -cyclin F antibody (red). Hoechst dye 33342 reveals the nucleus in panels b, d and f (blue). (B) Colocalization of CRS-cyclin B1 and cyclin F. The CRS<sup>Ala</sup>-B1<sup>Ala</sup> and cyclin F cotransfection (top row) shows that the cyclin B1 Ala mutant (green) and cyclin F (red) colocalize (yellow) in the cytoplasm. The CRS<sup>Glu</sup>-B1<sup>Glu</sup> and cyclin F cotransfection (bottom row) indicates that the cyclin B1 Glu mutant (green) and cyclin F (red) also colocalize (yellow), but in the nucleus.

with the following cyclin B1 derivatives: CRS<sup>Ala</sup>-B1<sup>Ala</sup>, NLS-B1<sup>Ala</sup>, CRS<sup>Glu</sup>-B1<sup>Glu</sup> or NLS-B1<sup>Glu</sup>. Transient transfections show that each cyclin B1 derivative predominantly colocalized with  $\Delta$ NLS1/2-cyclin F to the cytoplasm (data not shown). Figure 9B shows the localization of NLS-B1<sup>Glu</sup> (panel c) and  $\Delta$ NLS1/2-cyclin F (panel d). While NLS-B1<sup>Glu</sup> alone is ~100% nuclear (Figure 8A, panel c), upon co-expression with  $\Delta$ NLS1/2-cyclin F, both NLS-B1<sup>Glu</sup> and  $\Delta$ NLS1/2-cyclin F exhibit cytoplasmic localization (Figure 9B, panel e). Thus,  $\Delta$ NLS1/2-cyclin F recruits the nuclear NLS-B1<sup>Glu</sup> to the cytoplasm. Conversely, these results suggest that one role of wild-type cyclin F may be to localize cyclin B1 to the nucleus when the NES of cyclin B1 has been inactivated by phosphorylation.

## Discussion

In this study, we found that a region containing the cyclin box of cyclin F interacts with both CRS<sup>Ala</sup>-CRS<sup>Ala</sup> and CRS<sup>Glu</sup>-CRS<sup>Glu</sup> mutants in a yeast two-hybrid screen. This interaction was confirmed *in vitro* and *in vivo* (data not shown) using full-length cyclin B1 constructs, and established the first known cyclin-cyclin association. The binding between cyclin B1 and cyclin F is strong, as shown by the ability of myr-cyclin F to retain nuclear-targeted cyclin B1 at the plasma membrane. We have also confirmed that cyclin B1 and cyclin F interact endogenously in coimmunoprecipitation experiments.

Both cyclin B1 and cyclin F accumulate at G<sub>2</sub> phase (Bai *et al.*, 1994), providing the opportunity for a potential




**Fig. 8.** (A) Individual localization of NLS-B1 and myr-cyclin F in COS-1 cells synchronized in G<sub>2</sub>-M phase. Panels a and c are NLS-B1 derivatives stained with mAb P5D4 (green). Panel e is myristylated cyclin F stained with  $\alpha$ -cyclin F antibody (red). Hoechst dye shows the nucleus in panels b, d and f (blue). (B) Colocalization of NLS-B1 and myr-cyclin F. The NLS-B1<sup>Ala</sup> and myr-cyclin F cotransfection (top row) shows that the NLS-B1<sup>Ala</sup> mutant (green) and myr-cyclin F (red) colocalize (yellow). The NLS-B1<sup>Glu</sup> and myr-cyclin F cotransfection indicates that the NLS-B1<sup>Glu</sup> mutant (green) and myr-cyclin F (red) colocalize (yellow).

interaction between the two cyclins. In addition, our immunofluorescence data consistently showed that ~35% of cells expressing cyclin F protein exhibited perinuclear/cytoplasmic localization, and ~65% of cells exhibited nuclear localization (our unpublished data), further providing the opportunity for cyclin F to bind to cyclin B1 in the cytoplasm. We have also shown that cyclin F mRNA is expressed in *Xenopus* oocytes and throughout early embryonic development, indicating that cyclin F is potentially present to interact with cyclin B1.

We have also found that cdc2 can be detected in the cyclin B1-cyclin F complex *in vivo*. Since it has been reported that cdc2 and cyclin F do not interact *in vitro* (Bai *et al.*, 1994), it is likely that cyclin B1 is serving as the link between cyclin F and cdc2. The binding of

cyclin F to cyclin B1 may allow cyclin F to interact indirectly with cdc2 and potentially help regulate its kinase activity in a surrogate manner. This is supported by the histone H1 kinase activity associated with the cyclin F immunoprecipitate, demonstrating that MPF maintains its kinase activity when complexed to cyclin F.

Immunofluorescence data show that cyclin B1 colocalizes with cyclin F, suggesting that cyclin F can bind indiscriminately to either CRS<sup>Ala</sup>-B1<sup>Ala</sup> or CRS<sup>Glu</sup>-B1<sup>Glu</sup>, the unphosphorylated or phosphorylated forms, respectively, of cyclin B1. In Figure 10, a model is proposed to explain why both the Ala and Glu derivatives of cyclin B1 colocalize with cyclin F. Figure 10A shows that cyclin F transports CRS<sup>Ala</sup>-B1<sup>Ala</sup> into the nucleus via the two NLS motifs of cyclin F, both of which have been shown in this



**Fig. 9.**  $\Delta$ NLS1/2-cyclin F localizes to the cytoplasm and retains NLS-B1<sup>Glu</sup> in the cytoplasm. (A)  $\Delta$ NLS1/2-cyclin F transfected cells were fixed and stained with an antibody directed against the epitope-tag of cyclin F (panel a, red). The nucleus is seen with Hoechst dye (panel b, blue) and indicates that  $\Delta$ NLS1/2-cyclin F is completely excluded from the nucleus. (B) NLS-B1<sup>Glu</sup> (panel c, green) is cotransfected with  $\Delta$ NLS1/2-cyclin F (panel d, red) and they colocalize (panel e, yellow).

study to be functional. Once in the nucleus, the functional NES of CRS<sup>Ala</sup>-B1<sup>Ala</sup> shuttles the complex back to the cytoplasm. In these immunofluorescence studies, the CRS<sup>Ala</sup>-B1<sup>Ala</sup>-cyclin F complex was not seen to colocalize to the nucleus at any time. A possible explanation is that the rate of nuclear export is faster than the rate of nuclear import. Thus, CRS<sup>Ala</sup>-B1<sup>Ala</sup> and cyclin F are seen to colocalize to the cytoplasm.


The model presented in Figure 10B explains the colocalization seen between CRS<sup>Glu</sup>-B1<sup>Glu</sup> and cyclin F. In this case, the CRS<sup>Glu</sup>-B1<sup>Glu</sup>-cyclin F complex traverses into the nucleus utilizing the two NLS regions of cyclin F. Recent studies have shown that phosphorylation of Ser residues within the CRS domain reduced the binding affinity for CRM1 and resulted in a decrease of nuclear export (Yang *et al.*, 1998). Thus, the NES of CRS<sup>Glu</sup>-B1<sup>Glu</sup> is inactivated due to the mimicked phosphorylation by the Glu residues, and cyclin B1 is retained in the nucleus with cyclin F. The phosphorylated form of cyclin B1, which may also be complexed to cdc2, would then be properly localized in the nucleus to trigger the onset of mitosis by phosphorylating nuclear components necessary for mitosis. Therefore, the inactivation of the NES by phosphorylation of the Ser residues within cyclin B1 may be the trigger that initiates mitosis.

With regard to wild-type cyclin B1, the upper panel of Figure 10C depicts the localization of unphosphorylated cyclin B1 into the nucleus via the two NLS motifs of its binding partner cyclin F. In the nucleus, the functional NES of unphosphorylated cyclin B1 would result in export of the complex to the cytoplasm. With respect to the phosphorylated form of cyclin B1 complexed to cyclin F, we propose that the complex enters the nucleus via the NLS regions of cyclin F as shown in the lower panel of Figure 10C. Once in the nucleus, MPF is properly localized to initiate mitosis.

The ability of cyclin F to regulate the localization of cyclin B1 is clearly shown by the myr-cyclin F and NLS-B1 immunofluorescence studies. In addition, deletion

of the two NLS regions of cyclin F resulted in its localization to the cytoplasm rather than to the nucleus. When this  $\Delta$ NLS1/2-cyclin F was cotransfected with various cyclin B1 derivatives, the resulting complexes were colocalized to the cytoplasm. The ability of  $\Delta$ NLS1/2-cyclin F to retain the NLS-B1<sup>Glu</sup> derivative in the cytoplasm is especially notable, suggesting the possibility that cyclin F itself may contain an NES whose activity is manifested in the absence of its NLS activity. Alternatively, cyclin F may contain some type of CRS whose function becomes apparent upon inactivation of its NLS function. The presence of an NES in cyclin F may also be consistent with our data in the myr-cyclin F/NLS-B1 colocalization to the cytoplasm; the myristylation signal appended to cyclin F in conjunction with an NES overcomes the NLS appended to cyclin B1.

Our data support a model in which cyclin F mediates the nuclear localization of cyclin B1. Most nuclear import/export factors interact directly with their cargo, but importin- $\beta$  uses the adaptor protein importin- $\alpha$  to interact with an NLS-containing protein (Gorlich *et al.*, 1995a,b). Importin- $\alpha$  binds to importin- $\beta$  to form a heterodimer, and during translocation through the nuclear pore complex (NPC), the heterodimer dissociates: importin- $\alpha$  and the NLS-containing protein enter and accumulate in the nucleoplasm as importin- $\beta$  accumulates at the NPC (Gorlich *et al.*, 1995b; Moroianu *et al.*, 1995). It has recently been shown that importin- $\beta$  can bind directly to a truncated form of *Xenopus* cyclin B1 (aa 121–397) and be imported into the nucleus without the presence of importin- $\alpha$  (Moore *et al.*, 1999). However, this truncated cyclin B1 is lacking the CRS/NES region that has been shown to regulate cyclin B1 localization, and the import was not seen with full-length cyclin B1. Thus, the interaction found between cyclin F and the CRS region of cyclin B1 (aa 78–127) supports the possibility that importin- $\alpha$  binds to the NLS region(s) of cyclin F, which can then interact with importin- $\beta$  to translocate cyclin B1 as well as cyclin F into the nucleus. Alternatively, the



**Fig. 10.** Model of cyclin B1–cyclin F colocalization. (A) CRS<sup>Ala</sup>-B1<sup>Ala</sup> and cyclin F colocalize in the cytoplasm, undergo nuclear transport, and then re-localize to the cytoplasm due to the functional NES of CRS<sup>Ala</sup>-B1<sup>Ala</sup>. (B) CRS<sup>Glu</sup>-B1<sup>Glu</sup> and cyclin F interact in the cytoplasm and then colocalize to the nucleus. The complex is then retained in the nucleus due to the inactive NES of CRS<sup>Glu</sup>-B1<sup>Glu</sup> and properly localized to trigger the onset of mitosis. (C) Upper panel: unphosphorylated cyclin B1 and cyclin F interact in the cytoplasm and enter the nucleus. Re-localization to the cytoplasm may then occur via the active NES of cyclin B1. Lower panel: phosphorylated cyclin B1 and cyclin F colocalize from the cytoplasm to the nucleus. MPF is active and ready to initiate mitosis. In each panel, cdc2 is complexed to cyclin B1, although it is not represented.

cyclin B1–cyclin F interaction could provide the link between both importin- $\beta$  and importin- $\alpha$ . Whichever the case may be, the interaction of cyclin F would now provide a boost in cyclin B1 translocation to the nucleus since the truncated cyclin B1 protein exhibited slow nuclear import mediated by importin- $\beta$  (Moore *et al.*, 1999). Thus, at the G<sub>2</sub>–M transition of the cell cycle, the role of cyclin F may be to increase nuclear import of cyclin B1, possibly through an interaction with importin- $\alpha$  in addition to importin- $\beta$ .

One of the few known roles of cyclin F is that it interacts with Skp1 through its F-box motif (Bai *et al.*, 1996). Skp1 is involved in G<sub>1</sub>–S and G<sub>2</sub>–M transitions of the cell cycle, and is linked to ubiquitin-mediated proteolysis of many proteins such as Sic1p, Cln2p and Clb5p (Bai *et al.*, 1996). Skp1 has been shown to act as a ubiquitin ligase when associated with Cdc53 and the F-box-containing protein Cdc4, forming a complex named SCF<sup>Cdc4</sup> (Skp1, Cdc53 and F-box protein Cdc4) (Skowrya *et al.*, 1997). Skowrya *et al.* (1997) proposed a model in which an F-box protein specifically recruits phosphoryl-

ated substrates to the SCF complex and forms a ubiquitylation complex. Thus, cyclin F may be the F-box protein that completes the ubiquitin ligase complex, forming an SCF<sup>cyclinF</sup> and targeting phosphorylated cyclin B1 for ubiquitin-mediated degradation. However, this may not be the case as the timing of cyclin degradation suggests that cyclin F is degraded prior to cyclin B1 (Bai *et al.*, 1994). Another role may be that cyclin F itself is destroyed by interaction with Skp1 and/or another F-box-containing protein that recognizes cyclin F. This implies that cyclin F degradation may be necessary for the degradation of cyclin B1. The destruction of cyclin B1 by the ubiquitylation pathway results in the inactivation of MPF, and the cell exits mitosis (Glotzer *et al.*, 1991). Further experiments will be necessary to determine if cyclin F is indeed involved in the proteolysis of cyclin B1.

In conclusion, we have shown a novel interaction between two cyclins. The CRS region of cyclin B1 binds a region including the cyclin box of cyclin F and, through this interaction, cyclin F is able to colocalize with cyclin B1 and mediate nuclear localization of the phosphorylated form of cyclin B1. Since cyclin F has two putative NLSs and is present at approximately the same time as cyclin B1 during the cell cycle, the translocation of cyclin B1 into the nucleus is likely to be regulated by cyclin F. In this manner, one role of cyclin F in the cell cycle may be to regulate the nuclear translocation of cyclin B1 at the G<sub>2</sub>–M transition.

## Materials and methods

### Two-hybrid constructs

The two-hybrid plasmids pBTM116, pVP16 and pLexA-lamin (Vojtek *et al.*, 1993) were generous gifts from S.Hollenberg and J.A.Cooper (Fred Hutchinson Cancer Research Center). Construction of CRS<sup>Ala</sup>-CRS<sup>Ala</sup> and CRS<sup>Glu</sup>-CRS<sup>Glu</sup> was achieved through digestion of the *Bsu*36I and *Bam*HI sites from full-length cyclin B1 constructs made previously in our laboratory (Li *et al.*, 1997). A linker oligonucleotide containing a stop codon was added after the last CRS domain of each construct. Each construct was then cut with *Eco*RI and *Bam*HI and inserted into the pBTM116 vector, which was constructed by P.Bartel and S.Fields, at the corresponding sites. Human CRS regions were synthesized and inserted into the pBTM116 vector, the sequence was verified through dideoxy nucleotide sequencing. Human cyclin F (Bai *et al.*, 1994), a generous gift from S.J.Elledge, was cloned into the pVP16 plasmid for two-hybrid analysis at the *Bam*HI and *Not*I sites using an oligonucleotide linker.

### Two-hybrid screen

The two-hybrid screen was performed according to previously published protocols (Vojtek *et al.*, 1993; Vojtek and Hollenberg, 1995). Each LexA-CRS<sup>Ala</sup>-CRS<sup>Ala</sup> and LexA-CRS<sup>Glu</sup>-CRS<sup>Glu</sup> fusion construct was cotransformed with a 9.5 d.p.c. mouse embryo cDNA library fused to pVP16 into the L40 strain of *Saccharomyces cerevisiae*. The transformants were selected on –His plates for 3–4 days at 30°C with the addition of 5 mM 3-aminotriazole (3-AT) (Sigma) for the CRS<sup>Ala</sup>-CRS<sup>Ala</sup> transformants and 50 mM 3-AT for the CRS<sup>Glu</sup>-CRS<sup>Glu</sup> transformants. An interaction was scored positive for the ability of the yeast to grow beyond bait plus empty pVP16 vector and to overcome 3-AT inhibition. False positives were discriminated by testing cDNA-pVP16 against LexA-lamin. The remaining cDNA-pVP16 clones were then shuttled into *Escherichia coli* HB101 cells, rescued, and then sequenced as previously described (Vojtek and Hollenberg, 1995). The sequences were then entered into the NCBI BLAST program for identification.

### RT-PCR assays

Total RNA from oocytes and staged embryos was prepared using TRIzol reagent as described by the manufacturer (Gibco-BRL, Life Technologies). RT-PCR was performed using 200 ng of total RNA,

50 pmol of sense primer (ATATGCACACGCTTTATTAGC) and 50 pmol of antisense primer (ATGGACGTGCTTGTCTGTGC) for the cyclin F gene to generate a 325 bp fragment, and 50 pmol of sense primer (TACATCAGTCTCGAACTC) and 50 pmol of antisense primer (ACAGCATATACTGTACCAG) for the *c-src* gene to generate a 347 bp fragment.

#### **In vitro binding assay**

The cyclin F-pVP16 construct was cleaved by *Bam*HI and *Nor*I and inserted into pGEX-3X to generate GST-cyclin F. The GST-cyclin F fusion protein was produced by *E.coli* BL21 cells. The cells were induced with 1 mM isopropyl- $\beta$ -D-thiogalactopyranoside (IPTG) for 3 h. The bacterial pellet was resuspended in 50 ml of lysis buffer [50 mM Tris pH 7.5, 150 mM NaCl, 10% glycerol, 1 mM dithiothreitol (DTT), 1 mM EDTA, 1 mM phenylmethylsulfonyl fluoride (PMSF), 10  $\mu$ g/ml aprotinin, 10  $\mu$ g/ml pepstatin and 10  $\mu$ g/ml leupeptin] and sonicated five times for 15 s each. The pellet was resuspended in lysis buffer and 6 M urea, then slowly dialyzed. GST-cyclin F was then incubated with glutathione-agarose beads (Sigma) at 4°C and washed three times. Coupled *in vitro* transcription-translation (Promega) was performed according to the manufacturer's instructions for the full-length CRS<sup>Ala</sup>-B1<sup>Ala</sup>, CRS<sup>Glu</sup>-B1<sup>Glu</sup> and CRS<sup>WT</sup>-B1<sup>WT</sup> derivatives. Each contained an epitope-tag derived from vesicular stomatitis virus glycoprotein (VSV-G) appended at the C-terminus, which was recognized by murine mAb P5D4 (Kreis and Lodish, 1986) as previously described (Li *et al.*, 1997). The *in vitro* translated products were individually incubated with GST-cyclin F or GST proteins overnight at 4°C in binding buffer (20 mM Tris pH 7.5, 25 mM NaCl, 10% glycerol, 1 mM DTT, 1 mM EDTA, 1 mM PMSF, 10  $\mu$ g/ml aprotinin, 10  $\mu$ g/ml pepstatin and 10  $\mu$ g/ml leupeptin). The reactions were then washed four times in binding buffer, boiled for 3 min in sample buffer, and analyzed on 7.5% SDS-PAGE followed by immunoblotting with mAb P5D4 (1:2000). Enhanced chemiluminescence (ECL) (Amersham) was then used to detect proteins. The membrane was then reprobed with GST antibody (Santa Cruz Biotechnology, Inc.) at 1:4000 and detected by ECL.

#### **Coimmunoprecipitation**

The cyclin F gene was cloned into pcDNA3 from the pAB23BXN vector. 293T cells were cultured in Dulbecco's modified Eagle's medium (DMEM), supplemented with 10% fetal bovine serum (FBS) and incubated at 37°C in 10% CO<sub>2</sub>. Sub-confluent cells were transfected with 20  $\mu$ g of pcDNA3 or cyclin F-pcDNA3 by calcium phosphate precipitation (Chen and Okayama, 1987) in a 10 cm dish. Twenty-four hours after transfection, the cells were harvested and lysed in lysis buffer (20 mM Tris pH 8.0, 150 mM NaCl, 0.5% NP-40, 1 mM Na<sub>3</sub>VO<sub>4</sub>, 1 mM NaF, 1 mM benzamide, 0.1 mM *p*-nitrophenylphosphate, 1 mM DTT, 1 mM PMSF, 10  $\mu$ g/ml aprotinin, 10  $\mu$ g/ml pepstatin and 10  $\mu$ g/ml leupeptin). Lysates were pre-cleared with 40  $\mu$ l of protein A-Sepharose beads and incubated with 2  $\mu$ g of p34<sup>cdc2</sup> antibody (Santa Cruz Biotechnology, Inc.) overnight at 4°C. Protein A-Sepharose beads were then added and the immunoprecipitated samples were washed four times, boiled for 3 min in sample buffer, and analyzed on 10% SDS-PAGE. The membrane was immunoblotted with  $\alpha$ -cyclin F antibody (Santa Cruz Biotechnology, Inc.) at 1:8000 and reprobed with  $\alpha$ -cyclin B1 antibody (Santa Cruz Biotechnology, Inc.) at 1:2000, followed by ECL (Amersham).

HeLa cells were cultured in 10% FBS in 5% CO<sub>2</sub> to ~60% confluency and then synchronized in G<sub>2</sub>-M phase by double-thymidine block as described elsewhere (Bai *et al.*, 1994). Cells were harvested 9 h after release with 0.1% NP-40 lysis buffer (0.1% NP-40, 20 mM Tris-HCl pH 7.5, 100 mM NaCl, 1 mM DTT, 1 mM Na<sub>3</sub>VO<sub>4</sub>, 10  $\mu$ g/ml aprotinin, 10  $\mu$ g/ml leupeptin, 1 mM PMSF, 200  $\mu$ M *N*-acetyl-leucinyll-leucinyll-norleucine and 5 mM *N*-ethylmaleimide). Lysates were pre-cleared with protein A-Sepharose beads and then immunoprecipitated with 2  $\mu$ g of  $\alpha$ -cdc2,  $\alpha$ -cyclin B1 or  $\alpha$ -cyclin B1 antibody plus 10  $\mu$ g of corresponding blocking peptide (Santa Cruz Biotechnology, Inc.). After washing four times, the samples were boiled for 3 min in sample buffer and analyzed by 10% SDS-PAGE. The gel was transferred to nitrocellulose membrane and then immunoblotted for cdc2, cyclin B1 or cyclin F, followed by ECL (Amersham).

#### **Histone H1 kinase assay**

HeLa cells cultured in 10% FBS were lysed in 0.1% NP-40 lysis buffer and then pre-cleared with protein A-Sepharose followed by immunoprecipitation with 2  $\mu$ g of  $\alpha$ -cdc2,  $\alpha$ -cyclin F or  $\alpha$ -cyclin B1 plus 10  $\mu$ g of corresponding blocking peptide (Santa Cruz Biotechnology, Inc.). The immunoprecipitates were washed four times prior to the

addition of 50  $\mu$ l of kinase buffer (50 mM Tris-HCl pH 7.5, 10 mM MgCl<sub>2</sub>, 1 mM DTT, 20 mM EGTA, 50 mM ATP, 10 mCi of [ $\gamma$ -<sup>32</sup>P]ATP) and 74  $\mu$ g/ml H1 histone (Boehringer Mannheim) for 10 min at 30°C. Sample buffer was added to stop the reaction and 50  $\mu$ l of each sample were analyzed by 10% SDS-PAGE, followed by autoradiography.

#### **Immunofluorescence microscopy**

COS-1 cells were plated onto glass coverslips and cultured in DMEM supplemented with 10% FBS at 37°C in 5% CO<sub>2</sub>. Cells were transiently transfected by calcium phosphate precipitation with 10  $\mu$ g of each DNA in a 60 mm dish. An epitope-tag consisting of 12 amino acids (EPDGAALWVHLL) was synthesized and appended to the C-terminus of cyclin F. The myristylated cyclin F construct was constructed by appending a 13 aa myristylation sequence from p60<sup>src</sup> (Cross *et al.*, 1984; Pellman *et al.*, 1985). The  $\Delta$ NLS1/2-cyclin F construct was derived from the epitope-tagged cyclin F by deletion of residues 35–57 ( $\Delta$ NLS1) and residues 568–574 ( $\Delta$ NLS2). Each cyclin B1 derivative contained a VSV-G epitope-tag at the C-terminus. Sixteen hours after transfection, the cells were re-fed and then synchronized by double-thymidine block. Cells were treated with 2 mM thymidine for 15 h to arrest cells in S phase, rinsed three times, and released with fresh 10% FBS medium for 9 h. The cells were then retreated with 2 mM thymidine for 15 h. Five and a half hours after release >50% of the cells had a rounded mitotic phenotype. Cells were then fixed at this G<sub>2</sub>-M phase with 3% paraformaldehyde in phosphate-buffered saline (PBS) for 15 min and permeabilized with 0.1% Triton X-100, 0.2 M glycine and 2.5% FBS in PBS for 45 min. Primary antibody P5D4 (1:1000) was used to detect VSV-G-tagged cyclin B1 derivatives with FITC-conjugated goat anti-mouse secondary antibody (Boehringer Mannheim). To detect epitope-tagged cyclin F, affinity-purified rabbit antisera were made (Lenormand *et al.*, 1999) and visualized with rhodamine-conjugated goat anti-rabbit secondary antibody (Boehringer Mannheim). Coverslips were mounted on slides with 100 mM Tris pH 8.0, 90% glycerol, 1 mg/ml phenylene-diamine and 1  $\mu$ g/ml Hoechst dye 33342 to detect DNA.

#### **Acknowledgements**

We thank S.Hollenberg and J.A.Cooper for the kind gifts of pBTM116, pVP16 and pLexA-lamin plasmids; P.Bartel and S.Fields for construction of the pBTM116 vector; V.Malhotra for providing mAb P5D4; and S.J.Elledge for kindly providing human cyclin F. We also thank K.C.Hart, R.W.Dellinger, A.N.Meyer, J.A.Tynan and other members of the D.J.D. laboratory for critical review of the manuscript. Support from NIH/NCI CA34456 is gratefully acknowledged. M.K. also gratefully acknowledges support from NIH/GM07313, and E.A.B. gratefully acknowledges support from NIH/NCI T32-CA09523.

#### **References**

- Bai,C., Richman,R. and Elledge,S.J. (1994) Human cyclin F. *EMBO J.*, **13**, 6087–6098.
- Bai,C., Sen,P., Hofmann,K., Ma,L., Goebel,M., Harper,J.W. and Elledge,S.J. (1996) SKP1 connects cell cycle regulators to the ubiquitin proteolysis machinery through a novel motif, the F-box. *Cell*, **86**, 263–274.
- Booher,R.N., Alfa,C.E., Hyams,J.S. and Beach,D.H. (1989) The fission yeast *cdc2/cdc13/suc1* protein kinase: regulation of catalytic activity and nuclear localization. *Cell*, **58**, 485–497.
- Boulikas,T. (1996) Nuclear import of protein kinases and cyclins. *J. Cell Biol.*, **60**, 61–82.
- Chen,C. and Okayama,H. (1987) High-efficiency transformation of mammalian cells by plasmid DNA. *Mol. Cell. Biol.*, **7**, 2745–2752.
- Clute,P. and Pines,J. (1999) Temporal and spatial control of cyclin B1 destruction in metaphase. *Nature Cell Biol.*, **1**, 82–87.
- Cross,F.R., Garber,E.A., Pellman,D. and Hanafusa,H. (1984) A short sequence in the p60<sup>src</sup> N terminus is required for p60<sup>src</sup> myristylation and membrane association and for cell transformation. *Mol. Cell. Biol.*, **4**, 1834–1842.
- Draetta,G. and Beach,D. (1988) Activation of cdc2 protein kinase during mitosis in human cells: cell cycle-dependent phosphorylation and subunit rearrangement. *Cell*, **54**, 17–26.
- Draetta,G., Luca,F., Westendorf,J., Bruzuela,L., Ruderman,J. and Beach,D. (1989) cdc2 protein kinase is complexed with both cyclin A and B: evidence for proteolytic inactivation of MPF. *Cell*, **56**, 829–838.

- Evans,T., Rosenthal,E.T., Youngblom,J., Distel,D. and Hunt,T. (1983) Cyclin: a protein specified by maternal mRNA in sea urchin eggs that is destroyed at each cleavage division. *Cell*, **33**, 389–396.
- Fesquet,D. et al. (1993) The *MO15* gene encodes the catalytic subunit of a protein kinase that activates cdc2 and other cyclin-dependent kinases (CDKs) through phosphorylation of Thr 161 and its homologues. *EMBO J.*, **12**, 3111–3121.
- Gautier,J., Minshull,J., Lohka,M., Glotzer,M., Hunt,T. and Maller,J.L. (1990) Cyclin is a component of maturation promoting factor from *Xenopus*. *Cell*, **60**, 487–494.
- Gautier,J., Solomon,M.J., Booher,R.N., Bazan,J.F. and Kirschner,M.W. (1991) cdc25 is a specific tyrosine phosphatase that directly activates p34<sup>cdc2</sup>. *Cell*, **67**, 197–211.
- Glotzer,M., Murray,A.W. and Kirschner,M.W. (1991) Cyclin is degraded by the ubiquitin pathway. *Nature*, **349**, 132–138.
- Gorlich,D., Kostka,S., Kraft,R., Dingwall,C., Laskey,R.A., Hartmann,E. and Prehn,S. (1995a) Two different subunits of importin cooperate to recognize nuclear localization signals and bind them to the nuclear envelope. *Curr. Biol.*, **5**, 383–392.
- Gorlich,D., Vogel,F., Mills,A.D., Hartmann,E. and Laskey,R.A. (1995b) Distinct functions for the two importin subunits in nuclear protein import. *Nature*, **377**, 246–248.
- Gould,K.L. and Nurse,P. (1989) Tyrosine phosphorylation of the fission yeast cdc2+ protein kinase regulates entry into mitosis. *Nature*, **342**, 39–45.
- Gould,K.L., Moreno,S., Owen,D.J., Sazer,S. and Nurse,P. (1991) Phosphorylation at Thr161 is required for *Schizosaccharomyces pombe* p34<sup>cdc2</sup> function. *EMBO J.*, **10**, 3297–3309.
- Hagting,A., Karlsson,C., Clute,P., Jackman,M. and Pines,J. (1998) MPF localization is controlled by nuclear export. *EMBO J.*, **17**, 4127–4138.
- Honda,R., Ohba,Y., Nagata,A., Okayama,H. and Yasuda,H. (1993) Dephosphorylation of human p34<sup>cdc2</sup> kinase on both Thr-14 and Tyr-15 by human cdc25B phosphatase. *FEBS Lett.*, **318**, 331–334.
- Hunt,T. (1991) Cyclins and their partners: from a simple idea to complex reality. *Semin. Cell Biol.*, **2**, 213–222.
- Izumi,T. and Maller,J.L. (1991) Phosphorylation of *Xenopus* cyclins B1 and B2 is not required for cell cycle transitions. *Mol. Cell. Biol.*, **11**, 3860–3867.
- Kobayashi,H., Stewart,E., Poon,R., Adamczewski,J.P., Gannon,J. and Hunt,T. (1992) Identification of the domains in cyclin A required for binding to and activation of, p34<sup>cdc2</sup> and p32<sup>cdk2</sup> protein kinase subunits. *Mol. Biol. Cell*, **3**, 1279–1294.
- Kornbluth,S., Sebastian,B., Hunter,T. and Newport,J. (1994) Membrane localization of the kinase which phosphorylates p34<sup>cdc2</sup> on threonine 14. *Mol. Biol. Cell*, **5**, 273–282.
- Kreis,T.E. and Lodish,H.F. (1986) Oligomerization is essential for transport of vesicular stomatitis viral glycoprotein to the cell surface. *Cell*, **46**, 929–937.
- Krek,W. and Nigg,E.A. (1991) Differential phosphorylation of vertebrate p34<sup>cdc2</sup> kinase at the G<sub>1</sub>/S and G<sub>2</sub>/M transitions of the cell cycle: identification of major phosphorylation sites. *EMBO J.*, **10**, 305–316.
- Labbe,J.-C., Capony,J.-P., Caput,D., Cavadore,J.-C., Derancourt,J., Kaghdad,M., Lelias,J.-M., Picard,A. and Doree,M. (1989) MPF from starfish oocytes at first meiotic metaphase is a heterodimer containing one molecule of cdc2 and one molecule of cyclin B. *EMBO J.*, **8**, 3053–3058.
- Lenormand,J.-L., Dellinger,R.W., Knudsen,K., Subramani,S. and Donoghue,D.J. (1999) Speedy: a novel cell cycle regulator of the G<sub>2</sub>/M transition. *EMBO J.*, **18**, 1869–1877.
- Li,J., Meyer,A.N. and Donoghue,D.J. (1995) Requirement for phosphorylation of cyclin B1 for *Xenopus* oocyte maturation. *Mol. Biol. Cell*, **6**, 1111–1124.
- Li,J., Meyer,A.N. and Donoghue,D.J. (1997) Nuclear localization of cyclin B1 mediates its biological activity and is regulated by phosphorylation. *Proc. Natl Acad. Sci. USA*, **94**, 502–507.
- Maridor,G., Gallant,P., Golsteyn,R. and Nigg,E.A. (1993) Nuclear localization of vertebrate cyclin A correlates with its ability to form complexes with cdk catalytic subunits. *J. Cell Sci.*, **106**, 535–544.
- Meijer,L., Arion,D., Golsteyn,R., Pines,J., Brizuela,L., Hunt,T. and Beach,D. (1989) Cyclin is a component of the sea urchin egg M-phase specific histone H1. *EMBO J.*, **8**, 2275–2282.
- Millar,J.B., McGowan,C.H., Lenaers,G., Jones,R. and Russell,P. (1991) p80<sup>cdc25</sup> mitotic inducer is the tyrosine phosphatase that activated p34<sup>cdc2</sup> kinase in fission yeast. *EMBO J.*, **10**, 4301–4309.
- Minshull,J., Pines,J., Golsteyn,R., Standart,N., Mackie,S., Colman,A., Blow,J., Ruderman,J.V., Wu,M. and Hunt,T. (1989) The role of cyclin synthesis, modification and destruction in the control of cell division. *J. Cell Sci., Suppl.*, **12**, 77–97.
- Moore,J.D., Yang,J., Truant,R. and Kornbluth,S. (1999) Nuclear import of cdk/cyclin complexes: identification of distinct mechanisms for import of cdk2/cyclin E and cdc2/cyclin B1. *J. Cell Biol.*, **144**, 213–224.
- Moroianu,J., Hijikata,M., Blobel,G. and Radu,A. (1995) Mammalian karyopherin  $\alpha 1\beta$  and  $\alpha 2\beta$  heterodimers:  $\alpha 1$  or  $\alpha 2$  subunit binds nuclear localization signal and  $\beta$  subunit interacts with peptide repeat-containing nucleoporins. *Proc. Natl Acad. Sci. USA*, **92**, 6532–6536.
- Mudryj,M., Devoto,S.H., Hiebert,S.W., Hunter,T., Pines,J. and Nevins,J.R. (1991) Cell cycle regulation of the E2F transcription factor involves an interaction with cyclin A. *Cell*, **65**, 1243–1253.
- Mueller,P.R., Coleman,T.R., Kumagai,A. and Dunphy,W.G. (1995) Myt1: a membrane-associated inhibitory kinase that phosphorylates cdc2 on both threonine-14 and tyrosine-15. *Science*, **270**, 86–90.
- Nugent,J.H., Alfa,C.E., Young,T. and Hyams,J.S. (1991) Conserved structural motifs in cyclins identified by sequence analysis. *J. Cell Sci.*, **99**, 669–674.
- Nurse,P. (1990) Universal control mechanism regulating onset of M-phase. *Nature*, **344**, 503–508.
- Parker,L.L., Atherton-Fessler,S. and Pivnicka-Worms,H. (1992) p107<sup>wee1</sup> is a dual-specificity kinase that phosphorylates p34<sup>cdc2</sup> on tyrosine 15. *Proc. Natl Acad. Sci. USA*, **89**, 2917–2921.
- Pellman,D., Gerber,E.A., Cross,F.R. and Hanafusa,H. (1985) An N-terminal peptide from p60<sup>src</sup> can direct myristylation and plasma membrane localization when fused to heterologous proteins. *Nature*, **314**, 374–377.
- Pines,J. and Hunter,T. (1989) Isolation of a human cyclin cDNA: evidence for cyclin mRNA and protein regulation in the cell cycle and for interaction with p34<sup>cdc2</sup>. *Cell*, **58**, 833–846.
- Pines,J. and Hunter,T. (1991) Human cyclins A and B1 are differentially located in the cell and undergo cell cycle-dependent nuclear transport. *J. Cell Biol.*, **115**, 1–17.
- Pines,J. and Hunter,T. (1994) The differential localization of human cyclins A and B is due to a cytoplasmic retention signal in cyclin B. *EMBO J.*, **13**, 3772–3781.
- Pondaven,P., Meijer,L. and Beach,D. (1990) Activation of M-phase-specific histone H1 kinase by modification of the phosphorylation of its p34<sup>cdc2</sup> and cyclin components. *Genes Dev.*, **4**, 9–17.
- Poon,R.Y., Yamashita,K., Adamczewski,J.P., Hunt,T. and Shuttleworth,J. (1993) The cdc2-related protein p40<sup>MO15</sup> is the catalytic subunit of a protein kinase that can activate p32<sup>cdk2</sup> and p34<sup>cdc2</sup>. *EMBO J.*, **12**, 3123–3132.
- Rempel,R.E., Sleight,S.B. and Maller,J.L. (1995) Maternal *Xenopus* Cdk2–cyclin E complexes function during meiotic and early embryonic cell cycles that lack a G<sub>1</sub> phase. *J. Biol. Chem.*, **270**, 6843–6855.
- Russell,P. and Nurse,P. (1987) Negative regulation of mitosis by *wee1+*, a gene encoding a protein kinase homolog. *Cell*, **49**, 559–567.
- Skowyra,D., Craig,K.L., Tyers,M., Elledge,S.J. and Harper,J.W. (1997) F-box proteins are receptors that recruit phosphorylated substrates to the SCF ubiquitin-ligase complex. *Cell*, **91**, 209–219.
- Solomon,M.J., Glotzer,M., Lee,T.H., Philippe,M. and Kirschner,M.W. (1990) Cyclin activation of p34<sup>cdc2</sup>. *Cell*, **63**, 1013–1024.
- Solomon,M.J., Lee,T. and Kirschner,M.W. (1992) Role of phosphorylation in p34<sup>cdc2</sup> activation: identification of an activating kinase. *Mol. Biol. Cell*, **3**, 13–27.
- Toyoshima,F., Moriguchi,F., Wada,A., Fukuda,M. and Nishida,E. (1998) Nuclear export of cyclin B1 and its possible role in the DNA damage-induced G<sub>2</sub> checkpoint. *EMBO J.*, **17**, 2728–2735.
- Vojtek,A.B. and Hollenberg,S.M. (1995) Ras–Raf interaction: two-hybrid analysis. *Methods Enzymol.*, **255**, 331–342.
- Vojtek,A.B., Hollenberg,S.M. and Cooper,J.A. (1993) Mammalian Ras interacts directly with the serine/threonine kinase Raf. *Cell*, **74**, 205–214.
- Westendorf,J.M., Swenson,K.I. and Ruderman,J.V. (1989) The role of cyclin B in meiosis I. *J. Cell Biol.*, **108**, 1431–1444.
- Yang,J., Bardes,E.S.G., Moore,J.D., Brennan,J., Powers,M.A. and Kornbluth,S. (1998) Control of cyclin B1 localization through regulated binding of the nuclear export factor CRM1. *Genes Dev.*, **12**, 2131–2143.

Received August 6, 1999; revised and accepted January 21, 2000