



HAL
open science

^{13}C nuclear magnetic resonance studies of malate and citrate synthesis and compartmentation in higher plant cells

Elizabeth Gout, Richard Bligny, Nadine Pascal, Roland Douce

► **To cite this version:**

Elizabeth Gout, Richard Bligny, Nadine Pascal, Roland Douce. ^{13}C nuclear magnetic resonance studies of malate and citrate synthesis and compartmentation in higher plant cells. *Journal of Biological Chemistry*, 1993, 268 (6), pp.3986-3992. hal-02707364

HAL Id: hal-02707364

<https://hal.inrae.fr/hal-02707364>

Submitted on 1 Jun 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

¹³C Nuclear Magnetic Resonance Studies of Malate and Citrate Synthesis and Compartmentation in Higher Plant Cells*

(Received for publication, July 31, 1992)

Elizabeth Gout‡, Richard Bligny§, Nadine Pascal§, and Roland Douce¶

From the ‡Laboratoire de Résonance Magnétique en Biologie et Médecine and the §Laboratoire de Physiologie Cellulaire Végétale, Unité de Recherche Associée Centre National de la Recherche Scientifique 576, Département de Biologie Moléculaire et Structurale, Centre d'Etudes Nucléaires et Université Joseph Fourier, 85X F-38041 Grenoble Cédex, France

The synthesis of malate and citrate by sycamore cells (*Acer pseudoplatanus* L.) perfused with $\text{KH}^{13}\text{CO}_3$ was analyzed using ^{13}C NMR. To perform *in vivo* experiments, cells were compressed in a 25-mm tube and perfused with an arrangement enabling tight control of the circulating nutrient medium. An original method using paramagnetic Mn^{2+} that induced a complete loss of the vacuolar malate and citrate signals was developed to discriminate between cytoplasmic and vacuolar pools of malate and citrate. Our results indicated the following. (a) The accumulation of appreciable amounts of malate in sycamore cells required rather high (1 mM) concentrations of bicarbonate at all the pH values tested. (b) Malate was equally labeled at C-1 and C-4, suggesting that malate labeled at C-1 was produced by randomization of C-1 and C-4 by mitochondrial fumarase. Indeed, the separation of the intact organelles from the lysed protoplasts indicated that fumarase activity was essentially limited to the mitochondria. Similarly, citrate was equally enriched at C-1 and C-5 + C-6 carboxyls. (c) Malate appeared first in the cytoplasmic compartment; and when a threshold of cytoplasmic malate concentration was attained, malate molecules were expelled into the vacuole, where they accumulated. On the other hand, citrate accumulated steadily in the vacuole. Pulse-chase experiments demonstrated the central role played by the tonoplast in governing the vacuolar influx of citrate and the permanent exchange of malate between the cytoplasm and the vacuole.

P-enolpyruvate carboxylase occurs in all plants and catalyzes the carboxylation of P-enolpyruvate to oxalacetate, which in turn can be reduced to malate by malate dehydrogenase. The reaction catalyzed by P-enolpyruvate carboxylase is highly exergonic (the ΔG for this reaction is in the vicinity of 30 kJ mol^{-1}) (1, 2). Furthermore, Gout *et al.* (3) observed that any increase in cytoplasmic pH stimulates the synthesis of malate. Consequently, *in vivo* in the presence of $^{13}\text{CO}_2$, malate labeled at C-4 should be the product of carboxylation via P-enolpyruvate carboxylase. Chang and Roberts (4) and Stidham *et al.* (5) used ^{13}C NMR to observe the incorporation of $^{13}\text{CO}_2$ into malate by maize root tips and intact leaves of *Kalanchoë tubiflora*. Surprisingly, both groups observed a

significant incorporation of ^{13}C label into C-1 of malate and suggested therefore that malate labeled at C-1 was produced by randomization of C-1 and C-4 by mitochondrial fumarase. However, reports by two independent groups showed that rat liver (6) and *Saccharomyces cerevisiae* (7) contain two isozymes of fumarase that are localized in different intracellular compartments, the mitochondria and the cytosol. Consequently, it is possible that the unexpected incorporation of $^{13}\text{CO}_2$ into C-1 of malic acid was caused by equilibration with fumarate in the cytosol, where fumarase activity would exchange the label between C-1 and C-4.

In this investigation, we have used protoplasts from sycamore cells as a source of subcellular fractions and have concluded that in plant cells fumarase is confined to the mitochondria. In addition, ^{13}C NMR spectroscopy of intact sycamore cells was used to follow malate and citrate accumulation *in vivo*. One great advantage of ^{13}C NMR spectroscopy is that the resolving power of the technique allows for simultaneous identification and quantification of individual carbon atoms of the same molecule, in addition to distinguishing between molecules (8).

MATERIALS AND METHODS

Plant Material—The strain of sycamore (*Acer pseudoplatanus* L.) used in the study was grown as a suspension in a liquid nutrient medium according to the method of Bligny (9), except that Mn^{2+} was excluded to prevent excessive broadening of the resonance of vacuolar compounds (10). The cell suspensions were maintained in exponential growth by frequent subcultures. The cell wet weight was measured after straining culture aliquots onto a glass-fiber filter.

Preparation of Protoplasts—Washed cells (130 g, wet weight) were suspended in their culture medium containing 0.5 M mannitol, 10 mM Mops,¹ 1% (w/v) cellulase (Onozuka RS, Yakult Pharmaceutical Co., Nishinomiya, Japan), and 0.1% (w/v) pectolyase Y-23 (Seishin Pharmaceutical Co., Nishinomiya) adjusted to pH 5.7. The cells were incubated with constant shaking (20 cycles/min) at 25 °C. Digestion of young cells at 25 °C for 45 min resulted in a high yield of protoplasts. After digestion, the suspension was filtered through one layer of Miracloth (Krantex, Alfortville, France), which retained any undigested cell aggregates; and protoplasts were collected by centrifugation at $150 \times g$ for 10 min and then washed twice with 300 ml of suspension medium (0.5 M mannitol, 10 mM phosphate buffer (pH 7.5), 10 mM KCl, 5 mM MgCl_2 , 1% (w/v) polyvinylpyrrolidone (*M*, ~25,000, Serva), 0.1% (w/v) bovine serum albumin). Protoplasts were stored in suspension medium (600 ml) and were normally used within 1–2 h.

Gentle Rupture of Protoplasts and Separation of Organelles from Cytosolic Fraction—Since sycamore cell protoplasts have an average diameter of 20–30 μm , a rapid and effective procedure for the gentle rupture of intact protoplasts (*i.e.* for stripping the cell membrane) is to pass protoplasts through a fine nylon mesh (Nybolt PA, 20 μm)

* The costs of publication of this article were defrayed in part by the payment of page charges. This article must therefore be hereby marked "advertisement" in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.

¶ To whom correspondence should be addressed: Lab. de Physiologie Cellulaire Végétale, CENG, 85X F-38041 Grenoble Cédex, France. Tel.: 33-76-88-37-85; Fax: 33-76-88-50-91.

¹ The abbreviations used are: Mops, 4-morpholinepropanesulfonic acid; CDTA, *trans*-1,2-diaminocyclohexane-*N,N,N',N'*-tetraacetic acid.

affixed to the cut end of a 100-ml disposable syringe (11). Thus, if protoplasts (equivalent to 10 g, wet weight, 50 ml) are taken up and expelled through the 20- μ m nylon mesh, they will be completely ruptured. To separate the cytosolic fraction from the cell organelles, we subjected the broken protoplast fraction to centrifugation to yield a pellet largely free of cytosol and a supernatant enriched in cytosolic enzymes. Centrifugation was carried out in three steps (100 \times g for 5 min, 400 \times g for 5 min (RS-4 rotor, Kubota KN-70 centrifuge), and 20,000 \times g for 20 min (SS-34 rotor, Sorvall)). Each supernatant was centrifuged in a new tube, and the three successive pellets were combined together (cell organelles). This procedure ruptures all the protoplasts, leaving the fragile mitochondria largely intact.

Isolation of Mitochondria—The broken protoplasts were centrifuged for 10 min at 400 \times g in a table-top Kubota KN-70 centrifuge (RS-4 rotor) to remove the bulk of amyloplastids containing big starch grains. The supernatant was centrifuged for 20 min at 20,000 \times g (SS-34 rotor), and the mitochondrial pellet was resuspended in ~0.5 ml of suspension medium.

Perchloric Extract—For perchloric acid extraction, cells (9 g, wet weight) were quickly frozen in liquid nitrogen and ground to a fine powder with a mortar and pestle with 1 ml of 70% (v/v) perchloric acid. The frozen powder was then placed at -10 °C and subsequently thawed. The thick suspension thus obtained was centrifuged at 10,000 \times g for 10 min to remove particulate matter, and the supernatant was neutralized with 2 M KHCO₃ to pH ~6.5. The supernatant was then centrifuged at 10,000 \times g for 10 min to remove KClO₄; the resulting supernatant was lyophilized and stored in liquid nitrogen. For the NMR measurements, this freeze-dried material was redissolved in 2.5 ml of water containing 10% D₂O (perchloric acid extract).

The ¹³C NMR spectra of neutralized perchloric acid extracts were measured on a Bruker NMR spectrometer (AMX 400, narrow bore) equipped with a 10-mm multinuclear probe tuned at 100.62 MHz. Acquisition used 18- μ s pulses (60°) at 4-s intervals (spectra of carboxyl groups were unchanged when a pulse interval of 10 s was employed). Two levels of proton decoupling were used: 2.5 watts during the data acquisition (0.54 s) and 0.4 watts during the delay period (3.46 s). Spectra were acquired over a period of 2 h (1800 scans). Free induction decays were accumulated using 16,000 data points and zero-filled to 32,000 prior to Fourier transformation. A 1-Hz line broadening was applied. Chemical shifts were obtained by reference to the hexamethyldisiloxane resonance at 2.7 ppm. Spectra of standard solutions of known carbon compounds at pH 7.5 were compared with those of a perchloric acid extract of sycamore cells. The definitive assignments were made after running a series of spectra obtained by addition of the authentic compounds to the perchloric acid extracts (for an introduction to high-resolution NMR spectroscopy and its application to *in vivo* and *in vitro* studies, see Ref. 12).

In Vivo NMR Measurements—To get a better signal-to-noise ratio, an experimental arrangement was designed to analyze the maximum cell volume and to optimize the homogeneity of the cell incubation conditions. Cells (9 g, wet weight) were slightly compressed by hand between two polymer filters to a volume of 17–18 ml (13) and perfused under slight pressure at a flow rate of 50 ml min⁻¹ with a well-oxygenated (O₂-bubbling) nutrient medium (manganese-free culture medium containing 100 μ M phosphate) circulated via a 4-liter reservoir. The pH of this circulating medium (external pH (pH_e)) was controlled after passage through the compressed cells and was maintained constant in the reservoir using a pH-stat coupled to a titrimeter (Uretron 6, Tacussel, Lyon, France) to monitor the addition of HCl or KOH.

The ¹³C NMR spectra were measured on a Bruker spectrometer (AMX 400, wide bore) equipped with a dual ¹³C and ³¹P 25-mm probe tuned at 100.62 MHz. Acquisition used 30- μ s pulses (60°) at 4-s intervals (spectra of carboxyl groups were not significantly changed when longer intervals were chosen). Two levels of proton decoupling (Waltz pulse sequence) were used: 9.0 watts during the data acquisition (0.38 s) and 0.5 watts during the delay period (3.64 s). Spectra were acquired during periods ranging from 15 min to 2 h. Free induction decays were accumulated using 16,000 data points and zero-filled to 32,000 prior to Fourier transformation. A 5-Hz line broadening was applied. Chemical shifts were obtained by reference to hexamethyldisiloxane resonance at 2.7 ppm. Spectra of standard solutions of known carbon compounds at pH 7 were compared with those of perchloric acid extracts of sycamore cells as described previously (14). The definitive assignments were made after running a series of spectra obtained by addition of the authentic compounds to the perchloric acid extracts.

Perfusion in Presence of [¹³C]Bicarbonate—[¹³C]Bicarbonate (5

mM; isotopic enrichment of 99%) was added to the perfusion medium at time 0. The perfusate (1 liter) was recirculated. It was verified that due to cell respiration the isotopic enrichment was reduced to ~60% within 10 h.

Measurement of Sucrose, Malate, and Citrate—Sucrose, malate, and citrate were determined according to previous publications (15, 16).

Measurement of Marker Enzyme Activities—All assays were first performed on blanks to detect any unspecific drift or endogenous substrate or enzyme contamination. They were then optimized with respect to the concentration of each component and to the pH of the reaction mixture. We verified that, under these conditions, activities were linear with respect to time for at least 2 min and were proportional to the amount of protein. Fumarase (EC 4.2.1.2), ADP-glucose pyrophosphorylase (EC 2.7.7.27), catalase (EC 1.11.1.6), alcohol dehydrogenase (EC 1.1.1.2), and citrate synthase (EC 4.1.3.7) assays were performed according to previous publications (17, 18).

RESULTS

Intracellular Location of Fumarase—This was investigated in protoplasts from suspension cultures of sycamore cells. The following markers enzymes were used: mitochondria, citrate synthase; plastids, ADP-glucose pyrophosphorylase; peroxisomes, catalase; and cytosol, alcohol dehydrogenase. The gentle rupture of intact protoplasts passed through a fine nylon mesh followed by centrifugation carried out in three steps (see "Material and Methods") produced a supernatant that did not contain fumarase activity (Table I). The observation that almost all of the alcohol dehydrogenase activity was in the supernatant is consistent with the absence of fumarase in the cytosolic compartment. Furthermore, very little latency was found for alcohol dehydrogenase in carefully prepared lysates of protoplasts (data not shown), indicating that almost all the protoplasts had been ruptured. These results demonstrate that fumarase is confined within a membrane-bound cell organelle. To localize fumarase activity more precisely in the pellet containing cell organelles, intact amyloplastids and mitochondria were isolated from sycamore cells. As expected (data not shown), fumarase was not associated with sycamore cell amyloplastids and was confined within the mitochondria (~1100 nmol/min/mg of mitochondrial protein). These results together strongly support the unique location of the fumarase in mitochondria in plants in contrast to what was previously observed in yeast (7) and rat liver (6), where a substantial fraction of fumarase was found in the cytosol.

Accumulation of Citrate and Malate in Sycamore Cells during Utilization of [¹³C]Bicarbonate—*In vivo* ¹³C NMR spectra obtained under aerobic conditions at pH 6.5 (Fig. 1A) showed that the resonances of highest intensity corresponded to those

TABLE I

Subcellular localization of fumarase, ADP-glucose pyrophosphorylase (plastidial marker), alcohol dehydrogenase (cytosolic marker), and citrate synthase (mitochondrial marker) in mechanically ruptured sycamore cell protoplasts followed by centrifugation (20,000 \times g for 20 min)

Preparation of intact and broken protoplasts (total extract) and centrifugation of intact organelles were carried out as described under "Materials and Methods." These data are from a representative experiment and have been reproduced four times.

Enzyme	Total extract nmol/min/10 ⁶ cells	Distribution	
		Supernatant	Pellet
		% total activity	
Fumarase	32	0	100
ADP-glucose pyrophosphorylase	3	3	90
Catalase	520	10	85
Alcohol dehydrogenase	34	90	7
Citrate synthase	18	0	100




FIG. 1. Proton-decoupled ^{13}C NMR spectra (100.62 MHz) of compressed sycamore cells (*in vivo*) (A) and of their perchloric extracts (B). A, cells (9 g, wet weight) were packed in a 25-mm NMR tube as described under "Materials and Methods" and continuously perfused with a well-aerated manganese-free culture medium maintained at pH 6.5. The cell volume composed ~50% of the total (cell + perfusion medium) volume. To reduce the intracellular concentration of sucrose, the circulating medium contained 2.5 mM instead of 50 mM sucrose (we have verified that cell respiration and growth were not modified by lowering the sucrose concentration of the nutrient medium). The spectrum is the result of 1800 transients (2 h). B, perchloric extracts were prepared from 9 g of oxygenated cells, wet weight, as described in the text. The resolution of the carboxyl groups was considerably improved by the addition of 1 mM CDTA to the extracts. The samples (2.5 ml) containing 250 μl of $^2\text{H}_2\text{O}$ were analyzed for 2 h at 25 $^\circ\text{C}$. The carboxyl group area (left) and part of the amino acid methylene groups (right) are shown on expanded scales. Peak assignments are as follows: *suc*, succinate; *cit*, citrate; *iso*, isocitrate; *mal*, malate; *Asp*, aspartate; *Glu*, glutamate; *s*, sucrose.

of the glucosyl and fructosyl moieties of sucrose and were estimated to correspond to an intracellular level of $\sim 70 \mu\text{mol g}^{-1}$, wet weight, in good agreement with previous biochemical determination (14). The other major resonances in the chemical shift range of 50–40 ppm and those around 180 ppm arose from citrate and malate. The four resonances at 45.7, 76.3, 179.2, and 182.0 ppm coincided with those of C-2 + C-4, C-3, C-1 + C-5, and C-6 of citrate, respectively, whereas the two resonances at 43.2 and 71.1 ppm coincided with those of the methylene carbon (C-3) and the carbon atom with the hydroxy group (C-2) of malate, respectively. To obtain higher resolution and therefore more accurate quantification of the resonances, ^{13}C NMR spectroscopy was performed on tissue extracts (Fig. 1B). The supernatants from two homogenates were deproteinized with perchloric acid (16), and the pH dependencies of their ^{13}C spectra were examined independently. The chemical shifts obtained from this experiment (data not shown) matched precisely the corresponding data for citrate and malate. These results established unequivocally that the resonances at 45.7, 76.3, 179.2, 182.0, 43.2, and 71.1 ppm in spectra of intact sycamore cells arose from citrate and

malate. They also showed that the peaks at 181.7 and 180.5 ppm in the perchloric extracts at pH 7 arose from C-1 and C-4 (carboxyl groups) of malate, respectively. These two last peaks emerged from the background noise in the spectrum of intact sycamore cells incubated at alkaline pH (18). Interestingly, titration curves plotting chemical shift *versus* pH for citrate and malate in crude cell extracts (supernatant obtained after a 10-min centrifugation at $18,000 \times g$ following disruption of cells by freeze-thawing) indicated that the *in vivo* position of carboxyl groups corresponds to citrate and malate at pH 5.8. This suggests that the bulk of citrate and malate molecules accumulated in the vacuolar compartment.

Fig. 2 illustrates the changes that occurred in the sycamore cell spectrum when [^{13}C]bicarbonate (5 mM; isotopic enrichment of 99%) was added to the nutrient medium. In this experiment, the cells were maintained for >10 h in a continuously oxygenated circulating solution maintained at pH 7.5 (at this pH, the concentration of bicarbonate in the perfusion medium did not significantly decrease within 10 h). The major resonance at 161.2 ppm was attributable to bicarbonate present in the perfusion medium (pH 7.5) and the cytoplasmic compartment (pH 7.5), whereas the peak observed at 121.5 ppm was identified as the signal from free CO_2 gas present in all the cell compartments and the perfusion medium. Important changes occurred during the first 3 h. During this time, the ^{13}C resonances of C-1 and C-4 of malate increased considerably until a new steady state was obtained (Figs. 2–4). Since the chemical shifts of the carboxyl groups of malate are sensitive to pH <6.5 (5), ^{13}C NMR can distinguish the vacuolar malate pool at acidic pH from the cytoplasmic malate pool at the slightly alkaline pH as previously shown by Chang and Roberts (4). The signals from C-4 of malate in the cytoplasm and vacuole were clearly discernible, whereas C-1 signals from the two pools of malate partially overlapped (Fig. 3A). A close examination of an expanded scale of the spectra between 174 and 184 ppm indicates in fact that malate appeared first in the cytoplasmic compartment (the position of both carboxyl groups (C-1 and C-4) corresponded to malate at pH >7).

When 0.5 mM Mn^{2+} was added to the perfusion medium, this paramagnetic ion accumulated specifically in the vacuole (see Ref. 19) and, as already observed for vacuolar phosphate (19), suppresses the peaks of vacuolar carboxylates. It was therefore possible to observe separately for the first time the cytoplasmic malate directly and the vacuolar malate by the difference in the corresponding spectra obtained in the absence of Mn^{2+} (Fig. 3B). When a threshold of cytoplasmic malate concentration was attained, malate molecules were slowly expelled into the vacuole, where they accumulated, reaching a constant level after ~ 5 h (Fig. 4). Thus, when sycamore cells were treated with [^{13}C]bicarbonate for 1 h, the intensities of the signals from the C-1 and C-4 carboxyls of cytoplasmic malate were higher than those of vacuolar malate, whereas after a 3-h exposure with labeled bicarbonate, the intensities of the signals from carboxyls of vacuolar malate were much higher than those of cytoplasmic malate (Fig. 3). We have observed (data not shown) that the concentration of cytoplasmic malate attained at equilibrium (that is, when its rate of formation by the P-enolpyruvate carboxylase-malate dehydrogenase complex matches its rate of utilization) is strongly dependent on the activity of P-enolpyruvate carboxylase, which exhibits a rather low affinity for bicarbonate (Refs. 1 and 2 and see below). Figs. 2 and 4 also indicate that, after a lag, vacuolar citrate increased throughout the experiment because the chemical shifts matched those of citrate added to crude cell extracts at pH 5.8. It is interesting to note

FIG. 2. Representative proton-decoupled ^{13}C NMR spectra (100.6 MHz) of sycamore cells after addition of 5 mM $\text{H}^{13}\text{CO}_3^-$ in perfusion culture medium maintained at pH 7.5. Cells (9 g) packed in a 25-mm NMR tube as described for Fig. 1 and under "Materials and methods" were perfused with a well-aerated manganese-free culture medium containing 2.5 mM sucrose. At time 0, 5 mM $\text{H}^{13}\text{CO}_3^-$ was added to the perfusion culture medium. Spectra were obtained after 1, 2, 5, and 10 h. The pH of the culture medium was maintained constant at 7.5. The spectra were the result of 900 transients (1 h). *Inset*, perchloric extracts were prepared as described for Fig. 1 and in the text from cells incubated for 1 and 10 h in the presence of 5 mM HCO_3^- at pH 7.5. Spectra were obtained as described for Fig. 1.





FIG. 3. Representative proton-decoupled ^{13}C NMR spectra (100.62 MHz) of sycamore cells showing labeling of carboxylates after addition of 5 mM $\text{H}^{13}\text{CO}_3^-$ in perfusion culture medium maintained at pH 7.5. *A*, for experimental conditions, see Fig. 2 legend. 0h, standard spectrum; 1h, 2h, and 3h, spectra obtained after the indicated times of perfusion of the compressed cells with culture medium containing 5 mM $\text{H}^{13}\text{CO}_3^-$. Note that it is possible to discriminate between the C-4 peak of malate present in the cytoplasm (*cyt*) (pH 7.5) and that present in the vacuole (*vac*) (pH 5.7) at least during the first 2 h of incubation in the presence of $\text{H}^{13}\text{CO}_3^-$ at pH 7.5. However, the C-1 peak of cytoplasmic and vacuolar malate and C-6 of citrate were not clearly separated. *B*, experimental conditions were as described for *A*, except that 0.5 mM Mn^{2+} was added to the perfusion medium. Spectra were obtained after 15 min and 1, 2, and 3 h of perfusion of the compressed cells with culture medium containing 5 mM $\text{H}^{13}\text{CO}_3^-$ and 0.5 mM Mn^{2+} . The spectrum obtained at 15 min was the result of 225 transients; spectra obtained at 0, 1, 2, and 3 h were the result of 900 transients. Note that the addition of Mn^{2+} to the perfusion culture medium suppresses the peaks of vacuolar malate and citrate carboxylates (see Fig. 2A). The spectrum obtained at 2 h (*A - B*) is a difference spectrum of normal (without Mn^{2+}) (*A*) and Mn^{2+} -containing (*B*) cells obtained after 2 h in the presence of 5 mM $\text{H}^{13}\text{CO}_3^-$. Such a difference spectrum gives an accurate picture of the vacuolar citrate and malate carboxylates.

that when the accumulation of vacuolar malate ceased, signals from vacuolar citrate increased steadily. With time, other peaks appeared at 178.3, 175.2, and 175.5 ppm; these have been assigned to aspartate and glutamate carboxyl groups (the C-5 carboxyl of glutamate at 182 ppm was not labeled (Fig. 3B)). Titration curves plotting chemical shift versus pH

for aspartate and glutamate in crude cell extracts indicated that the position of carboxyl groups corresponded to glutamate and aspartate at pH >7. This suggests that these amino acids accumulated in the cytoplasmic compartment. This was also confirmed by the fact that added Mn^{2+} in the perfusion medium did not eclipse the peaks corresponding to these




FIG. 4. Time course evolution of cytoplasmic and vacuolar malate and citrate in sycamore cells perfused with culture medium containing 5 mM $\text{H}^{13}\text{CO}_3^-$ at pH 7.5. The concentrations of malate, citrate, and sucrose in the cell sample (average of the total sample within the detector) were determined either biochemically (15, 16) or by ^{13}C NMR analysis (13). The relative intensities of carboxylate peaks were determined by reference to the hexamethylsiloxane external standard. Experimental conditions were as described for Figs. 2 and 3, and the curves are from a representative experiment that has been reproduced five times. Note the unexpected decrease in intracellular sucrose. In fact, we have observed that at alkaline external pH (7.5) (see Ref. 3), the pH gradient across the plasma membrane is negligible, thus preventing the active transport of sucrose (or hexose) ($+\text{H}^+$) through the plasma membrane via a specific carrier. *cit*, citrate; *mal(cyt-vac)*, cytoplasmic and vacuolar malate.

carboxyls (resonances of aspartate and glutamate carboxyls are not detectable in the presence of Mn^{2+} ions).

When ^{13}C bicarbonate was used as a precursor and after subtraction of the control spectrum obtained with unlabeled bicarbonate to correct for natural abundance (see Fig. 1), C-1 and C-4 of malate present either in the cytoplasmic compartment or in the vacuole were labeled with almost equal probability (Fig. 3). Comparison with experiments performed in the presence of unlabeled bicarbonate demonstrated (data not shown and Ref. 3) that there was no label in the methylene carbon and the carbon with the hydroxy group of malate (C-2 and C-3) (some experiments were conducted for >10 h). This is well exemplified in a typical decoupled ^{13}C NMR spectrum of a perchloric acid extract of sycamore cells at pH 7.5 in the presence of 1 mM CDTA (Fig. 2). A similar pattern was observed for short-time exposures (10 min) with labeled bicarbonate (Fig. 3B). On the other hand, the citrate (C-1 + C-5) and C-6 signals increased with time and showed the same intensity, while under the same conditions, citrate C-2, C-3, and C-4 remained unlabeled.

Effect of Cytoplasmic pH and Bicarbonate Concentration on Rate of Malate and Citrate Synthesis by Sycamore Cells—Since phosphoenolpyruvate carboxylase has a relatively low affinity for bicarbonate and since the pH of the cytosol is unlikely to be as high as the pH optimum of P-enolpyruvate

carboxylase, we were prompted to examine the effect of cytoplasmic pH on the rate of malate accumulation in sycamore cells.

Table II indicates the effect of pH_c on the intracellular pH values in sycamore cells. In agreement with Fox and Ratcliffe (20), the cytoplasmic pH (pH_c) was independent of pH_e over the range 6–7.5 (see also Ref. 3). However, a loss of pH control was observed in response to the addition of 5 mM bicarbonate to the perfusion medium (Table II), especially when the external pH was acidic. Indeed, in the presence of 5 mM bicarbonate, pH_c fell below its original value to reach a new steady state. The difference between the original pH_c value and that attained after addition of 5 mM bicarbonate increased as the ΔpH across the plasma membrane increased (Table II). On the other hand, pH_c increased by up to 1 pH unit when external pH was increased from 7.5 to 9, irrespective of the presence of bicarbonate in the external medium (Table I). This observation strongly suggests that sycamore cells do not possess appropriate mechanisms to counteract the passive efflux of H^+ and/or the passive influx of OH^- when the outwardly directed H^+ gradient across the plasma membrane is reversed. Consequently, in the presence of 5 mM bicarbonate in the perfusion medium, pH_c increased progressively from 7.0 to 8.2 as pH_e increased from 6 to 9. We have therefore studied the effect of pH_c from 7 to 8.5 on the accumulation of intracellular malate and citrate in the presence of ^{13}C bicarbonate.

Table II indicates that in the absence of bicarbonate, the rate of malate and citrate accumulation in sycamore cells was negligible up to pH_e 7.5. Unexpectedly, at pH_e >7.5, the accumulation of organic acids remained negligible. Indeed, the presence of higher amounts of bicarbonate in the cytosolic compartment ($\log([\text{HCO}_3^-]/[\text{CO}_2]) = \text{pH} - \text{pK}'$ ($\text{pK}' = 6.4$)) would be expected to stimulate the activity of the P-enolpyruvate carboxylase. It is therefore possible that the large volume of the alkaline perfusion medium, which facilitates the rapid diffusion of the respiratory CO_2 , drains continuously the cell CO_2 content. In support of this suggestion, Table I indicates that the addition of 5 mM bicarbonate to the perfusion medium triggered a marked increase in the rate of malate and citrate synthesis at all of the pH values tested (the maximum rate of organic acid synthesis was already attained in the presence of 2 mM bicarbonate (data not shown)). It is noteworthy that the elimination of bicarbonate from the perfusion medium led to a progressive consumption of malate previously accumulated in the vacuole. Such a result strongly suggests that when the activity of P-enolpyruvate carboxylase declines owing to the collapse of cytosolic bicarbonate concentration, the net flux of malate toward the vacuole is stopped, and vacuolar malic acid diffuses across the tonoplast to be further metabolized in the cytoplasmic compartment.

In contrast, under the same conditions, citrate appeared to be very stable and was not further metabolized. To substantiate this, the flow of ^{13}C label in these experiments was further studied in a "chase" experiment carried out at pH 7.5, in which unlabeled bicarbonate was added to the sycamore cells after the elimination of ^{13}C bicarbonate from the perfusion medium. The time courses of the various signals obtained in this experiment are shown in Fig. 5. The time course shows that the vacuolar malate C-1 and C-4 peaks, which had built up in the presence of ^{13}C bicarbonate, decreased exponentially with a half-time of 3 h after the addition of unlabeled bicarbonate. Concurrent with these changes in the vacuolar malate peaks, ^{13}C enrichment at C-6 and C-1 + C-5 of citrate increased progressively. Interestingly, during the course of this chase experiment, unlabeled malate increased steadily from its initial value of 3 $\mu\text{mol/g}$, wet weight (Fig. 5).

TABLE II

Malate and citrate synthesis at different external pH values (pH_e) in the absence or presence of added bicarbonate (HCO₃⁻ + CO₂ = 5 mM)

The cytoplasmic pH values (pH_c) were determined as described by Gout *et al.* (3). It was verified that the pH_c values remained quite stable throughout the experiments. The total amounts of malate and citrate were measured from ¹³C NMR spectra after calibration with authentic compounds. The data are from a representative experiment and have been reproduced five times.

	pH _e									
	6		7		7.5		8		9	
CO ₂ + HCO ₃ ⁻ (mM)	0	5	0	5	0	5	0	5	0	5
pH _c	7.5	7.0	7.5	7.3	7.5	7.5	7.7	7.7	8.0	8.2
Synthesized malate (μmol/h/g cell, wet wt)	0	2.0	0	3.5	0	4.5	0.4	4.5	0.6	4.3
Synthesized citrate (μmol/h/g cell, wet wt)	0	4.0	0	3.6	0	3.3	0	3.3	0	3.0




FIG. 5. Time course evolution of vacuolar malate and citrate in sycamore cells during pulse-chase experiment carried out with 5 mM H¹³CO₃⁻/H¹²CO₃⁻. Experimental conditions were as described for Fig. 4, except that after 3 h of experiment, H¹³CO₃⁻ was replaced by H¹²CO₃⁻ (---). Note that during the chase, C-4 of malate (C4-mal) steadily decreased, whereas the total amount of vacuolar malate continuously increased. *cit*, citrate.

DISCUSSION

The results presented demonstrate that sycamore cells slightly compressed between two circular Teflon plates can survive for long periods as long as a well-aerated nutrient medium is pumped through the system under slight pressure. Such a system enables ¹³C NMR spectra of plant cells to be continuously recorded under various situations.

In good agreement with Fox and Ratcliffe (20), our results indicate that the cytoplasmic pH was independent of the carefully controlled external pH over the range 5.5–7.5. If one assumes that all biological membranes systems, including the plasma membrane, exhibit an intrinsic permeability to protons, we are forced to conclude that the steady-state value of ΔpH across the plasma membrane reflects a balance between the outward proton flux driven by the plasma membrane ATPase and the inward proton flux due to passive proton leakage. This means that up to pH 5.5, the efficiency of the H⁺ pump to react to back-leakage of protons was not limited. However, the loss of pH control observed in response to the addition of 5 mM HCO₃⁻ to the perfusion medium is very likely attributable to the fact that the equilibrium pH gradient

across the plasma membrane induces a rapid intake of CO₂ molecules (CO₂ diffuses very rapidly across all the biological membranes studied so far) into the cytoplasmic compartment, leading to a marked acidification of the cytoplasm ("acid load effect") (see Refs. 21 and 22). Under these conditions, the accumulation of HCO₃⁻ in the cytoplasm is considerably enhanced, and the distribution of HCO₃⁻ between the cytoplasm and the perfusion medium (which is predicted from the Henderson-Hasselbach equation) is inversely proportional to the distribution of protons. Under these conditions, the H⁺ pump does not work fast enough to counterbalance the massive CO₂ intake linked to the rapid production of protons in the cytoplasm. Interestingly, the fact that alkaline conditions also cause loss of cytoplasmic pH control is very likely attributable to a net flux of hydroxyl ions across the plasma membrane.

The response of intracellular pH to modifications of environmental factors has recently been reviewed by Kurkdjian and Guern (21). These authors consider the consumption (or synthesis) of organic acids such as malate, a process called biochemical pH-stat by Davies (23), as one of the mechanisms that allows the homeostasis of the cytoplasmic pH. Obviously, our results do not fit with this theory because the pool of malate attained at equilibrium in the cytosolic compartment is rather small, and, in addition, the bulk of this organic acid is excreted into the vacuolar space, where it accumulates. Furthermore, our results indicate that the synthesis of appreciable amounts of malate in the cytoplasmic and vacuolar compartments requires a rather high concentration of bicarbonate at all the external pH values tested. This raises the question of the regulation of cytoplasmic malate concentration in plant cells and, in particular, the problem of malate movement between the vacuole and the cytosol (and vice versa).

Apparently, it is the concentration of malate on both sides of the tonoplast membrane that governs the efflux or influx of malate. When a threshold of cytosolic malate concentration is attained, malate molecules are slowly expelled into the vacuole, where they accumulate. Conversely, when malate is no longer synthesized, malate concentration declines in the cytosol, leading to a slow efflux of malate from the vacuole. According to Lüttge (24) and Smith (25), plant cells accumulate large amounts of malic acid in the vacuoles following fixation of bicarbonate in the cytosol. The process has an overall stoichiometry of 2 mol of H⁺ accumulated per mol of malate and appears to be directly energized by the proton pumps of the tonoplast. This creates a potential gradient that is thought to drive the electrophoretic influx of malate²⁻ anions from the cytosol. Conversely, the efflux of the undissociated acid can be considered as a passive flux. Since during the course of the chase experiment unlabeled malate increased steadily, whereas the vacuolar [¹³C]malate decreased exponentially, we are forced to conclude that both efflux (passive?) and influx (via a specific carrier; for example, see Marigo *et al.* (26)) of malate²⁻ ions occur simultaneously across the

tonoplast *in vivo*. In other words, the tendency of malic acid molecules to flow down their concentration gradient is balanced by the electrogenic influx of malate²⁻ ions via the carrier.

In marked contrast, citrate ions, also entering the vacuolar space via a specific carrier (27), behave differently because they remain sequestered in the vacuole and removed from the equilibrium controlled by cytoplasmic enzymes. Indeed, the concentration of citrate and the labeled citrate carboxylate groups increased steadily throughout the experiment. These results indicate, in good agreement with a previous observation (14), that in these cells citrate exhibits a high metabolic inertness.

Our results demonstrate that C-1 and C-4 of malate become equally labeled when intact sycamore cells are treated with [¹³C]bicarbonate. This concurs with the findings of Stidham *et al.* (5) and Chang and Roberts (4), who inspected the incorporation of ¹³C from [¹³C]bicarbonate into malate in intact leaves of *K. tubiflora* and maize root tips. Likewise, the results of Osmond *et al.* (28) demonstrated that assimilation of ¹³CO₂ in the dark in a wide range of crassulacean acid metabolism plants leads to accumulation of a mixed population of [1-¹³C]- and [4-¹³C]malic acid (see also Ref. 29). Since the cytosolic compartment of sycamore cells is devoid of fumarase activity, these results strongly suggest that malate labeled at C-1 is produced by randomization of C-1 and C-4 by mitochondrial fumarase because fumarate is a symmetrical molecule, and it is assumed that the two CH groups in it would react identically. We are therefore forced to consider that *in vivo* a permanent movement of malate molecules between the mitochondrial matrix and the cytosol occurs probably via the dicarboxylate carrier of the inner mitochondrial membrane (30), catalyzing a strict counter-exchange of malate/malate. This implies that most of the product of the reaction catalyzed by mitochondrial fumarase is released into the bulk phase and is not necessarily "channeled" through a supramolecular complex bound to the inner mitochondrial membrane (metabolon) containing all the tricarboxylic cycle enzymes (31, 32).

If malate equally labeled on both carboxyls (because of scrambling through fumarate) is metabolized by the citric acid cycle via citrate synthase, we should see a ¹³C enrichment only at C-1 and C-6 of citrate. On the other hand, if newly formed oxalacetate molecules labeled exclusively at C-4 in the cytosolic compartment enter the matrix space via the specific oxalacetate carrier reported in all the plant mitochondria isolated so far (30), we should see a preferential labeling at C-6 of citrate (for an explanation, see Ref. 33). Our results indicate that incorporation of [¹³C]bicarbonate led to citrate highly labeled at C-1 and C-6, thereby confirming that *in vivo* labeled malate molecules synthesized by a sequence involving P-enolpyruvate carboxylase and malate dehydrogenase completely scrambled by mitochondrial fumarase are partly diverted into the tricarboxylic cycle to form citrate equally labeled at both C-1 and C-6. Such a situation is strongly favored because the cytoplasmic malate concentration at-

tained at the equilibrium is rather high (15–20 mM) (Fig. 4), and, in addition, endogenous controls polarizing the malate/citrate exchange can facilitate citrate efflux as previously shown with intact isolated mitochondria (30). In addition, citrate is very actively incorporated into the vacuole since no labeled citrate was observed in the cytoplasm. Unfortunately, our results cannot discriminate between pyruvate formed by the glycolytic pathway and that produced by the mitochondrial NAD⁺-linked malic enzyme (30) because both pathways lead to unlabeled acetyl-CoA, which condenses with oxalacetate to form citrate.

Finally, the results presented in this paper illustrate the resolving power of ¹³C NMR spectroscopy in distinguishing between individual carbon atoms of appropriate metabolites such as malate and citrate. This makes it possible to study intact cells as they respond to physiological perturbations.

REFERENCES

1. Latzko, E., and Kelly, G. J. (1983) *Physiol. Veg.* **21**, 805–813
2. O'Leary, M. H. (1982) *Annu. Rev. Plant Physiol.* **33**, 297–315
3. Gout, E., Bligny, R., and Douce, R. (1992) *J. Biol. Chem.* **267**, 13903–13909
4. Chang, K., and Roberts, J. K. M. (1989) *Plant Physiol. (Bethesda)* **89**, 197–203
5. Stidham, M. A., Moreland, D. E., and Siedow, J. N. (1983) *Plant Physiol. (Bethesda)* **73**, 517–520
6. Suzuki, T., Sato, M., Yoshida, T., and Tuboi, S. (1989) *J. Biol. Chem.* **264**, 2581–2586
7. Wu, M., and Tzagoloff, A. (1987) *J. Biol. Chem.* **262**, 12275–12282
8. Roberts, J. K. M., and Jardetzky, O. (1981) *Biochim. Biophys. Acta* **639**, 53–76
9. Bligny, R. (1977) *Plant Physiol. (Bethesda)* **59**, 502–505
10. Martin, J.-B., Bligny, R., Rébeillé, F., Douce, R., Leguay, J. J., Mathieu, Y., and Guern, J. (1982) *Plant Physiol. (Bethesda)* **70**, 1156–1161
11. Leegood, R. C., and Walker, D. A. (1983) in *Isolation of Membranes and Organelles from Plant Cells* (Hall, J. L., and Moore, A. L., eds) pp. 185–210, Academic Press Limited, London
12. Pfeffer, P. E., and Gerasimowicz, W. V. (1989) in *Nuclear Magnetic Resonance in Agriculture* (Pfeffer, P. E., and Gerasimowicz, W. V., eds) pp. 3–70, CRC Press, Inc., Boca Raton, FL
13. Roby, C., Martin, J.-B., Bligny, R., and Douce, R. (1987) *J. Biol. Chem.* **262**, 5000–5007
14. Genix, P., Bligny, R., Martin, J.-B., and Douce, R. (1990) *Plant Physiol. (Bethesda)* **94**, 712–722
15. Rébeillé, F., Bligny, R., Martin, J.-B., and Douce, R. (1985) *Biochem. J.* **226**, 679–684
16. Journet, E. P., Bligny, R., and Douce, R. (1986) *J. Biol. Chem.* **261**, 3193–3199
17. Journet, E. P., and Douce, R. (1985) *Plant Physiol. (Bethesda)* **79**, 458–467
18. Lunn, J. E., Droux, M., Martin, J., and Douce, R. (1990) *Plant Physiol. (Bethesda)* **94**, 1345–1352
19. Roby, C., Bligny, R., Douce, R., Tu, S.-I., and Pfeffer, P. E. (1988) *Biochem. J.* **252**, 401–408
20. Fox, G. G., and Ratcliffe, R. G. (1990) *Plant Physiol. (Bethesda)* **93**, 512–521
21. Kurkdjian, A., and Guern, J. (1989) *Annu. Rev. Plant Physiol. Plant Mol. Biol.* **40**, 271–303
22. Mathieu, Y., Guern, J., Pean, M., Pasquier, C., Beloeil, J.-C., and Lallemand, J.-Y. (1986) *Plant Physiol. (Bethesda)* **82**, 846–852
23. Davies, D. D. (1986) *Physiol. Plant. (Bethesda)* **67**, 702–706
24. Lüttge, U. (1987) *New Phytol.* **106**, 593–629
25. Smith, J. A. C. (1987) in *Plant Vacuoles: Their Importance in Solute Compartmentation in Cells and Their Applications in Plant Biotechnology* (Marin, B., ed) pp. 79–87, Plenum Press, New York
26. Marigo, G., Bouysson, H., and Laborie, D. (1988) *Bot. Acta* **101**, 187–191
27. Rentsch, D., and Martinoia, E. (1991) *Planta (Berl.)* **184**, 532–537
28. Osmond, C. B., Holtum, J. A. M., O'Leary, M. H., Roeske, C., Wong, O. C., Summons, R. E., and Avadhani, P. N. (1988) *Planta (Berl.)* **175**, 184–192
29. Kalt, W., Osmond, C. B., and Siedow, J. N. (1990) *Plant Physiol. (Bethesda)* **94**, 826–832
30. Douce, R. (1985) *Mitochondria in Higher Plants: Structure, Function, and Biogenesis*, Academic Press, New York
31. Srere, P. A. (1987) *Annu. Rev. Biochem.* **56**, 89–124
32. Cornish-Bowden, A. (1991) *Eur. J. Biochem.* **195**, 103–108
33. Stryer, L. (1988) *Biochemistry*, W. H. Freeman & Co., New York