



Détection par Elisa du virus de la marbrure de la vigne (GFkV = grapevine fleck virus)

Bernard Walter, Pascal Cornuet

► To cite this version:

Bernard Walter, Pascal Cornuet. Détection par Elisa du virus de la marbrure de la vigne (GFkV = grapevine fleck virus). *Agronomie*, 1993, 13 (7), pp.651-657. hal-02715940

HAL Id: hal-02715940

<https://hal.inrae.fr/hal-02715940>

Submitted on 1 Jun 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Elisa detection of grapevine fleck virus (GFkV)

B Walter, P Cornuet

INRA Colmar, Laboratoire de Pathologie Végétale, Vigne et Vin, BP 507, F68021 Colmar, France

(Received 3 February 1993; accepted 12 May 1993)

Summary — The association of grapevine fleck virus (GFkV) with fleck symptoms on grapevine was established using Elisa. The results confirmed that GFkV is probably the virus responsible for fleck. Its detection in vines by Elisa was not reliable throughout the vegetative cycle: under our conditions, in Alsace, June–July was found to be the best period for testing.

grapevine / virus / fleck / Elisa / etiology

Résumé — **Détection par Elisa du virus de la marbrure de la vigne (GFkV = grapevine fleck virus).** L'association du Grapevine fleck virus (GFkV) avec les symptômes de la marbrure de la vigne a été établie par Elisa. La détection par Elisa du GFkV dans les plants de vigne n'est pas possible pendant la totalité du cycle végétatif : dans nos conditions, en Alsace, les mois de juin et juillet constituent la meilleure période pour le dépistage.

vigne / virus / marbrure / Elisa / étiologie

INTRODUCTION

Detection of pathogens which are transmissible through the wood (viruses, mollicutes, bacteria) is an essential step in the selection of the grapevine (Walter, 1991a). More than 30 major virus and virus-like diseases of grapevines have been described (Bovey and Martelli, 1992). Among them, nepoviruses and closteroviruses are well characterized and can be detected by Elisa or molecular hybridization techniques. The virus-like disease are detected by grafting onto indexing varieties (Walter, 1991b).

Only recently a phloem-limited non-mechanically transmissible isometric virus was demonstrated to be associated with grapevine fleck disease (Boscia *et al*, 1991a,b).

Fleck disease is latent in European grapevine varieties and in most American rootstocks. The following symptoms develop on *Vitis rupestris*

used as indexing variety: clearing of veinlets and translucent spots, wrinkling and upward curling of the leaves. The disease can strongly affect root development of rootstocks as well as their grafting ability (Triolo and Materazzi, 1986). No vector is known and dissemination is primarily through infected propagation material. Seed transmission does not occur. The disease can be eliminated by heat therapy or meristem tip culture (Goheen and Luhn, 1973; Ottenwaelter *et al*, 1973; Barlass *et al*, 1982).

In France, sanitary certification presently requires freedom from fanleaf, leafroll, corky bark, *Rupestris* stem pitting and Kober stem grooving for rootstocks and *V. vinifera*. In addition, fleck has to be absent from rootstock varieties.

In the present paper we describe the purification of grapevine fleck virus (GFkV) from roots and leaves of infected grapevines and the development of Elisa for routine detection of the virus.

MATERIALS AND METHODS

Source material

Different *Vitis* spp accessions obtained from INRA Vassal were used for purification of the virus. After preliminary assays, *V. vinifera* cv Bogaskere (accession Y200) was chosen. Y200 indexed positive on *V. rupestris* and was also infected by vein mosaic and vein necrosis but was free from fanleaf and arabis mosaic viruses, leafroll, corky bark, *Rupestris* stem pitting and Kober stem grooving.

For Elisa, we used other *Vitis* spp from collections (INRA Vassal, INRA Côtmar, INRA Bordeaux) and, in some cases, plants obtained after heat treatment and *in vitro* culture of shoot apices of fleck-diseased mother plants (Bass and Legin, 1981).

Indexing

Detection of fleck disease was made by grafting onto *Vitis rupestris* du Lot, either by using woody cuttings or by green grafting as described by Walter *et al* (1990).

Virus purification

The purification procedure was basically the same as that described by Boulila *et al* (1990) using fresh young and fast growing rootlets or leaves of greenhouse-forced cuttings of the donor vine. After polyethyleneglycol precipitation (PEG 6.000) and high-speed centrifugation, the virus suspension was centrifuged through different types of gradients.

Sucrose gradients were formed by 3 layers of 15, 25 and 35% sucrose in citrate buffer 0.02 M pH 6.1, left to diffuse overnight at 4°C; centrifugation was for 150 min at 24 000 rpm in a Beckman SW 28 rotor at 5°C.

Nicodenz (Sigma) gradients were constituted of 4 layers, each 1 ml, of 60, 50, 40 and 30% with an upper layer of 6 ml 5% Nicodenz in citrate buffer, centrifuged for 18 h at 31 000 rpm at 5°C in Beckman SW 41 tubes.

Cesium sulfate was used at 1.4 g/cm³ in citrate buffer and centrifuged in a Beckman SW 41 rotor for 17 h at 35 000 rpm at 5°C.

The virus was obtained from the fractions of the gradients after dialysis in citrate buffer and high-speed centrifugation (Beckman 50.2 Ti, 2 h, 250 000 g, 5°C).

Electron microscopy

Carbon-coated pioloform (Agar Scientific R 1275) grids were floated for 15 min on a drop of virus sus-

pension. After desiccation, staining was made on a drop of 2% aqueous uranyl acetate. The preparations were observed with a Hitachi HU 11 CS microscope.

Serology

The first assays were made with an antiserum obtained from GP Martelli (University of Bari, Italy).

To prepare antisera, purified virus preparations from sucrose or Nicodenz gradients were emulsified with complete (for the first injection) or incomplete (for booster injections) Freund's adjuvant. Rabbits were injected subcutaneously at 7-d intervals. Antiserum collection began 14 d after the third injection. Immunoglobulins were purified by the rivanol precipitation method (Hardie and Van Regenmortel, 1977). Conjugation of purified immunoglobulins with alkaline phosphatase or biotin was made using 0.06% glutaraldehyde (Avrameas, 1969).

Detection of GFkV in grapevines

Fresh leaves or cortical scrapings were crushed in the presence of 5 vol Tris-HCl 0.2 M buffer pH 8.2 containing 2% polyvinylpyrrolidone, 0.8% NaCl and 0.05% Tween 20. In some cases, this extract was serially diluted from 10⁻¹ to 10⁻⁵ in the same buffer. For Elisa, coating immunoglobulins at 0.25 µg/ml were incubated for 3 h at 37°C. After overnight incubation of the plant extract at 4°C, an immunoglobulin-biotin conjugate (in some cases pre-adsorbed with 5% v/v healthy grapevine extract) was added for 2 h at 38°C. Detection was with a streptavidin-alkaline phosphatase conjugate (Interchim Jackson) at 1/10 000 for 30 min and paranitrophenylphosphate at 1 mg/ml at 38°C.

RESULTS

Virus purification

Elisa (fig 1) and EM observations of sucrose gradient fractions showed that 2 zones were enriched with GFkV particles. An intermediate step in the purification consisting of clarification with ether (1 v/1 v) or butanol (8.5%; v/v) after PEG precipitation and before high-speed centrifugation, yielded more highly purified virus suspensions.

Virus particles were paraspherical, ≈ 30 nm in diameter, some appearing empty, others apparently intact (fig 2).




Fig 1. Elisa detection of GFkV in sucrose gradient fractions. For the purification, leaf extract was PEG-precipitated, clarified with butanol and centrifuged at 200 000 g for 90 min. The pellet was resuspended in citrate buffer 0.02 M pH 6.1 and 1-ml suspension (corresponding to 250 g leaves) was centrifuged in a Beckman SW 28 rotor at 24 000 rpm for 150 min at 5°C through a gradient obtained after overnight diffusion of 3 layers of 35, 25 and 15% sucrose in citrate buffer. Centrifugation was from left to right. Each gradient fraction was diluted 1/50 in Tris-HCl buffer. Data are OD readings at 405 nm, after 1-h incubation of the substrate.

Association of GFkV with fleck symptoms

A very good correlation was found between the presence of fleck symptoms following indexing on *V. rupestris* and positive Elisa response: 25 accessions from various geographical origins, which included *V. vinifera* and rootstock hybrids, were positive in both tests; 95 were negative in both; 3 were positive by indexing but negative by Elisa and 6 were positive by Elisa but negative by indexing.

Plants obtained after *in vitro* culturing of shoot tips from heat-treated vines (65 *V. rupestris* and 23 *V. vinifera* Muscat d'Alexandrie) had been found to be free from fleck after > 30 d treatment at 35°C (Bass and Legin, 1981). The absence of fleck as revealed by absence of symptoms (table I) or negative indexing (table II) was highly correlated with the absence of GFkV in Elisa. Only 2 plants of cv Muscat d'Alexandrie indexing negative on *V. rupestris* after 50 d heat treatment were positive in Elisa (table II). This difference may be due to a lower sensitivity of indexing compared to Elisa, or to a natural reinfection of these 2 plants that were indexed immediately after planting outdoors, whereas they were checked by Elisa some years later.




Fig 2. Electron micrograph of a NycoDenz gradient fraction enriched in GFkV after staining with aqueous uranyl acetate. Bar: 100 nm (photo courtesy of G Stocky).

Distribution of virus in plants

In the first experiment in May, very young leaves and cortical scrapings of *V. vinifera* Muscat d'Alexandrie and *V. rupestris* were extracted and

Table I. Presence of GFkV in *Vitis rupestris* with and without symptoms after increasing periods of heat treatment at 35°C.

Heat treatment (d)	Symptoms	Elisa
0	21+ 3–	21+ 3–
26	9+	9+
156	13–	13–
184	6–	6–
382	13–	13–

Table II. Presence of GFkV in *Vitis vinifera* (Muscat d'Alexandrie) that indexed positive or negative on *V. rupestris* after increasing periods of heat treatment at 35°C.

Heat treatment (d)	Index/ <i>V. rupestris</i>	Elisa
21	3+	3+
36	2+ 2-	2+ 2-
50	4+ 9-	4+ 7- 2+
426	3-	3-

sap dilutions from 10^{-1} to 10^{-5} were assayed in Elisa. For Muscat, the 10^{-5} dilution of leaf extracts consistently gave a positive reaction, whereas cortical scraping extract only reacted up to 10^{-3} . In contrast, leaves of *V. rupestris* reacted to a dilution of 10^{-1} and cortical scrapings up to 10^{-2} . These differences were confirmed when comparing optical density (OD) readings (table III).

In the second experiment, half of a Muscat d'Alexandrie plant was harvested on September 30, 1992 and cortical scrapings, petioles and leaves were analyzed by Elisa. The sample con-

Table III. Distribution of GFkV in young leaves and wood shavings harvested in May. Data are Elisa OD readings of various dilutions of the extracts.

<i>V. vinifera</i> Muscat d'Alexandrie		<i>V. rupestris</i> du Lot	
Leaves	Wood	Leaves	Wood
Fleck			
10^{-1}	>2.000 1.673	0.503	0.956
10^{-2}	>2.000 0.778	0.092	0.399
10^{-3}	1.128 0.203	0.046	0.099
Control			
10^{-1}	0.045 0.068	0.094	0.122

sisted of a main branch from which cortical scrapings were harvested near the beginning of 4 1-yr-old branches that were numbered 1 to 4 from the bottom to the top of the main branch. From each 1-yr-old cane, cortical scrapings, petioles and leaves were collected from the bottom, middle and apical portions (table IV). OD readings ranged from 0.348 to 1.243 in leaf extracts but no conclusion could be drawn about the distribution of the virus in the plant. In petioles and cortical scrapings, Elisa readings were more consistent.

In the third experiment (October 10, 1992) leaves, petioles and cortical scrapings from the bottom, middle and top parts of 4 canes of *Vitis rupestris* were analyzed (table V). Cortical scrapings did not show significant variations, whereas petioles and leaves gave increasing readings from the bottom to the top of the canes.

Table IV. Distribution of GFkV in leaves, petioles and wood shavings of different portions of 1-yr-old canes and main branch of *Vitis vinifera* Muscat d'Alexandrie.

Portion of cane	Cane 1	Cane 2	Cane 3	Cane 4
Top				
Leaves	1.045	0.348	0.590	1.013
Petioles	0.671	0.618	0.701	0.853
Wood	1.000	0.700	0.806	0.963
Middle				
Leaves	1.077	1.015	1.043	0.944
Petioles	1.004	0.868	0.769	0.867
Wood	0.887	0.954	1.040	0.903
Bottom				
Leaves	0.779	0.813	0.746	1.243
Petioles	0.929	0.782	0.917	0.777
Wood	1.014	1.090	0.965	0.901
Main branch	1.082	0.971	0.689	-

Data are Elisa OD readings of 1/10 extracts after 1-h incubation of nitrophenylphosphate. Healthy controls: leaves = 0.023; petioles = 0.016; wood = 0.015. Buffer control = 0.013.

Table V. Distribution of GFkV in leaves, petioles and wood shavings of different portions of 1-yr-old canes of *Vitis rupestris* du Lot.

Portion of cane	Cane 1	Cane 2	Cane 3	Cane 4
Top				
Leaves	0.715	0.847	0.850	0.885
Petioles	0.582	0.800	0.558	0.578
Wood	0.756	0.891	0.754	0.771
Middle				
Leaves	0.779	0.752	0.589	0.529
Petioles	0.516	0.482	0.377	0.439
Wood	0.635	0.848	0.825	0.779
Bottom				
Leaves	0.298	0.398	0.458	0.429
Petioles	0.503	0.371	0.408	0.476
Wood	0.852	0.958	0.796	0.789

Data are Elisa OD readings of 1/10 extracts after 1-h incubation of nitrophenylphosphate. Healthy controls: leaves = 0.098; petioles = 0.045; wood = 0.008. Buffer control = 0.007.




Fig 3. Variation of Elisa end-point dilution for the detection of GFkV over the vegetative period. For detection of GFkV in 3 *V. vinifera* Muscat d'Alexandrie plants (A) and 3 *V. rupestris* plants (B), leaves were harvested at various dates from May 7th to October 14th. Extracts were diluted from 10⁻¹ to 10⁻⁵ as described in the *Materials and Methods*. The height of the diagrams corresponds to the dilution end-point of extracts reacting in Elisa.

Variation of Elisa response during growth

Large variations in Elisa response were noticed when leaves of vines were analyzed during the entire vegetative period.

The Elisa response decreased rapidly during the hotter summer period, especially in the case of *V. vinifera* Muscat d'Alexandrie (fig 3). Under our conditions, the sensitivity of Elisa was maximal during June and July for *V. vinifera* and *V. rupestris*. This leads to the conclusion that certain periods should be avoided for reliable Elisa detection of GFkV in our climatic conditions: the beginning of the vegetative period for *V. rupestris* and late summer (August and September) for *V. vinifera* (fig 4).




Fig 4. Variation of Elisa response (OD 405 nm) in leaf extract of grapevine during the vegetative period. OD variations in Elisa of a 10⁻¹ dilution of the extracts of leaves of *V. vinifera* Muscat d'Alexandrie (A) and *V. rupestris* (B) harvested at various dates during the vegetative period. Data are OD at 405 nm after 30 min incubation of the substrate for diseased (open circles) and healthy (dark triangles) vines.

DISCUSSION

In this paper, we confirm that GFkV — an isometric virus previously described by Boulila *et al* (1990) and Boscia *et al* (1991b) — is associated with the disease known as fleck (marbrure) and so probably the causal agent of it. Though Koch's postulates have not yet been fully confirmed, we demonstrated a high correlation between the presence of fleck symptoms and the presence of GFkV, both in various grapevine accessions from different geographical origins and in heat-treated vines.

In a series of experiments we have analyzed some factors which could influence reliable detection of GFkV for a routine diagnosis in grapevines, *ie* when, during the year, a vine should be indexed and what type of tissue should be sampled.

In general, leaves and cortical scrapings were comparable as antigen sources for Elisa; under our conditions, there were no significant differences when comparing tissues from the top or bottom of *V. vinifera* (Muscat d'Alexandrie). Only petioles and leaves from *V. rupestris* showed increasing Elisa readings when harvested from lower to upper parts of the canes. In contrast, Boscia *et al* (1991b) reported that Elisa readings were highest in the basal leaves of LN 33 and Primitivo di Gioia.

Studying Elisa responses in some *V. vinifera* and *V. rupestris*, we can conclude that the best period for routine testing is, under our conditions, June and July. The dramatic decrease in Elisa readings during the hottest summer period may bring the response very close to the background level.

Considerable seasonal fluctuations have been reported in the efficiency of detection of nepoviruses in grapevines: in some cases, virus detection was impossible in the hottest periods of the year and was most reliable during intensive growth stages (Kölber *et al*, 1985) or at full flowering (Bovey *et al*, 1980; Kölber *et al*, 1981; Lehoczky *et al*, 1983, 1984; Rüdel *et al*, 1983). Under other conditions, detection of nepoviruses was possible by Elisa in grapevine leaves during the whole vegetative period (Walter and Etienne, 1987).

ACKNOWLEDGMENTS

Grateful thanks are expressed to Prof Martelli and co-workers for supplying the first antiserum, and also to Prof Van Regenmortel for revising the manuscript.

REFERENCES

- Avrameas S (1969) Coupling of enzymes to proteins with glutaraldehyde. Use of the conjugates for the detection of antigens and antibodies. *Immunochemistry* 6, 43-52
- Barlass M, Skene KGM, Woodham RC, Krake LR (1982) Regeneration of virus-free grapevines using *in vitro* apical cultures. *Ann Appl Biol* 101, 292-295
- Bass P, Legin R (1981) Thermo-thérapie et multiplication *in vitro* d'apex de vigne. Application à la séparation ou à l'élimination de diverses maladies de type viral et à l'évaluation des dégâts. *CR Acad Agric Fr* 67, 922-933
- Boscia D, Martelli GP, Savino V, Castellano MA (1991b) Identification of the agent of grapevine fleck disease. *Vitis* 30, 97-105
- Boscia D, Savino V, Martelli GP, Castellano MA (1991a) Association of the phloem-limited non-mechanically transmissible isometric virus with grapevine fleck disease. In: *Proc 10th Meet ICVG* (IC Rumbo, R Bovey, D Gonsalves, WB Hewitt, GP Martelli, eds) Ores Publishing, Volos 1990, 173-174
- Boulila M, Boscia D, Di Terlizzi B, Castellano MA, Minafra A, Savino V, Martelli GP (1990) Some properties of a phloem-limited non-mechanically-transmissible grapevine virus. *J Phytopathol* 129, 151-158
- Bovey R, Brügger JJ, Gügerli P (1980) Detection of fanleaf virus in grapevine tissue extract by enzyme-linked immunosorbent assay (Elisa) and immune electron microscopy (IEM). In: *Proc 7th Meeting ICVG*. Vineland Res Station Canada Agric Branch, Niagara Falls, 259-275
- Bovey R, Martelli GP (1992) Directory of Major Virus and Virus-Like Diseases of Grapevines. Description, Historical Review and Bibliography. MFCIC, ICVG, pp 111
- Goheen AC, Luhn C (1973) Heat inactivation of viruses in grapevines. *Riv Patol Vég Ser IV* 9, 287-289
- Hardie G, Van Regenmortel MHV (1977) Isolation of specific antibody under conditions of low ionic strength. *J Immuno Methods* 15, 305-314
- Kölber M, Beczner L, Pacsa S, Lehoczky J (1985) Detection of grapevine chrome mosaic virus in field-grown vines by Elisa. *Phytopathol Mediterr* 24, 135-140
- Kölber M, Lehoczky J, Pacsa S, Kobza S (1981) Detection of grapevine fanleaf virus (GFV) with Elisa technique in the leaf samples collected in various phenophases and in deep-frozen leaf extracts. *Kertgazdaság* 13, 11-22
- Lehoczky J, Kölber M, Beczner L, Pacsa S (1984) Distribution of the grapevine chrome mosaic disease in Hungary and detection of its virus (GCMV) in the leaves of outdoor vines by Elisa technique. *Kertgazdaság* 16, 41-52
- Lehoczky J, Kölber M, Farkas G (1983) Effect of spring and summer high temperature on detection

- of grapevine fanleaf virus and yellow mosaic virus with Elisa. *Kertgazdaság* 15, 37-45
- Ottenwaelter MM, Hevin M, Leclair P, Doazan JP, Rives M (1973) Heat therapy eliminates the ability to transmit the causal agent of marbrure in several *V. vinifera* clones and in *V. rupestris* du Lot (St George). *Riv Patol Veg Ser IV* 9, 281-285
- Rüdel M, Alebrand M, Altmayer B (1983) Untersuchungen über den Einsatz des Elisa-Tests zum Nachweis verschiedener Rebviren. *Wein Wissenschaft* 38, 177-185
- Triolo E, Materazzi A (1986) La Maculatura Infettiva della vite: influenza di isolati diversi sull'attitudine alla propagazione vegetativa di *Vitis rupestris* St George. In: IV Symp Int Sélection Clonale de la Vigne, Changins, 1986. *Rech Agron Suisse* 3 (26), 320-324
- Walter B (1991a) Sélection de la vigne : le dépistage des maladies de la vigne transmissibles par les bois et plants. *Bull OIV* 727-728, 691-701
- Walter B (1991b) New or improved procedures for the detection and identification of viruses or agents of virus-like diseases of grapevine and for diagnosis. In: *Proc 10th Meet ICVG* (IC Rumbos, R Bovey, D Gonsalves, WB Hewitt, GP Martelli, eds) Ores Publishing, Volos, 1990, 226-238
- Walter B, Bass P, Legin R, Martin C, Vernoy R, Collas A, Vesselle G (1990) The use of a green-grafting technique for the detection of virus-like diseases of the grapevine. *J Phytopathol* 128, 137-145
- Walter B, Etienne L (1987) Detection of the grapevine fanleaf viruses away from the period of vegetation. *J Phytopathol* 120, 355-364