^{13}C -NMR characterization of formaldehyde bonds in model mixtures and proteins containing lysine

D. Tome, N. Naulet, A. Kozlowski

► To cite this version:

D. Tome, N. Naulet, A. Kozlowski. ^{13}C -NMR characterization of formaldehyde bonds in model mixtures and proteins containing lysine. *International Journal of Peptide and Protein Research*, 1985, 25, pp.258-266. hal-02722002

HAL Id: hal-02722002

<https://hal.inrae.fr/hal-02722002>

Submitted on 1 Jun 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Inventaire
N°

DUB
72

¹³C-n.m.r. characterization of formaldehyde bonds in model mixtures and proteins containing lysine

D. TOME,* N. NAULET** and A. KOZLOWSKI*

*National Institute for Agricultural Research – Laboratory of Animal Feed Technology;

**University of Nantes – Organic Chemistry, ERA C.N.R.S. 315, Nantes, France

Received 12 April, accepted for publication 3 July 1984

Reactions involving formaldehyde, lysine and another amino acid (tyrosine, arginine, asparagine, glutamine or cysteine) were studied by ¹³C-n.m.r. analysis. The products of the crosslinking between lysine and tyrosine were found to be acid-resistant and were isolated using ion exchange chromatography. With the other selected amino acids, the crosslinked products with lysine are acid-labile and could not be isolated; however, n.m.r. allows the determination of their structure. In addition, the formation of methyllysine and of formyllysine have been observed. The ¹³C-n.m.r. chemical shifts of the formaldehyde derived hydroxymethyl, methylene, methyl and formyl carbons with the amino acids side chains help in the interpretation of ¹³C-enriched formaldehyde treated protein ¹³C-n.m.r. spectra. With BSA, we have observed the formation of hydroxymethyllysine, hydroxymethylhistidine, hydroxymethyl-asparagine or -glutamine, methyllysine, and a crosslink between lysine and arginine. These results show the great importance of lysine in the treatment of protein with formaldehyde. Use of ¹³C-n.m.r. is considered highly suitable for obtaining more knowledge about formaldehyde binding to peptides and proteins.

Key words: amino acids; carbon 13 n.m.r.; formaldehyde; lysine; protein

Formaldehyde treatment of proteins, which is of importance in a number of procedures (1–4), can affect different chemical groups and particularly the ε-amino group of lysine (5, 6). It is known that this latter group reacts with formaldehyde, resulting in *N*-hydroxymethyl (7–9), *N*-methyl (8), *N*-formyl (10, 11). The *N*-hydroxymethyl may then react on other nucleophilic sites, leading to the formation of methylene bridges; the amino acids involved have been tentatively identified as arginine (12, 13), tyrosine, histidine or tryptophan (4, 14), asparagine or glutamine (12, 15) and cysteine (16).

The above scheme can explain most of

the modifications produced in peptides and proteins by formaldehyde treatment, but the mechanism by which they proceed has not yet been definitively established. The main drawbacks hindering identification of the reaction products are binding reversibility and low concentration. These difficulties can be overcome by use of ¹³C-enriched formaldehyde and ¹³C-n.m.r. spectroscopy (17). However, before obtaining a single step ¹³C-n.m.r. identification of the ¹³C-formaldehyde proteins bonds, we need to know the ¹³C-chemical shifts of the formaldehyde derived hydroxymethyl, methylene, methyl or formyl carbons of the different reactive groups suspected to be

involved in the reaction. Some knowledge has already been obtained using model compounds (13, 18, 19) and amino acids (7, 8, 16, 17).

This paper describes ^{13}C -n.m.r. characterization of the bonds of formaldehyde or ^{13}C -formaldehyde in model systems containing lysine or protected lysine in the presence of various protected amino acids. The obtained results allow interpretation, at least to some extent, of the ^{13}C -n.m.r. spectra of ^{13}C -formaldehyde treated proteins.

EXPERIMENTAL PROCEDURES

Amino acids and N^α acetyl amino acids (protected amino acids) were obtained from Sigma and used without further purification. Bovine serum albumin (BSA) (Sigma) was extracted with hexane prior to use. A 10-M solution of formaldehyde was prepared by depolymerisation of pure paraformaldehyde (Merck) in distilled water at 85° . A 6.5-M solution of ^{13}C -formaldehyde (91% enriched) was obtained from Prochem.

Different model systems were designed in which either lysine and one protected amino acid (e.g. lysine and N^α acetylglutamine), or two protected amino acids (e.g. N^α acetyllysine and N acetyltyrosine) were allowed to react either with formaldehyde or with ^{13}C -formaldehyde. The experiments were carried out in aqueous phase using different pH, time and temperature conditions. The amino acid concentration was kept in the 0.1–0.5 M range and the molar ratio of the added formaldehyde or ^{13}C -formaldehyde to amino acids was fixed at 1–4. pH adjustments were made with concentrated HCl or NaOH. "Acid-labile" formaldehyde (Fh) was steam-distilled off after addition of *o*-phosphoric acid (0.5 ml for 3 ml of solution).

The compounds A1 and A2 (Table 1) were prepared and isolated as follows: protected lysine (0.5 M) and protected tyrosine (0.5 M) were mixed with formaldehyde (0.5 M) or ^{13}C -formaldehyde (0.5 M) and allowed to stand at pH 7.5 and 37° for 1 week. Acid-labile formaldehyde was first eliminated, then deacetylation of amino groups was performed by hydrolysis in 6 M hydrochloric acid for 24 h at 110° . The hydrolysate was then concen-

trated under reduced pressure at 40° and fractionated by ion exchange chromatography on a column (60×1.6 cm) of Dowex 50X4 in pyridine acetate buffer 0.3 M, pH 4.7. The column was regenerated by a pyridine acetate buffer 2.23 M, pH 5.4. The detection was carried out on an aliquot of the individual fractions by absorbance measurement at 560 nm after reaction with ninhydrin. The isolated compounds were concentrated under reduced pressure at 40° in order to get rid of the pyridine acetate, and redissolved in water.

BSA was used at a concentration of 100 g/l in sodium phosphate buffer 0.1 M, pH 7.5, or in sodium acetate buffer 0.2 M, pH 4.5. The protein was treated with 0.33 M ^{13}C -formaldehyde. Free ^{13}C -formaldehyde was eliminated by means of dialysis against phosphate buffer.

^{13}C -n.m.r. spectra were recorded in the pulsed Fourier Transform mode on a Bruker WH 90 spectrometer operating at 22.635 MHz, in 10 mm tubes. The field-frequency lock was provided by deuterated benzene (C_6D_6) put into a coaxial tube, and ^{13}C chemical shifts measured with respect to the central peak of C_6D_6 and calculated from TMS according to the relation: $\delta^{13}\text{C}(\text{C}_6\text{D}_6) = \delta^{13}\text{C}(\text{TMS}) + 127$ p.p.m. Typical running conditions were: 6000 Hz bandwidth, acquisition time 0.679 s, 30° flip angle without pulse delay, quadrature detection, broad band noise decoupling. Conditions for quantitative measurement were: 90° flip angle; pulse delay 200 s, broad band noise decoupling only during acquisition. Resonance lines were assigned by comparing experimental data with those previously reported in the literature. The use of resonance decoupling and of the APT pulses sequence (Attached Proton Test) helped in this assignment.

The nomenclature used is that of the IUPAC-IUB Commission on Biochemical Nomenclature [*European J. Biochem.* (1975) 53, 1–14].

RESULTS AND INTERPRETATION

In a preliminary study, mixtures of two protected amino acids with a reactive side chain (7) were allowed to react in the presence of

TABLE 1

¹³C-n.m.r. chemical shifts (TMS) of the compounds identified in model mixtures of amino acids in the presence of formaldehyde. (a) N^α acetyl amino acids; (b) after deacetylation; (c) asparagine or glutamine in the N^α acetyl form. (1) confirmed by the use of ¹³C-enriched formaldehyde; (2) see in the text

	Origin of the carbons																		
	Formaldehyde (1)				Lysine				Tyrosine, arginine, asparagine or glutamine										
	CH ₂ OH	CH ₂	CH ₃		α	β	γ	δ	ε	α	β	γ	δ	ω	1	2	3	4	5
Lysine (a)	—	—	—	54.9	31.1	22.3	26.7	39.8	—	—	—	—	—	—	—	—	—	—	—
N ^ε , hydroxymethyllysine (a)	71.7	—	—	54.9	31.3	22.6	25.2	49.3	—	—	—	—	—	—	—	—	—	—	—
N ^ε , methyllysine (b)	—	—	33.1	54.3	31.2	22.5	24.3	48.5	—	—	—	—	—	—	—	—	—	—	—
Tyrosine (a)	—	—	—	—	—	—	—	—	56.8	30.9	—	—	—	—	129.3	130.6	115.8	155.2	115.8
C3, hydroxymethyl-tyrosine (a)	60.0	—	—	—	—	—	—	—	56.8	30.9	—	—	—	—	129.1	130.0	126.9	153.7	116.1
A1 N ^ε lysine, C3 tyrosine, methane (b)	47.1	—	—	54.7	30.0	21.6	25.1	47.0	56.2	35.5	—	—	—	—	126.9	132.3	118.3	154.8	116.2
A2 N ^ε , N ^ε lysine (C3 tyrosine methane) ₂ (b)	—	55.3	—	54.5	30.0	21.4	23.2	53.6	56.3	35.4	—	—	—	—	127.1	132.7	117.3	154.4	115.9
A3 (2)	81.2–49.3	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Arginine (a)	—	—	—	—	—	—	—	—	55.1	29.1	24.9	41.1	157.2	—	—	—	—	—	—
N ^ω , hydroxymethyl-arginine (a)	65.5	—	—	—	—	—	—	—	55.1	29.1	24.9	41.2	156.4	—	—	—	—	—	—
B1 N ^ε lysine, N ^ω arginine, methane (a)	—	59.8	—	54.9	31.1	22.2	26.5	49.5	55.1	28.8	24.9	41.1	152.4	—	—	—	—	—	—
B2 (2)	66.3	59.1	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
B3 (2)	64.7–64.1	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Asparagine (a)	—	—	—	—	—	—	—	—	50.5	37.1	173.4	—	—	—	—	—	—	—	—
N ^γ , hydroxymethyl-asparagine (a)	63.2	—	—	—	—	—	—	—	50.4	37.7	173.4	—	—	—	—	—	—	—	—
C1 N ^ε lysine, N ^γ asparagine methane (a)	51.8	—	—	54.7	30.2	21.9	25.1	44.8	50.9	37.7	173.9	—	—	—	—	—	—	—	—
C2 N ^α lysine, N ^γ asparagine methane (c)	—	51.6	—	59.7	29.3	21.9	26.8	39.7	50.9	37.7	174.4	—	—	—	—	—	—	—	—
Glutamine (a)	—	—	—	—	—	—	—	—	53.0	26.8	31.6	173.2	—	—	—	—	—	—	—
N ^δ hydroxymethyl-glutamine (a)	63.2	—	—	—	—	—	—	—	53.0	26.9	32.2	173.2	—	—	—	—	—	—	—
D1 N ^ε lysine, N ^δ glutamine methane (a)	—	51.9	—	54.7	30.1	21.9	25.4	44.8	53.0	27.2	31.7	173.2	—	—	—	—	—	—	—
D2 N ^α lysine, N ^δ glutamine methane (c)	—	51.8	—	59.8	29.3	21.9	26.7	39.7	53.4	27.2	31.7	173.2	—	—	—	—	—	—	—

formaldehyde. In most cases, only the reaction products of each side chain with formaldehyde were traced on the ^{13}C -n.m.r. spectra (7-9). In several cases, however, new reactions were observed, four of which, resulting in the formation of methylene bridges between cysteine and respectively lysine, arginine, asparagine and glutamine, have already been described in a previous paper (16). Other reactions, considered in this work, result in the formation of methylene bridges between lysine and tyrosine, arginine, asparagine or glutamine (Table 1). These reactions which involve lysine seem also predominant when BSA is treated with ^{13}C -formaldehyde.

Reactions between lysine, tyrosine and formaldehyde

In acidic medium, protected tyrosine does not react with formaldehyde even if protected lysine is added. However, at pH higher than 6, the phenol ring becomes much more reactive.

In a system containing protected tyrosine and formaldehyde left at pH 7.5 and 37° for several days, a hydroxymethyl derivative is formed by addition of formaldehyde on the carbon 3 of the ring. This derivative is acid-labile and lost through formaldehyde steam-distillation. The reaction is never completed and not very favoured. If, under the same experimental conditions, protected lysine is added to the mixture in an equimolar ratio, different products are formed. Three of them are acid-resistant and have been isolated using ion exchange chromatography following deacetylation (Fig. 1); they have been identified by n.m.r. as methyllysine and products A1 and A2. ^{13}C -n.m.r. spectra carried out under quantitative conditions gave evidence of a ratio of 1:1 between lysine and tyrosine in A1 and 1:2 in A2. A third product A3 is characterized by two peaks at 81.6 and 49.5 p.p.m., deriving from formaldehyde. These two peaks always show the same intensity and they increase in intensity with an excess of formaldehyde; these products do not withstand the acid-labile formaldehyde elimination step and disappear with a subsequent increase of the peak at 47.3 p.p.m. (A1). Therefore, it was assumed that product A3 could result from a cyclisation in A1 through a methylene bridge between the

FIGURE 1

Elution profile from Dowex 50X4 ion exchange chromatography of the reaction products between protected lysine, protected tyrosine and formaldehyde (after deacetylation). The mixture (1 ml) was applied to the column (60×1.6 cm) and the fractions were eluted with pyridine acetate 0.3 M, pH 4.7 at a flow rate of 60 ml/min. ^{13}C -n.m.r. analysis of each peak: 1-tyrosine; 2-lysine + methyllysine; 3-product A1; 4-product A2

nitrogen of lysine side chain and the oxygen of the phenol ring of tyrosine.

After 5 days at pH 7.5 and 37° , the solution still contains free formaldehyde and approximately 70% unreacted amino acids, 13% A1, 7% A2, 7% A3 and 3% methyllysine.

When an excess of formaldehyde and lysine is used, A2 decreases and A1, A3 and methyllysine increase. In addition, it was shown, with the help of ^{13}C -formaldehyde, that dimethyllysine, with the methyl carbons at 42.8 p.p.m., and formyl lysine, characterized by the formyl carbons at 164.8 p.p.m., were built up. These latter products are found at low concentrations.

Reaction between lysine, arginine and formaldehyde

Protected arginine reacts with formaldehyde yielding an hydroxymethyl derivative on the guanidyl function, characterized by the ^{13}C -n.m.r. peak of the hydroxymethyl carbon at 65.2 p.p.m. This derivative is the only one which is produced in a large range of pH and

represents nearly 35% of arginine when reagents are initially mixed in an equimolar ratio.

The reactions between protected lysine, protected arginine and formaldehyde are pH dependent. In acidic medium (pH 2–4), only the hydroxymethyl derivative of each side chain can be traced on the ^{13}C -n.m.r. spectra (hydroxymethyl carbons at 71.3 and 65.2 p.p.m.); when the pH is higher than 4, three new products B1, B2 and B3 can be detected after several hours at room temperature.

The crosslinked product B1, which shows a methylene bridge between the side chain of lysine and arginine at 59.9 p.p.m., is always found in the largest amount. Its concentration increases when pH is increased with a concomitant disappearance of hydroxymethyllysine.

Product B2 is characterized by two methylene carbons deriving from formaldehyde at 66.3 and 59.1 p.p.m. These peaks are always of the same intensity and make up less than 5% of the mixture. In light of the recorded chemical shifts, B2 can be considered a compound resulting from an hydroxymethylation of the free nitrogen of the guanidyl function of B1.

The product B3 is characterized by two methylene carbons at 64.2 and 64.8 p.p.m. deriving from formaldehyde. These peaks are always of the same intensity and make up about 3% of the products. They have not yet been identified and several hypotheses can be put forth. The most likely, if we consider their chemical shifts, is the formation of a dimethylene ether bridge between the side chains of lysine and arginine.

All these derivatives are acid labile and break up during the steam-distillation step. Therefore, we did not succeed in isolating them; however, if the reacting medium is acidified at room temperature, just after the derivative formation, B1 can still be found in the mixture. This gives some evidence of a relative stability of the methylene bridge.

After reaction of protected lysine, protected arginine and formaldehyde in an equimolar ratio at pH 7.5 and 37° for 1 day, it was found that formaldehyde had been entirely consumed and the mixture was approximately made up of 15% unreacted amino acids, 15% hydroxymethyl arginine, 60% B1, 10% B2 and B3.

Reactions between lysine, asparagine or glutamine and formaldehyde

The amide groups of protected asparagine and glutamine slowly react with formaldehyde to yield the hydroxymethyl derivatives (7).

In the presence of protected lysine, after several days at room temperature and in acidic medium only, the product of the cross-linking by a methylene bridge between the amide group of asparagine or glutamine and the ϵ -amino group of lysine (products C1 and D1) can be traced on the ^{13}C -n.m.r. spectra. The methylene signal is at 51.8 p.p.m. If protected lysine is replaced by lysine, a secondary reaction is also observed on the α -amino group with the methylene bridge at 51.6 p.p.m. (C2 and D2). These amide-formaldehyde-amine condensation products are unstable and easily decompose when pH is higher than 5; therefore they cannot be isolated.

The proportions of the reaction products when the equilibrium is reached are the same for both amino acids. After reaction of an equimolar mixture of the three reagents at pH 2.5 and 37° for several days, only 20% of the formaldehyde has been used and the solution contains 75% unreacted amino acids, 5% hydroxymethyl-asparagine or -glutamine, 5% hydroxymethyllysine and 10% of the crosslinked products (C1 or D1).

Reactions between ^{13}C -formaldehyde and BSA

Typical ^{13}C -n.m.r. spectra obtained from BSA treated with ^{13}C -enriched formaldehyde, either in acid (pH 4.5) or neutral (pH 7.5) solution, are shown in Fig. 2.

The intensified resonances occur in a frequency range 30–80 p.p.m. which is the region of the hydroxymethyl, methylene and methyl carbons resulting from the reaction of formaldehyde with amino acids side chains. The intensified resonances are not completely resolved because of an important line broadening. Moreover, due to the very low concentration on each bond, a long accumulation time is needed in order to obtain spectra. Denaturation of the modified protein would probably increase the resolution; however, all the attempts carried out either by thermal treatment or by adding chemical agents, resulted in precipitation of the protein. Despite these difficulties, five intensified peaks for which it is possible to

FIGURE 2

90-MHz ^{13}C -n.m.r. spectra of ^{13}C -formaldehyde treated BSA at 37° 1-pH 4.5, 10 h; 2-pH 4.5., 1 week; 3-pH 7.5, 1 week; 4-pH 7.5, 1 week, dialyzed against water. Each spectrum results from 60 000 transients obtained by using the 8K-frequency domain data point (spectral width 6,000 Hz; recycle time 0.679 s.; flip angle 30°). F-peaks coming from the ^{13}C -enriched formaldehyde solution (formaldehyde monomer, 82.3 p.p.m.; methanol, 49.3 p.p.m.). P-peaks coming from the protein. a, b, c, d, e, i, i': ^{13}C -formaldehyde - BSA links (see in the text).

postulate some structure are subsequently designated a, b, c, d, e.

Both in acidic and neutral medium, the first peak appears after some hours at 71.8 p.p.m. (a), which is the chemical shift of hydroxymethyllysine and hydroxymethylhistidine (7). In neutral medium only, another *N*-hydroxymethyl is formed at 70.2 p.p.m. (b) near the value of a *N*-terminal hydroxymethyl (8). These bonds decrease as the reaction goes on and disappear when ^{13}C -formaldehyde is eliminated by dialysis. The reversibility of these linkages is in accordance with the properties of the *N*-hydroxymethyls.

Within a few days, a broad band appears between 50 and 70 p.p.m. In acidic medium its formation is accompanied by a gelification of the solution, preventing any obtention of resolved spectra, and the only additional identified peak is at 63.1 p.p.m. (c), which is exactly the chemical shift of the amide hydroxymethyl on asparagine or glutamine. In neutral medium, the products remain soluble; in that case, in addition to the aforementioned amide hydroxymethyl at 63.1 p.p.m. (c), there is another peak at 60.2 p.p.m. (d) very near the chemical shift of a lysine-arginine crosslink (59.9 p.p.m.) or of a tyrosine hydroxymethyl (60 p.p.m.). A well resolved peak appears, after several days, in neutral medium, at 33.1 p.p.m. (e); considering this chemical shift, it may be assumed that the compound is the *N*-methyllysine. The peaks c, d, e build up progressively during the reaction and are not affected when ^{13}C -formaldehyde is eliminated by dialysis. At last, the two signals i and i', around 50 p.p.m., are very weak and badly resolved; their assignment needs further investigation.

Identical spectra have been obtained with other proteins (lysozyme, β -lactoglobulin, ribonuclease).

DISCUSSION

From the above results and those of previous works, we are able to predict the ^{13}C -n.m.r. chemical shifts of several formaldehyde-derived hydroxymethyl, methylene, methyl and formyl carbons which may occur during the reaction between formaldehyde and proteins (Table 2).

TABLE 2

¹³C-n.m.r. chemical shifts (TMS) of the formaldehyde derived formyl (R-CHO), hydroxymethyl (R-CH₂OH), methyl (R-CH₃) and methylene (R-CH₂-R') carbons with the amino acids side chains

System	Formyl	Hydroxymethyl	Methylene	Methyl
Lysine	164.8	71.9	—	33.1–42.8 ^d
Serine ^a	—	89.1	—	—
Histidine ^a	—	71.7	—	—
Alanine ^b	—	69.5	—	—
Tryptophane ^a	—	68.4	—	—
Cysteine	—	65.7	—	—
Arginine	—	65.2	—	—
Asparagine	—	63.2	—	—
Glutamine	—	63.2	—	—
Tyrosine	—	60.0	—	—
Lys:Arg	—	—	59.9	—
Lys:Cyst ^c	—	—	52.7–57.9 ^d	—
Lys:Asn	—	—	51.2	—
Lys:Gln	—	—	51.2	—
Lys:Tyr	—	—	47.1–55.3 ^d	—
Cyst:Arg ^c	—	—	43.8	—
Cyst:Gln ^c	—	—	41.4	—
Cyst:Asn ^c	—	—	41.4	—

^aRef. 7.

^bModel for *N*-terminal, from ref. 8.

^cRef. 16.

^dDisubstituted on lysine nitrogen.

Hydroxymethyl carbons absorb in the frequency range 60–90 p.p.m. and, with the exception of sulfhydryl, amide and phenol function, form according to an equilibrated reaction (7, 9, 20, 21). *N*-hydroxymethylation is the most prominent on lysine side chain but also occurs on α -amino groups; no evidence for so-called Schiff-base formation was obtained (8, 22). Moreover, spontaneous methylation and formylation may proceed when lysine and formaldehyde are in excess, leading to *N*-methyllysine, *N*-*N'*-dimethyllysine and *N*-formyllysine (8, 10, 11). The formation of hydroxymethyls on amino groups (lysine and *N*-terminal) and of methyllysine, when BSA is treated by ¹³C-formaldehyde, is consistent with these observations. In addition, hydroxymethylations on histidine and amide groups which have also been suspected to act (23–25), are confirmed by the present results. On the other hand, hydroxymethylation on the phenol

ring of tyrosine, which is unfavoured, probably will not occur.

The observed methylene bridges, formed between amino acids side chains, implicate lysine or cysteine which are able to link either together, or with arginine, asparagine or glutamine. In addition, lysine reacts with the phenol ring of tyrosine. Identification of some of these cross-linked systems brings support to hypotheses previously put forth (4, 12, 14); however, the mechanism involved in their formation is not yet clearly understood. For instance, it seems surprising not to record lysine-lysine or cysteine-cysteine bridges. The ¹³C-n.m.r. chemical shifts of the methylene carbons in a variety of cross-linked systems are between 40 and 60 p.p.m. The experimental values are in agreement with those predicted for lysine-arginine and lysine-amide systems (13, 18), but somewhat at lower field for lysine-tyrosine (18). In addition, the de-

shielding effect of the substitution of another methylene on the nitrogen of lysine side chain, in lysine-tyrosine and lysine-cysteine systems, is consistent with the effects previously reported for model systems of the same kind (26, 27).

The methylene bridge formed between lysine and arginine side chains, which was found to occur very easily in model mixtures, presumably represents the major crosslink in formaldehyde-BSA reaction. Although this bridge is acid labile, it is relatively stable. The n.m.r. study enables the visualization of such a bridge, previously proposed as well in model mixtures as in proteins (12, 13), but difficult to observe by other techniques.

The only acid resistant methylene bridge occurs between lysine and tyrosine. Its formation in model mixtures, probably through a Mannich reaction (28), is now definitely established; moreover a disubstitution on the nitrogen leads to the linkage of two tyrosine molecules on the lysine side chain. From our experiment using BSA, there is no evidence that this lysine-tyrosine bridge, often suspected to constitute most of the irreversibly bound formaldehyde on proteins (4), was formed. Actually it probably occurs at low concentration and its detection may be hindered because of the short spin lattice relaxation time (T_1) of this methylene type (19).

The other bridges are more unstable. The favoured lysine-cysteine crosslink may occur (16) but rarely happens due to the lack of cysteine which is mostly in the cystine form in the studied proteins; however, formaldehyde may induce, in some cases, the reduction of cystine with liberation of reactive cysteine. The lysine-amide crosslink, previously postulated (12) could play a role in acidic medium and partially explains the observed gelification. Finally cysteine-arginine and cysteine-amide bridges have also been observed in model mixtures (16), but are very unfavoured.

CONCLUSION

According to current studies, lysine possesses a highly reactive ϵ -amino group allowing various covalent reactions with formaldehyde. Our ^{13}C -n.m.r. experiments describe the

structure of several hydroxymethyl, methyl and methylene derivatives which are linked to this amino group in model mixtures. The ^{13}C -enriched formaldehyde treatment of proteins proved useful to point out the formation of methyllysine and a lysine-arginine crosslink as major products. However, further experiments are needed in order to identify definitively any of the components produced during the reaction between proteins and formaldehyde.

REFERENCES

1. Gustavson, K.H. (1956) *The Chemistry of Tanning Processes*. Academic Press, New York
2. Walker, J.F. (1964) *Formaldehyde*. Robert E. Krieger, Huntington, NY
3. Zelter, S.Z., Leroy, F. & Tissier, J.F. (1970) *Ann. Biol. Anim. Biochim. Biophys.* 10, 111–121
4. Bizzini, B. & Raynaud, M. (1974) *Biochimie* 56, 297–303
5. Davis, P. & Tabor, B.E. (1963) *J. Polymer Sci. A1*, 799–815
6. Tome, D., Bertrand, D., Viroben, G. & Delort-Laval, J. (1979) *Ann. Technol. Agric.* 28, 299–318
7. Tome, D. & Naulet, N. (1981) *Int. J. Peptide Protein Res.* 17, 501–507
8. Tome, D., Naulet, N. & Martin, G.J. (1982) *J. Chim. Phys.* 79, 361–366
9. Naulet, N., Tome, D. & Martin, G.J. (1983) *Org. Magn. Reson.* 21, 564–566
10. Tyihak, E., Trezl, L. & Rusnak, I. (1980) *Pharmazie* 35, 18–20
11. Trezl, L., Rusnak, I., Tyihak, E., Szarvas, T. & Szende, B. (1983) *Biochem. J.* 214, 289–292
12. Fraenkel-Conrat, H. & Olcott, H.S. (1948) *J. Am. Chem. Soc.* 70, 2673–2684
13. Taylor, S.K., Davidson, F. & Ovenall, W. (1978) *Photogr. Sci. Eng.* 22, 134–138
14. Fraenkel-Conrat, H. & Olcott, H.S. (1948) *J. Biol. Chem.* 174, 827–843
15. Koga, D., Imoto, T., Nakamura, Y., Fujikawa, S. & Yagishita, K. (1978) *Agric. Biol. Chem.* 42, 1147–1156
16. Naulet, N. & Tome, D. (1984) *J. Chim. Phys.* 81, 173–178
17. Jentoft, J.E., Jentoft, N., Gerken, T.A. & Dearborn, D.G. (1979) *J. Biol. Chem.* 254, 4366–4370
18. Dewar, M.K., Johns, R.B., Kelly, D.P. & Yates, J.F. (1975) *Aust. J. Chem.* 28, 917–924
19. Kelly, D.P., Dewar, M.K., Johns, R.B., Shao Wei-Let & Yates, J.F. (1977) in *Protein Cross-*

- linking, *Biochemical and Molecular Aspects* (Friedman, M., ed.), pp. 641–647, Plenum Press, New York
20. French, O. & Edsall, J.T. (1945) *Advan. Protein Chem.* **2**, 277–355
21. Feraud, P.F. & Le Henaff, Ph. (1968) *Bull. Soc. Chim.* **5**, 1868–1978
22. Kitamoto, Y. & Maeda, H. (1980) *J. Biochem.* **87**, 1519–1530
23. Fraenkel-Conrat, H., Cooper, M. & Olcott, H.S. (1945) *J. Am. Chem. Soc.* **67**, 950–954
24. Martin, C.J. & Marini, M.A. (1967) *J. Biol. Chem.* **242**, 5736–5743
25. Dunlop, P., Marini, M., Fales, H.M., Sokoloski, E. & Martin, C.J. (1973) *Bioorg. Chem.* **2**, 235–247
26. De Breet, A.J.J., Dankeelman, W., Huysmans, W.G.B. & De Wit, J. (1977) *Angew. Makromol. Chem.* **62**, 7–31
27. Tomita, B. & Hatono, S. (1978) *J. Polymer Sci.* **16**, 2509–2525
28. Short, J. H. & Ours, W. (1975) *J. Heterocycl. Chem.* **12**, 869–876

Address:

D. Tome

National Institute for Agricultural Research (INRA)
Laboratory of Animal Feed Technology
Rue de la Géraudière
44072- Nantes Cedex
France