



HAL
open science

Relationships between genetic markers and production traits in pigs

Christine Renard, Jean Pierre Bidanel, A. Palovics, M. Vaiman, Gérard Guérin, J.P. Runavot

► To cite this version:

Christine Renard, Jean Pierre Bidanel, A. Palovics, M. Vaiman, Gérard Guérin, et al.. Relationships between genetic markers and production traits in pigs. *Annales de zootechnie*, 1988, 37, pp.205. 10.1051/animres:19880345 . hal-02723640

HAL Id: hal-02723640

<https://hal.inrae.fr/hal-02723640>

Submitted on 10 Aug 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Evaluation in pure and crossbreeding of the Large-White hyperprolific line in multiplier and commercial herds

G. PETIT ⁽¹⁾, J.P. RUNAVOT ⁽¹⁾, J. GRUAND ⁽²⁾, C. LEGAULT ⁽³⁾

(1) *Institut Technique du Porc, Pôle Amélioration de l'Animal, B.P. 3, 35650 Le Rheu*

(2) *Institut National de la Recherche Agronomique, Station Expérimentale de Sélection Porcine, 86480 Rouille*

(3) *Institut National de la Recherche Agronomique, Station de Génétique quantitative et appliquée, 78350 Jouy-en-Josas*

In 49 multiplier and commercial herds, reproductive performance of 92 purebred Large-White and 169 Large-White × Landrace crossbred females sired by 8 Large-White hyperprolific (H) boars was compared with that of contemporary females from the same genetic types (126 purebreds and 266 crossbreds), born in the same herd and reared in the same conditions.

Purebred and crossbred daughters of H boars showed a definite advantage in prolificacy + 0.96 and + 0.61 in total number of piglets at birth, + 0.87 and + 0.52 in number of piglets born alive. As no systematic recording system had been implemented regarding transfer of piglets between litters, information on litter size at weaning is not fully reliable. No significant difference is observed for age at first farrowing, interval between farrowings and interval from weaning to fertile mating.

The comparison of the two types of progeny for traits recorded in the « on-the-farm » programme (age and backfat thickness at 100 kg liveweight) shows a slight genetic delay of the H line in production traits. The overall economic interest of using the H line is discussed.

Relationships between genetic markers and production traits in pigs

Christine RENARD ⁽¹⁾, J.P. BIDANEL ⁽²⁾, G. PETIT ⁽³⁾, Agnès PALOVICS ⁽⁴⁾,
M. VAIMAN ⁽⁵⁾, G. GUERIN ⁽⁴⁾, J.P. RUNAVOT ⁽³⁾

(1) *Institut National de la Recherche Agronomique, L.R.A., 78350 Jouy-en-Josas*

(2) *Institut National de la Recherche Agronomique, Station de Génétique quantitative et appliquée, 78350 Jouy-en-Josas*

(3) *Institut Technique du Porc, Région Ouest, B.P., 35650 Le Rheu*

(4) *Institut National de la Recherche Agronomique, Laboratoire de Génétique Biochimique, 78350 Jouy-en-Josas*

(5) *Centre d'Etudes Atomiques, I.P.S.N.-D.P.S.-S.P.E.-L.R.A., 91191 Gif-sur-Yvette Cedex*

A total of 243 female pigs were tested for 9 different genetic polymorphisms, and the results were related to production traits. The genetic systems studied were the blood groups A, E and H, the proteins Phi, Po2, Pgd, Pil and Tf, and the class I lymphocyte antigens of the swine major histocompatibility complex (SLA). The production traits were growth rate, carcass quality (muscle/fat ratio) and meat quality (as measured by pH, water holding capacity and reflectance). The animals were from the Large-White (LW) and French Landrace (LF) breeds, which differed in allelic frequencies at most loci. Among the genetic systems studied, only SLA showed a marked association with production traits. Although there was no association with growth rate, in either breed, certain SLA haplotypes were associated with meat quality (reflectance) in the LW. In both breeds, there was a close association between SLA and carcass leanness, although the same haplotype was surprisingly associated with a lower fat content in the LW and with a higher fat content in the LF. This finding indicates that caution is required in the interpretation of such studies, without a fuller understanding of the metabolic processes concerned.