



HAL
open science

Assessment of the contributions of glycolysis and the pentose phosphate pathway to glucose respiration in ectomycorrhizas and non-mycorrhizal roots of spruce (*Picea abies* L. Karsten)

Isabelle Bilger, V. Guillot, Francis Martin, François F. Le Tacon

► **To cite this version:**

Isabelle Bilger, V. Guillot, Francis Martin, François F. Le Tacon. Assessment of the contributions of glycolysis and the pentose phosphate pathway to glucose respiration in ectomycorrhizas and non-mycorrhizal roots of spruce (*Picea abies* L. Karsten). *Annales des sciences forestières*, 1989, 46 (suppl.), pp.724s-727s. hal-02726568

HAL Id: hal-02726568

<https://hal.inrae.fr/hal-02726568>

Submitted on 2 Jun 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Assessment of the contributions of glycolysis and the pentose phosphate pathway to glucose respiration in ectomycorrhizas and non-mycorrhizal roots of spruce (*Picea abies* L. Karsten)

I. Bilger, V. Guillot, F. Martin and F. Le Tacon

Laboratoire de Microbiologie Forestière, Centre de Recherches Forestières de Nancy, Institut National de la Recherche Agronomique, Champenoux 54280 Seichamps, France

Introduction

The importance of carbon supply in mycorrhizal infection and symbiotic activity has long been recognized. The supply of carbohydrates by the higher plant to the fungus is a very basic trait of mycorrhizal symbiosis. Mycorrhizal plants assimilate more photosynthates than non-mycorrhizal ones, allocate a greater fraction of the assimilated carbon to the root systems and lose a greater fraction of the assimilated carbon to respiratory CO₂ than do non-mycorrhizal plants (for a review, see Martin *et al.*, 1987). The establishment of a carbon sink by the ectomycorrhizal hyphae may be attained by: 1) rapid carbohydrate degradation for respiration and for energy and reducing power production and 2) conversion of plant carbohydrates into fungal biomass. The high respiration rate of fungal tissues has been pointed out by several authors (France and Reid, 1983). Most studies of mycorrhizal respiration deal with mitochondrial respiration. Much less is known about the oxidative metabolism of glucose in mycorrhizal

roots. The substrate used as well as the pathways potentially involved in this process are not known.

The aim of this study was to determine the relative contribution of glycolysis and the pentose phosphate pathway to glucose oxidation in Norway spruce (*Picea abies*) ectomycorrhizas.

Materials and Methods

Plant material

Four year old plants of *Picea abies* L. Karsten, grown on a sandy soil, were sampled from a commercial bare-roots nursery (Merten, Vosges, eastern France). The plants were removed with attached soil, stored at 4°C and transferred to the laboratory. The root systems were washed with tap water and all soil particles were removed. The pyramidally branched ectomycorrhizas were pale brown, racemose with a prosenchymatous sheath, a thin mantle and an extensive Hartig net reaching to the endodermis. There were abundant extramatricial mycelia (*Hebeloma* sp.) interconnected with loosely woven, pale yellow mycelial cords (see Fig. 1 in Al-Abras *et al.*, 1988; Dell *et al.*, 1989).

Assuming that the same chitin/protein ratio occurs in the mycelial cords of *Hebeloma* sp. and the fungal component of the ectomycorrhizas, then approximately 50% of the protein in the ectomycorrhizas is fungal (Dell *et al.*, 1989).

Radiorespirometry

A radiorespirometry study was performed using ectomycorrhizal subsamples and the non-mycorrhizal apices of exploratory roots. This was done using a 10 ml continuous $^{14}\text{CO}_2$ -evolving and -trapping reaction flask. About 50 mg of fresh tissue were incubated in 5 ml of distilled water containing 10 nmol of [$1\text{-}^{14}\text{C}$]glucose or 11 nmol of [$6\text{-}^{14}\text{C}$]glucose for 90 min at 22°C. Experiments were started by the addition of 0.5 μCi (10.0 nmol) of suitably labeled [^{14}C]glucose. An airflow of 200 $\text{ml}\cdot\text{min}^{-1}$ was maintained and $^{14}\text{CO}_2$ was collected for 90 min. Effluent air was passed directly into a CO_2 -trapping scintillation fluid containing an organic amine (Carbomax, Kontron) in 10 ml vials and counted using a scintillation counter (Betamatic I, Kontron). Residual radiolabel in the flask was determined by counting aliquots. Antibiotics were added to the incubation solution at the following concentrations to prevent bacterial activity: 0.02% (w/v) penicillin, 0.04% (w/v)

streptomycin and 0.008% (w/v) aureomycin. Soluble compounds were then extracted according to Al-Abras *et al.* (1988) and radioactivity determined by counting 100 μl aliquots. Chitin was determined by measuring the amount of fungal glucosamine resulting from acid hydrolysis of chitin in mycorrhizal roots and mycelial cords using the method of Vignon *et al.* (1986).

Statistical analysis

Data are presented as means of 4 or 6 replicates. Variance analysis or mean comparison was performed on the logarithm of the percentages or ratios.

Theoretical

The approach used is based on the assumption that the initial yield of $^{14}\text{CO}_2$ from [$1\text{-}^{14}\text{C}$]glucose represented glycolysis and the pentose phosphate pathway, whereas that from [$6\text{-}^{14}\text{C}$]glucose represented only glycolysis (Ap Rees, 1980). The following set of equations enables the contribution of the pentose phosphate pathway (PPP) to be calculated.

$$\text{respiration rate} = \frac{(\text{released } ^{14}\text{C as } \text{CO}_2 \times 100)}{(\text{released } ^{14}\text{C as } \text{CO}_2 + \text{incorporated } ^{14}\text{C in soluble compounds})} \quad (1)$$

$$\text{specific yield of } ^{14}\text{CO}_2 = \frac{(\text{released } ^{14}\text{C as } ^{14}\text{CO}_2 \times 100)}{(\text{released } ^{14}\text{C as } ^{14}\text{CO}_2 + \text{incorporated } ^{14}\text{C})} \quad (2)$$

$$R_1 = \frac{(\text{released } ^{14}\text{C as } ^{14}\text{CO}_2 \text{ from } [6\text{-}^{14}\text{C}]\text{glucose})}{(\text{released } ^{14}\text{C as } ^{14}\text{CO}_2 \text{ from } [1\text{-}^{14}\text{C}]\text{glucose})} \quad (3)$$

$$R_2 = \frac{(\text{specific yield of } ^{14}\text{C from } [6\text{-}^{14}\text{C}]\text{glucose})}{(\text{specific yield of } ^{14}\text{C from } [1\text{-}^{14}\text{C}]\text{glucose})} \quad (4)$$

PPP contribution =

$$\frac{[(\text{specific yield of } ^{14}\text{CO}_2 \text{ from } [1\text{-}^{14}\text{C}]\text{glucose}) - (\text{specific yield of } ^{14}\text{CO}_2 \text{ from } [6\text{-}^{14}\text{C}]\text{glucose})]}{(1 - \text{specific yield of } ^{14}\text{CO}_2 \text{ from } [6\text{-}^{14}\text{C}]\text{glucose})} \quad (5)$$

Table I. Respiration rate and C6/C1 ratios of *Hebeloma* sp. ectomycorrhizas and non-mycorrhizal exploratory roots fed [1-¹⁴C]glucose and [6-¹⁴C]glucose.

	Respiration rate from [1- ¹⁴ C]glucose	Respiration rate from [6- ¹⁴ C]glucose	C6/C1 ratios		PPP contribution
	(%)	(%)	R ₁	R ₂	(%)
Ectomycorrhizas	53	7	0.1	0.13	49.5
Non-mycorrhizal roots	40	17	0.3	0.43	38.6
LSD (<i>P</i> = 0.05)	+	+	+	+	+

For definition of respiration rate, R₁, R₂ and relative contribution (%) of the pentose phosphate pathway (PPP) to glucose respiration, see Materials and Methods.

Results and Discussion

The radiorespirometric method was applied to non-mycorrhizal exploratory roots and young mycorrhizas. Both non-mycorrhizal roots and ectomycorrhizas showed virtual simultaneous emission of ¹⁴CO₂ from [1-¹⁴C]- and [6-¹⁴C]glucose with similar patterns (data not shown). These data indicated the operation of more than one oxidative pathway. The rapid and predominant release of ¹⁴CO₂ from [1-¹⁴C]glucose coupled with low emission from [6-¹⁴C]glucose, in both samples, implied both a minor role of the tricarboxylic acid cycle and relatively low recycling of labeled glucose through the non-oxidative part of the pentose phos-

phate pathway and/or mannitol cycle (Martin *et al.*, 1985).

Using an incubation period of 90 min in labeled glucose, the C6/C1 ratios, R₁ and R₂ (Table I), were found to range from 0.10 to 0.13 for mycorrhizal roots and 0.30 to 0.43 for non-mycorrhizal ones. The low C6/C1 ratios of the mycorrhizal roots suggests a high activity of the pentose phosphate pathway. The level of CO₂ released from [6-¹⁴C]glucose was always comparatively lower. In non-mycorrhizal exploratory roots, 38% of the carbohydrate oxidation was *via* the pentose phosphate pathway and 62% was *via* glycolysis. On the other hand, 50% of the glucose metabolism from mycorrhizal roots was catalyzed by the pentose phosphate pathway,

Table II. Respiration rate and C6/C1 ratios of various tissues from *Hebeloma* sp. ectomycorrhizas and non-mycorrhizal exploratory roots fed [1⁴C]glucose.

	Respiration rate from [1- ¹⁴ C]glucose	Respiration rate from [6- ¹⁴ C]glucose	C6/C1 ratios		PPP contribution
	(%)	(%)	R ₁	R ₂	(%)
Extramatrixal hyphae	50.8	10.1	0.21	0.19	45.3
Fungal sheath and root cortex including the Hartig net	51.5	7.9	0.13	0.14	48.4
Root stele	49.2	16.4	0.29	0.30	39.4
LSD (<i>P</i> = 0.05)	ns	+	+	+	+

For definition of respiration rate, R₁, R₂ and relative contribution (%) of the pentose phosphate pathway (PPP) to

demonstrating that the carbohydrate oxidative pathways are drastically altered in response to fungal colonization of the root.

To determine the distribution of the two catabolic pathways, mycorrhizal roots were further separated into extramatrical hyphae, symbiotic root tissues (mantle, Hartig net hyphae plus root cortex) and stele. The contribution of the pentose phosphate pathway was different in the various mycorrhizal tissues, being higher in symbiotic tissues (49.2%) and extramatrical hyphae (46.5%) (Table II). The contribution of the pentose phosphate pathway in the stele of mycorrhizal roots was identical to that of whole non-mycorrhizal roots and accounted for 40%.

These differences between mycorrhizas and non-mycorrhizal roots and between fungal and host tissues suggest that the contribution of the pentose phosphate pathway to respiration is higher in the fungal component than in the plant tissues. The fact that the pentose phosphate pathway activity was even higher in root tissues colonized by the fungal cells (mantle and Hartig net) than in extramatrical hyphae suggests that the contribution of this oxidative pathway is stimulated when the root is associated with a symbiotic fungus. This increase in the pentose phosphate pathway activity may be related to the higher metabolic activity of the Hartig net revealed by ultrastructural studies of the host-fungus interface (many mitochondria and ribosomes, extensive development of the endoplasmic reticulum, lack of large vacuoles) (Kottke and Oberwinkler, 1986).

Whether there is an increase in the activity of the pentose phosphate pathway

enzymes or changes in the respective polypeptide amounts during ectomycorrhiza formation must await further analysis.

References

- Al-Abras K., Bilger I., Martin F., Le Tacon F. & Lapeyrie F. (1988) Morphological and physiological changes in ectomycorrhizas of spruce [*Picea excelsa* (Lam.) Link] associated with ageing. *New Phytol.* 110, 535-540
- Ap Rees T. (1980) Assessment of the contributions of metabolic pathways to plant respiration. In: *The Biochemistry of Plants, A Comprehensive Treatise. Vol. 2 Metabolism and Respiration.* (Stumpf P.K. & Conn E.E., eds.), Academic Press, London, pp. 1-27
- Dell B., Botton B., Martin F. & Le Tacon F. (1989) Glutamate dehydrogenases in ectomycorrhizas of spruce [*Picea excelsa* (Lam.) Link] and beech (*Fagus sylvatica* L.). *New Phytol.* 111, 683-692
- France R.C. & Reid C.P.P. (1983) Interactions of nitrogen and carbon in the physiology of ectomycorrhizae. *Can. J. Bot.* 61, 964-984
- Kottke I. & Oberwinkler F. (1986) The cellular structure of the Hartig net: coenocytic and transfer cell-like organization. *Nord. J. Bot.* 7, 85-95
- Martin F., Canet D. & Marchal J.P. (1985) ¹³C nuclear magnetic resonance study of mannitol cycle and trehalose synthesis during glucose utilization by the ectomycorrhizal ascomycete *Cenococcum graniforme*. *Plant Physiol.* 77, 499-502
- Martin F., Ramstedt M. & Söderhäll K. (1987) Carbon and nitrogen metabolism in ectomycorrhizal fungi and ectomycorrhizas. *Biochimie* 69, 569-581
- Vignon C., Plassard C., Mousain D. & Salsac L. (1986) Assay of fungal chitin and estimation of mycorrhizal infection. *Physiol. Vég.* 24, 201-207