



HAL
open science

Targeted proteomics method comparison: SRM, PRM and SWATH-MS to quantify proteins in bovine muscle tissues

Joanna Bons, Husson Gauthier, François Delalande, Sarah Cianférani, Brigitte Picard, Muriel Bonnet, Christine Carapito

► To cite this version:

Joanna Bons, Husson Gauthier, François Delalande, Sarah Cianférani, Brigitte Picard, et al.. Targeted proteomics method comparison: SRM, PRM and SWATH-MS to quantify proteins in bovine muscle tissues. 12. European Proteomics Association (EuPA), Jun 2018, Santiago de Compostela, Spain. hal-02733603

HAL Id: hal-02733603

<https://hal.inrae.fr/hal-02733603>

Submitted on 2 Jun 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Targeted proteomics method comparison: SRM, PRM and SWATH-MS to quantify proteins in bovine muscle tissues

Joanna Bons¹, Gauthier Husson², François Delalande¹, Sarah Cianférani¹, Brigitte Picard³, Muriel Bonnet³,
Christine Carapito¹

¹Laboratoire de Spectrométrie de Masse Bio-Organique (LSMBO), IPHC, UMR 7178, Université de Strasbourg, CNRS, 25 rue
Becquerel, 67087 Strasbourg, France

²Current address: Ablynx, Technologiepark 21, 9052 Zwijnaarde, Belgium

³INRA, Clermont Université, VetAgro Sup, UMR1213 Herbivores, BP 10448, 63000, Clermont-Ferrand, France

Background: Targeted proteomics has become an approach of choice to validate and precisely/absolutely quantify protein biomarkers. Besides selected reaction monitoring (SRM), alternative methods have emerged over the last few years, among which parallel reaction monitoring (PRM) and sequential windowed acquisition of all theoretical spectral (SWATH-MS). These approaches perform differently in terms of instrumentation, multiplexing, and performances such as sensitivity, selectivity and accuracy (Borràs and Sabidó, 2017). A careful consideration of these points is required to select the best-suited targeted proteomics method.

In this study, we have evaluated and benchmarked three targeted methods to precisely quantify beef tenderness biomarkers in muscle tissues. Meat tenderness and lipid content (marbling) guide customer choice. Controlling these quality criteria by relying on protein biomarkers presents a high agronomical and economical interest (Picard *et al.*, 2015; Ceciliani *et al.*, 2017).

Methods: Twenty peptides, corresponding to ten previously identified protein biomarkers of beef adiposity and tenderness, were quantified in two muscles, *Longissimus thoracis* and *Rectus abdominis*, using a SRM assay on a triple quadrupole mass spectrometer (MS) (TSQ Vantage, Thermo Fisher Scientific), a PRM assay on a quadrupole-Orbitrap MS (Q Exactive Plus, Thermo Fisher Scientific) and a SWATH-MS method on a quadrupole-time-of-flight MS (TripleTOF 6600, Sciex). Analyses were performed on a cohort of 64 samples of bovine muscles, covering a wide dynamic range of adiposity and tenderness. Limits of quantification for each peptide were determined by establishing isotope-dilution calibration curves.

Results and Conclusions: We showed that (i) PRM performs better than SRM and SWATH-MS regarding sensitivity, (ii) SRM and SWATH-MS exhibit similar – and enhanced while comparing to PRM – dynamic range performance, and (iii) PRM and SRM selectivity is better than SWATH-MS one for the low-concentrated peptides. We also highlighted the advantage of PRM and DIA-SWATH over SRM for post-acquisition data refinement.

Keywords: Quantitative Proteomics, Selected Reaction Monitoring, Parallel Reaction Monitoring, Data-Independent Analysis.