

Characterisation of the biosynthesis of saponins during seed development in peas (Pisum sativum) and faba beans (Vicia faba)

Julie Marais, Ludivine Lebeigle, Vanessa Vernoud, Christine Le Signor, Marie-Aleth Lacaille-Dubois, Richard Thompson

▶ To cite this version:

Julie Marais, Ludivine Lebeigle, Vanessa Vernoud, Christine Le Signor, Marie-Aleth Lacaille-Dubois, et al.. Characterisation of the biosynthesis of saponins during seed development in peas (Pisum sativum) and faba beans (Vicia faba). International Conference "Advances in Grain Legume Cultivation and Use. Translating Legume Research into end-Users Reality", Sep 2017, Novi Sad, Serbia. 180 p. hal-02733774

HAL Id: hal-02733774 https://hal.inrae.fr/hal-02733774

Submitted on 2 Jun 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

International Conference

Advances in grain legume cultivation and use

BOOK OF ABSTRACS





Translating Legume Research Into End-Users Reality

27-28 SEPTEMBER 2017 NOVI SAD, SERBIA

International Conference

Advances in grain legume breeding, cultivation and uses for a more competitive value-chain

BOOK OF ABSTRACTS

27-28 SEPTEMBER 2017 NOVI SAD, SERBIA

Characterization of the biosynthesis of saponins during seed development in peas (*Pisum sativum*) and faba beans (*Vicia faba*)

J. Marais⁽¹⁾, L. Lebeigle⁽¹⁾, V. Vernoud⁽¹⁾, C. Le Signor⁽¹⁾, M-A. Lacaille-Dubois⁽²⁾, R. Thompson⁽¹⁾

(1)INRA , UMR 1347 Agroecology. rue Sully,, 17, F21000, DIJON, FRANCE. (2)Université Bourgogne-Franche Comté, Pharmacognosie. Boulevard Jeanne d'Arc, 7, F-21079, DIJON, FRANCE. richard.thompson@inra.fr

The use of pulses as ingredients for the production of food products rich in plant proteins is increasing. However, protein fractions prepared from pea or faba beans contain significant amounts of saponins, glycosylated triterpenes which can impart a bitter taste to the final food product. In addition, saponins have also been described to be involved in plant responses to biotic and abiotic stresses(1). In this study, we identified and characterized the genes involved in saponin biosynthesis during pea seed development(2), and optimized a saponin extraction protocol to follow the biosynthesis of these compounds during the development of pea and faba bean seeds. The identification of mutants affecting the function of key genes of the saponin biosynthetic pathway is currently underway in pea(3). Acknowledgement: This study is funded under the LEG'UP FUI (Unique Interministerial Fund) project (AAP No. 18) References [1] Faizal, A. and D. Geelen (2013). "Saponins and their role in biological processes in plants." Phytochemistry Reviews 12(4): 877-893. [2] Morita, M., M. Shibuya, et al. (2000). "Molecular cloning and functional expression of triterpene synthases from pea (Pisum sativum) - New alpha-amyrin-producing enzyme is a multifunctional triterpene synthase." European Journal of Biochemistry 267(12): 3453-3460. [3] Dalmais, M., J. Schmidt, et al. (2008). "UTILLdb, a Pisum sativum in silico forward and reverse genetics tool." Genome Biology 9(2).