



HAL
open science

Characterization of the biosynthesis of saponins during seed development in peas (*Pisum sativum*) and faba beans (*Vicia faba*)

Richard Thompson, Ludivine Lebeigle, Julie Marais, Vanessa Vernoud, Christine Le Signor, Marie-Aleth Lacaille-Dubois

► To cite this version:

Richard Thompson, Ludivine Lebeigle, Julie Marais, Vanessa Vernoud, Christine Le Signor, et al.. Characterization of the biosynthesis of saponins during seed development in peas (*Pisum sativum*) and faba beans (*Vicia faba*). 8. International Conference on Legume Genetics and Genomics (ICLGG), Noble Research Institute., Sep 2017, Siófok, Hungary. 128 p. hal-02733806

HAL Id: hal-02733806

<https://hal.inrae.fr/hal-02733806v1>

Submitted on 2 Jun 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

BOOK OF ABSTRACTS



8th International Conference on Legume Genetics and Genomics
18-22 September 2017
Siófok, Hungary

Characterization of the biosynthesis of saponins during seed development in peas (*Pisum sativum*) and faba beans (*Vicia faba*)

R. Thompson¹, L. Lebeigle¹, J. Marais¹, V. Vernoud¹, C. Le Signor¹,
M.-A. Lacaille-Dubois²

¹ INRA UMR1347 Agroécologie Dijon, France

² 2Dept. Medicine, Univ. Burgundy Franche-Comté, France

The use of pulses as ingredients for the production of food products rich in plant proteins is increasing. However, protein fractions prepared from pea or faba beans contain significant amounts of saponins, glycosylated triterpenes which can impart a bitter taste to the final food product. In addition, saponins have also been described to be involved in plant responses to biotic and abiotic stresses¹. In this study, we identified and characterized the genes involved in saponin biosynthesis during pea seed development², and optimized a saponin extraction protocol to follow the biosynthesis of these compounds during the development of pea and faba bean seeds. The identification of mutants affecting the function of key genes of the saponin biosynthetic pathway is currently underway in pea³. This study is funded under the LEG'UP FUI (Unique Interministerial Fund) project (AAP No. 18).

Keywords: *protein, flavour, food*

References

- [1] Faizal, A. and D. Geelen (2013). "Saponins and their role in biological processes in plants." *Phytochemistry Reviews* 12(4): 877-893.
- [2] Morita, M, et al. (2000). "Molecular cloning and functional expression of triterpene synthases from pea (*Pisum sativum*) - New alpha-amyrin-producing enzyme is a multifunctional triterpene synthase." *European Journal of Biochemistry* 267(12): 3453-3460.
- [3] Dalmais, M, et al. (2008). "UTILLdb, a *Pisum sativum* in silico forward and reverse genetics tool." *Genome Biology* 9(2).

How to refer your abstract:

R. Thompson, L. Lebeigle, J. Marais, V. Vernoud, C. Le Signor, M.-A. Lacaille-Dubois (2017) Characterization of the biosynthesis of saponins during seed development in peas (*Pisum sativum*) and faba beans (*Vicia faba*); ICLGG 2017 - Book of abstracts, ICLGG2017/P/64