



HAL
open science

Structural study of lipid droplet using synchrotron label-free multimodal imaging

Frederic Jamme, Isabelle Bouchez, Caroline Pénicaud, Stéphanie Passot,
Fernanda Fonseca, Yann Gohon, Mathieu Réfrégiers, Eva Pereiro, Bertrand
Cinquin, Marine Froissard

► To cite this version:

Frederic Jamme, Isabelle Bouchez, Caroline Pénicaud, Stéphanie Passot, Fernanda Fonseca, et al.. Structural study of lipid droplet using synchrotron label-free multimodal imaging. 4. International Conference on Physics and Biological Systems, Oct 2018, Gif-sur-Yvette, France. , 2018, 4th International Conference on Physics and Biological Systems 2018. hal-02733903

HAL Id: hal-02733903

<https://hal.inrae.fr/hal-02733903>

Submitted on 2 Jun 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.


F. Jamme¹, I. Bouchez², C. Pénicaud³, S. Passot³, F. Fonseca³, Y. Gohon², M. Réfrégiers¹, E. Pereiro⁴, B. Cinquin⁵, M. Froissard²

¹Soleil Synchrotron, France; ²IJPB, INRA AgroParisTech, CNRS, Université Paris Saclay, France
³GMPA, INRA AgroParisTech, Université Paris Saclay, France; ⁴ALBA synchrotron, Spain; ⁵LBPA, CNRS, ENS Paris Saclay, France

LIPID DROPLETS

Lipid droplet: a complex and dynamic organelle

In cells, neutral lipids (triglycerides TG and steryl esters SE) are stored in organelles called **lipid droplets (LD)** [1]. They are present in all organisms, from bacteria to plants and animals.



Lipid droplets: not well known but with rising interest

From biologists

- LD is not an inert fat depot but a **dynamic organelle** which regulates cell metabolism and signaling

From medical field

- LDs have a crucial role in diseases with **increasing prevalence** (obesity, diabetes) [2]
- Oleosins** (from peanut and hazelnut), seed LD associated proteins are **allergens** [3].


From industrials

- crushing**: oils for food and non food industries are extracted from seed LDs
- food processing industry, cosmetic and health**: oleosins harbor interfacial properties and could be used as emulsifying agents or in drug delivery systems [4]



Lipid droplet structure under debate

Many fundamental questions about LD biology remain unanswered [5]. How LDs form and grow? How proteins move to and from LDs? How LDs are related to protein degradation? How LD core is structured?





IMAGING LIPID DROPLETS IS NOT A TRIVIAL ISSUE

LD subpopulations in yeast


Various LD populations revealed using epifluorescence microscopy with tagged protein

- with long chain fatty acyl-CoA synthetase Faa4p
- with more or less (white arrows) sterol- Δ 24-methyltransferase Erg6p




Various LD populations revealed using TEM

- Nucleus-associated (blue arrows)
- ER-associated (yellow arrows)
- Vacuole-associated (purple arrows)



Advantages and disadvantages of some LD imaging techniques

APPROACH	ADVANTAGES	DISADVANTAGES
Lipid probes	Living cells, fast and easy protocol, dynamic studies	Modified LDs and lipid environment
Proteins with fluorescent tag	Living cells, dynamic studies	Modified LDs, GMOs, non exhaustive labeling
AFM-IR	Label-free, lipid quantification	Dried cells
Electron microscopy	High resolution, visualisation of subcellular structures	Fixed cells, fastidious protocol
Soft X-ray microscopy	30 nm resolution, Label-free	Cryofixed cells, soft X-ray source and microscope
Differential Interference Contrast (DIC)	Living cells, visualisation of subcellular structures	structure identification
Raman (CARS, SRS)	Label-free, living cells, dynamic studies	Resolution, structure identification
Deep UV imaging	Label-free, living cells, 110 nm resolution, dynamic studies	DUV source and microscope



LIPID DROPLET LABEL-FREE IMAGING

Multimodal DUV imaging on living cells at Soleil Synchrotron


- DISCO beamline is dedicated to biochemistry, chemistry and cell biology.
- 3 experimental endstations, SRCD, APEX, DUV imaging [6].




Images were recorded using

- TELEMOS, a full field, fully automatized inverted microscope, for fast imaging of live cells
- sei1Δ mutant cells with supersized LDs (1μm vs 250 nm for WT), containing TG and ES such as WT cells [7]


STEP 1



STEP 2



STEP 3




- Step 1**: Identification of LDs (white arrow) using DPH, a vital neutral lipid probe with an excitation spectrum compatible with DUV (340 nm). LDs are low autofluorescent organelles.
- Step 2**: combination of autofluorescence and transmittance without lipid probe under 275 nm excitation. LDs (white arrows) show heterogeneous transmittance properties from surface to central core
- Step 3**: LD structural analysis (slices and plot profile) revealed a concentric layer organization, with a low 1/TRANS central core and a high 1/TRANS (absorbance) external ring, which could correspond to ergosterol species (absorbance main peaks at 270 and 280 nm)

Soft X-ray imaging at Alba Synchrotron

- MISTRAL beamline is devoted to cryo nano-tomography in the water window for biological applications [8].

Images were recorded using

- Transmission X-ray Microscope (TXM), 270 eV to 1200 eV, cryo-temperature, rotation from -65° to +65°
- Mutant cells with LD contrasted content (TG triglycerides, SE steryl esters, TG+SE) [9]




- LDs appear as black round structures corresponding to high absorbance structures.
- TG+SE containing LDs and TG containing LDs are homogeneous (pink arrows)
- SE containing LDs appear as "donuts" with high absorbance material (SE) at the periphery and a central "hole"

CONCLUSION ON LD STRUCTURE

DUV fluorescence and absorbance on living cells + Soft X-ray tomography

LDs with contrasted signals
LD chemical heterogeneity



Our results are in favor of the concentric layer model