

Integrating blood transcriptome and immunity traits to identify markers of immune capacity in pigs

Tatiana Maroilley, Fany Blanc, Gaetan Lemonnier, Jerôme Lecardonnel, Jean Pierre Bidanel, Andrea Rau, Yvon Billon, Marie-José Mercat, Jordi Estellé, Claire Rogel Gaillard

▶ To cite this version:

Tatiana Maroilley, Fany Blanc, Gaetan Lemonnier, Jerôme Lecardonnel, Jean Pierre Bidanel, et al.. Integrating blood transcriptome and immunity traits to identify markers of immune capacity in pigs. 69. Annual Meeting of the European Federation of Animal Science (EAAP), European Federation of Animal Science (EAAP). INT., Aug 2018, Dubrovnik, Croatia. hal-02733943

HAL Id: hal-02733943 https://hal.inrae.fr/hal-02733943

Submitted on 2 Jun2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Integrating blood transcriptome and immunity traits to identify markers of immune capacity in pigs

Maroilley T¹, Blanc F¹, Lemonnier G¹, Lecardonnel J¹, Bidanel JP¹, Rau A¹, Billon Y², Mercat MJ³, Estellé J¹, Rogel-Gaillard C¹

¹ GABI, INRA, AgroParisTech, Université Paris-Saclay, Jouy-en-Josas, France

² GENESI, INRA, Le Magneraud, Surgères, France

³ IFIP - Alliance R&D, La Motte au Vicomte, BP 35651, Le Rheu, France

Understanding individual variability of immune capacity in livestock has become a priority to improve sustainability, with the aim to increase disease resistance and resilience in breeding programs. In this study, 550 60-day-old French Large White pigs vaccinated against Mycoplasma hyopneumoniae (M hyo) were monitored for 55 immunity traits (ITs) measured from blood samples (SUS_FLORA ANR funded project). The ITs included two types of parameters. First, parameters directly measured from blood: complete blood counts, counts of various cell subsets by flow cytometry, serum dosage of anti M hyo IgG and haptoblobin. Second, parameters measured after in vitro stimulation of total blood: phagocytosis, production of cytokines (IL-1β, IL-8, IL-10, IL-17, TNF α , IFN γ) after stimulation by LPS or mitogenic agents. All animals were genotyped with 60K Illumina SNP chips. A subset of 243 piglets was chosen for blood transcriptome analysis using Agilent microarrays. We explored covariations between blood expression profiles and IT levels, and could draw lists of the most correlated genes with each IT. Each list represented candidate blood biomarkers potentially predictive of IT variations. As an example, we found 134 genes associated with phagocytosis capacity and we identified a subset of genes that could significantly predict levels of eight ITs related to phagocytosis by a sPLS approach. This gene subset included CXCR1, CCR1 and TLR2. Few candidate biomarkers were previously shown to be genetically controlled for their transcription in blood by eGWAS. Thus, our results provide new data to decipher the genetic architecture of IT variations. A next step will be to understand how IT variations could reflect individual robustness while facing pathogens, and how blood biomarkers could be used as predictors of immune capacity.