



**HAL**  
open science

## Importance of sugar homeostasis within xylem parenchyma cells to sustain xylem development

Emilie Aubry, Nelly Wolff, Sylvie Dinant, Catherine Bellini, Rozenn Le Hir

### ► To cite this version:

Emilie Aubry, Nelly Wolff, Sylvie Dinant, Catherine Bellini, Rozenn Le Hir. Importance of sugar homeostasis within xylem parenchyma cells to sustain xylem development. 5. International Conference of Plant Vascular Biology, Jun 2019, Asilomar, Californie, United States. hal-02734023

**HAL Id: hal-02734023**

**<https://hal.inrae.fr/hal-02734023v1>**

Submitted on 2 Jun 2020

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Generously sponsoring graduate student, postdoc and early career scientists travel awards.



Thank you to our sponsors.



## **Local Organizing Committee**

Siobhan Brady, University of California, Davis, [sbrady@ucdavis.edu](mailto:sbrady@ucdavis.edu)

Andrew Groover, US Forest Service, Davis, [agroover@fs.fed.us](mailto:agroover@fs.fed.us)

Mechthild Tegeder, Washington State University, Pullman, [tegeder@wsu.edu](mailto:tegeder@wsu.edu)

## **International Advisory Board**

Christine Beveridge, University of Queensland, Australia

Annelie Carlsbecker, University of Uppsala, Sweden

Taku Demura, NAIST and RIKEN, Japan

Sylvie Dinant, INRA Centre de Versailles-Grignon, France

Thomas Greb, University of Heidelberg, Germany

Rainer Hedrich, University of Würzburg, Germany

Michele Holbrook, Harvard University, USA

Ji-Young Lee, Seoul National University, Korea

Mengzhu Lu, Chinese Academy of Forestry, China

William J Lucas, University California, Davis, USA

Yi-Fang Tsay, Academia Sinica, Taiwan

peptide signaling, which is crucial to permit non-cell autonomous action of a phloem-derived signal. The data suggest that via this signal, phloem acts as an organizer tissue in plant organ formation.

Key Words: Phloem, BRI1, CLE45, brassinosteroids

## **Importance of sugar homeostasis within xylem parenchyma cells to sustain xylem development**

Emilie Aubry<sup>1</sup>, Nelly Wolff<sup>1</sup>, Sylvie Dinant<sup>1</sup>, Catherine Bellini<sup>1,2</sup> and Rozenn Le Hir<sup>1</sup>

<sup>1</sup> Institut Jean-Pierre Bourgin INRA-AgroParisTech, CNRS, Université Paris-Saclay, 78000 Versailles, France. <sup>2</sup> Umeå Plant Science Centre, Department of Plant Physiology, Umeå University, 90736 Umeå, Sweden

The Arabidopsis floral stem xylem tissue is formed by xylary parenchyma cells and dead cells including xylary fibres and xylary vessels. Especially, these last two cells functions request the presence of an extra-thickened secondary cell wall whose synthesis requires an important amount of carbohydrate skeletons. However, the pathways by which carbon pools, transported as sugars, are supplied to the xylem cells remain unclear. Recently, the SWEET11 and SWEET12 genes coding for sugar transporters located at the plasma membrane of the vascular parenchyma cells were proposed to act in this mechanism (1). To gain more insights in this process, we analyzed the function of two additional SWEET genes coding for tonoplastic sugar transporters also expressed in the xylem parenchyma cells (2,3). We analysed combinations of sweet mutants using microscopy combined with high-throughput image analysis and vibrational spectroscopy techniques. Our analysis reveals defects in the number and the size of the xylary fibres in the *sweet16sweet17* mutant, in contrast to the *sweet11sweet12* mutant; that has been previously shown to be mostly affected in the xylary vessels (1). Interestingly, the quadruple mutant shows defects in both xylary fibres and vessels, suggesting an additive phenotype. Due to differences in the expression pattern of these genes in different categories of bundles connected or not to axillary buds, we also scrutinized the effects of these mutations depending on the vascular bundle type. Our results suggest that the sugar homeostasis within the xylary parenchyma cells is required to ensure a correct xylem development.

(1) Le Hir et al., 2015. *Molecular Plant*, 8, 1687-1690 (2) Chardon et al., 2013. *Current Biology*. 23, 8, 697-702 (3) Klemens et al., 2013. *Plant Physiology*. 163, 1338-1352

## **Long- and Short Distance Plant Electrical Communication**

Rainer Hedrich<sup>1</sup> and Khaled A.S. AL-Rasheid<sup>1,2</sup>

<sup>1</sup> Molecular Plant Physiology & Biophysics, Wuerzburg University, Germany. [hedrich@botanik.uni-wuerzburg.de](mailto:hedrich@botanik.uni-wuerzburg.de). <sup>2</sup> College of Science, King Saud University, PO Box 2455, Riyadh 11451, Saudi Arabia

Electrical signals represent a process that evolved very early in the history of life. Travel over long distances such signals provide for an efficient way of achieving cell-to-cell communication in living organisms. In plants, the phloem can be considered as a ‘green cable’ that allows the transmission of action potentials (APs) induced by stimuli such as touch, wounding and cold.

Charles Darwin over 100 years ago recognized that the Venus flytrap *Dionaea muscipula* is capturing animals. When an insect visits the *Dionaea* trap and tilts the mechanosensors on the inner surface, action potentials (APs) are fired. After a moving object elicits two APs, the trap snaps shut, engaging the victim and initiating the hunting cycle. Our studies show that within the different steps of the hunting cycle,