



HAL
open science

Measurements of VOC fluxes by dynamic plant and soil chambers in wheat and maize crop near Paris with a PTR-Qi-TOF-MS: Quantification and response to environmental and physiological drivers

Lais Gonzaga-Gomez, Christophe Boissard, Raluca Ciuraru, Florence Lafouge, Olivier Zurfluh, Pauline Buysse, Celine Decuq, Olivier Fanucci, Jean Christophe J. C. Gueudet, Valérie Gros, et al.

► To cite this version:

Lais Gonzaga-Gomez, Christophe Boissard, Raluca Ciuraru, Florence Lafouge, Olivier Zurfluh, et al.. Measurements of VOC fluxes by dynamic plant and soil chambers in wheat and maize crop near Paris with a PTR-Qi-TOF-MS: Quantification and response to environmental and physiological drivers. EGU 2017, European Geophysical Union General Assembly 2017, Apr 2017, Vienne, Austria. , 1 p., 2017. hal-02734062

HAL Id: hal-02734062

<https://hal.inrae.fr/hal-02734062>

Submitted on 2 Jun 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - ShareAlike 4.0 International License

Measurements of VOC fluxes by dynamic plant and soil chambers in wheat and maize crop near Paris with a PTR-Qi-TOF-MS: Quantification and response to environmental and physiological drivers.

Lais Gonzaga-Gomez (1), Christophe Boissard (2), Raluca Ciuraru (1), Florence Lafouge (1), Olivier Zurfluh (1), Pauline Buisse (1), Céline Decuq (1), Olivier Fanucci (1), Jean-Christophe Gueudet (1), Valérie Gros (2), Roland Sarda (2), Nora Zannoni (2), and Benjamin Loubet (1)

(1) UMR ECOSYS, INRA-AgroParisTech, Université Paris Saclay, Thiverval-Grignon, France (lais.gonzaga-gomez@inra.fr),
(2) UMR CNRS-CEA, LSCE, Gif-sur-Yvette, France

Volatile organic compounds (VOC) play an important role in the chemistry of the atmosphere as precursors of secondary pollutants such as ozone and organic aerosols. A large variety of VOC are exchanged between plants (BVOC) and the atmosphere. Their fluxes are strongly dependent on environmental factors (temperature, light, biotic and abiotic stress) and vary greatly among plant species. Only few studies focused on BVOC emissions by agricultural plants and were mostly carried in North America. However, agricultural lands occupy 51% of the total country area in France, with wheat being one of the most important crop.

We used a PTR-Qi-TOF-MS (national instrument within the ANAEE-France framework) and dynamic chambers to measure BVOC emissions from plant and soil compartments of a wheat and a maize crop near Paris (FR-GRI ICOS site). More than 700 masses were detected thanks to the resolution and sensitivity of this new instrument. We analyze the emission response to light, temperature and stomatal aperture in order to explain the mechanisms of BVOC exchanges by wheat plants. We investigate the emission differences between soil and plant compartment, and between wheat and maize crops. Acetone (m/z 59.049) was the predominant volatile compound in the emissions from wheat. Both methanol (m/z 33.033) and acetaldehyde (m/z 45.033) were also quite abundantly emitted but were less than half the acetone emissions. Other masses detected in relative importance in this study were m/z 63.026 (possible DMS), m/z 93.033 (not identified), m/z 69.069 (isoprene), m/z 57.069 (not identified), m/z 83.085 (possible green leaf volatiles), m/z 73.064 (methyl ethyl ketone). Their emissions were around 7 times smaller than the emissions of acetone. On the other hand we observed a deposition for, mainly, m/z 75.044 (hydroxyacetone) and m/z 61.028 (acetic acid). Methanol presented both positive and negative fluxes which could indicate either emission or absorption of this compound by the plant, respectively.