



HAL
open science

Genetic variation of alfalfa seed yield and its components in the year of establishment

Dura Karagic, Dragan B. Milic, Snezana M. Katanski, Branko R. Milosevic, Miroslav Z. Zoric, Bernadette Julier

► To cite this version:

Dura Karagic, Dragan B. Milic, Snezana M. Katanski, Branko R. Milosevic, Miroslav Z. Zoric, et al.. Genetic variation of alfalfa seed yield and its components in the year of establishment. 10. International Herbage Seed Group Conference (IHSG 2019), Oregon State University (OSU). Corvallis, USA., May 2019, Corvallis, Oregon, United States. hal-02734089

HAL Id: hal-02734089

<https://hal.inrae.fr/hal-02734089>

Submitted on 2 Jun 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Genetic variation of alfalfa seed yield and its components in the year of establishment

Đura Karagić^{1*}, Dragan B. Milić¹, Snežana M. Katanski¹, Branko R. Milošević¹, Miroslav Z. Zorić¹, Bernadette Julier²

¹Institute of Field and Vegetable Crops, Novi Sad, Serbia

²INRA - French National Institute for Agricultural Research, Lusignan, France

djura.karagic@ifvcns.ns.ac.rs

Abstract: In Serbia, alfalfa is grown on about 140,000 ha, out of which 3-7% are intended for seed production. Climatic factors (amount and distribution of precipitation) are the main determinant of alfalfa seed yield. Average alfalfa seed yield in Serbia is about 250 kg ha⁻¹, and seed production is characterized with huge variation (15-800 kg ha⁻¹). The most challenging is the seed production in the first year of plant life (establishment year). Even if breeding for seed yield is not highlighted, some genetic variation was evidenced. New alfalfa varieties are selected for high forage production, quality of forage and persistence to be successful on the market. However, an outstanding seed-yield potential is needed. In order to examine genetic variation for alfalfa seed yield and its components in the sowing year, a trial with 20 accessions was established at the experimental field of IFVCNS. These accessions, with dormancy ratings 3-6, are representative of the European commercial varieties: 11 from France, 4 from Serbia, 3 from Czech Republic, one from Sweden and one from Denmark. The experiment was established on 21 May 2018. The elementary plot size was 6 m². Seed yield (t ha⁻¹), and its components were recorded on each plot. Year of 2018 was not favorable for alfalfa seed production in Serbia. The results clearly demonstrate differences among varieties for total seed yield and its components in the year of establishment. Seed yields varied from 7.6 kg ha⁻¹ for the variety Luzelle up to 65 kg ha⁻¹ for the variety Etincelle. Higher seed yields were obtained with less dormant varieties (5-6), while lower yields were recorded with more dormant varieties (3-4). Overall analyses showed that even under unfavorable conditions, genetic variation could be important for alfalfa seed production in the year of establishment.

Keywords: alfalfa, seed yield components, *Medicago*, variety

Acknowledgments: This research has received funding from the European Union's Horizon 2020 project EUCLEG under grant agreement n°727312.