



HAL
open science

Towards the identification and quantification of candidate metabolites of tebuconazole fungicide

Najoi El Azhari, Eftichia Dermou, Lucio Boterri, Luigi Lucini, Karas Panagiotis, Riccardo Rossi, Dimitrios Karpouzas, Georgios Tsiamis, Fabrice Martin-Laurent, Marco Trevisan, et al.

► **To cite this version:**

Najoi El Azhari, Eftichia Dermou, Lucio Boterri, Luigi Lucini, Karas Panagiotis, et al.. Towards the identification and quantification of candidate metabolites of tebuconazole fungicide. 7. Conference on Pesticide Behaviour in Soils, Water and Air, Aug 2017, York, United Kingdom. <hal-02734291>

HAL Id: hal-02734291

<https://hal.inrae.fr/hal-02734291v1>

Submitted on 2 Jun 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



HAL Authorization

TOWARDS THE IDENTIFICATION AND QUANTIFICATION OF CANDIDATE METABOLITES OF TEBUCONAZOLE FUNGICIDE

Najoi El Azhari^{2,3}, Eftichia Dermou^{4,1}, Lucio Botteri^{1,2}, Luigi Lucini⁵, Panagiotis Karas^{6,1}, Riccardo Rossi¹, Dimitrios G. Karpouzas⁶, Georgios Tsiamis⁴, Fabrice Martin-Laurent², Marco Trevisan⁵, Federico Ferrari¹

¹*Aeiforia srl, Piacenza, Italy*, ²*INRA, Dijon, France*, ³*SATT GRAND EST, Dijon, France*, ⁴*University of Patras, Agrinio, Greece*, ⁵*Università Cattolica del Sacro Cuore, Piacenza, Italy*, ⁶*University of Thessaly, Larissa, Greece*

Tebuconazole belongs to the family of triazole fungicides, used for crop protection and human health applications. In the environment, the dissipation of the parent molecule leads to the formation of metabolites that are of unknown identity or toxicity.

In order to identify and determine the putative identity of those metabolites and their potential toxicity, a quadrupole time-of-flight (Q-TOF) approach is often used. Q-SAR approaches help to predict their toxicity by comparing them to a known database of molecules with known properties. All together the information on the candidate by-products may help to select relevant sub-set of metabolites for further quantification by LC or GC coupled with MS. It is thereby possible to select putative toxic compounds for further quantification using chemical analysis.

Previous work allowed the identification of potential metabolites of tebuconazole. Triazole, triazolyl acetic acid and p-chlorophenol were suspected to result from the decomposition of tebuconazole. Tebuconazole degradation kinetics was followed for 125 days by quantifying the dissipation of the parent molecule and the emergence of the three candidate metabolites by LC/MS for tebuconazole, triazol and triazolyl acetate and by GC/MS for p-chlorophenol. The data allowed the proposition of several metabolic pathways.