



HAL
open science

Micro-evolution of sex determination mechanisms and sex determining genes in the cavefish *Astyanax mexicanus*

Boudjema Imarazene, Amaury Herpin, Romain Feron, John H Postlethwait, Manfred Schartl, Sylvie Rétaux, Yann Guiguen

► **To cite this version:**

Boudjema Imarazene, Amaury Herpin, Romain Feron, John H Postlethwait, Manfred Schartl, et al.. Micro-evolution of sex determination mechanisms and sex determining genes in the cavefish *Astyanax mexicanus*. 8. International Symposium on the Biology of Vertebrate Sex Determination, Apr 2018, Kona (Hawaii), United States. hal-02734624

HAL Id: hal-02734624

<https://hal.inrae.fr/hal-02734624>

Submitted on 2 Jun 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

**EIGHTH INTERNATIONAL SYMPOSIUM ON
THE BIOLOGY
OF VERTEBRATE SEX DETERMINATION**



April 16-20, 2018 KONA, HAWAII

EIGHTH INTERNATIONAL SYMPOSIUM ON THE BIOLOGY OF VERTEBRATE SEX DETERMINATION

16-20, APRIL 2018

King Kamehameha's Kona Beach Hotel

Kailua-Kona, Hawaii

Organizers

Richard Behringer
University of Texas MD Anderson Cancer Center
Houston, Texas

Blanche Capel
Duke University
Durham, North Carolina

Martin Cohn
Howard Hughes Medical Institute
University of Florida
Gainesville, Florida

Monika Ward
University of Hawaii at Manoa
Honolulu, Hawaii

Cover: Statue of Ardhanarishvara, the half-male, half-female form of Shiva, India 11th century A.D.

MICRO-EVOLUTION OF SEX DETERMINATION MECHANISMS AND SEX DETERMINING GENES IN THE CAVEFISH, *ASTYANAX MEXICANUS*.

Boudjema Imarazene¹, Amaury Herpin¹, Romain Feron¹, John H. Postlethwait²,
Manfred Scharl³, Sylvie Retaux⁴, Yann Guiguen¹

¹ INRA, Fish Physiology and Genomics Laboratory, 35042 Rennes, France.

² Institute of Neuroscience, University of Oregon, Eugene, USA.

³ Department of Physiological Chemistry, University of Wuerzburg, Wuerzburg, Germany.

⁴ CNRS, UPR3294, N&D laboratory, DECA group, 91198 Gif sur Yvette, France.

Vertebrates display remarkable diversity of sex determining (SD) mechanisms, ranging from environmental to genetic sex determination (ESD and GSD). In teleosts, sex determination (SD) mechanisms include both environmental and genetic regulation and are extremely diverse, changeable, and labile. In addition, many sex-determining (SD) genes have been already found in this group of vertebrates, showing that SD genes can be extremely variable even between closely related species. Hence, teleost fish have emerged as interesting models to study the macroevolution of SD mechanisms within vertebrates. The cavefish *Astyanax mexicanus* belongs to the Characiform group. Different populations of *Astyanax* have been described, including pigmented river-dwelling and several depigmented blind cave populations. These populations are still inter-fertile and cave populations are known to have evolved from a surface ancestral population less than 100,000 years ago. In addition these *Astyanax* populations can be also reared under laboratory conditions quite easily and are amenable to genetic manipulations. Altogether, this makes surface and cave populations a particularly interesting evolutionary genetic model system for comparative and microevolution studies. We thus initiated studies on the microevolution of SD and SD genes in the cavefish *Astyanax mexicanus*, and used restriction site-associated DNA (RAD) sequencing and pool-sequencing approaches to identify sex-biased molecular markers in both cavefish originating from the Pachón cave and surface fish populations. Our first results led to the identification of a candidate master SD gene that potentially triggers SD in both morphs. Despite this conservation of a strict GSD and the same master SD gene in both populations, our results also reveal a quick evolutionary transition between a polygenic SD system in surface population and a simple monofactorial system in the Pachón cave population.