

Model-based estimation of aphid dispersal distances and its use to co-optimize sharka management strategies and the allocation of resistant cultivars

Coralie Picard, David Pleydell, Loup Rimbaud, Victor Picheny, Sylvie Dallot, Emmanuel Jacquot, Samuel Soubeyrand, Gael Thébaud

▶ To cite this version:

Coralie Picard, David Pleydell, Loup Rimbaud, Victor Picheny, Sylvie Dallot, et al.. Model-based estimation of aphid dispersal distances and its use to co-optimize sharka management strategies and the allocation of resistant cultivars. 17. Rencontres de Virologie Végétale (RVV 2019), Jan 2019, Aussois, France. 127 p. hal-02734861

HAL Id: hal-02734861 https://hal.inrae.fr/hal-02734861

Submitted on 2 Jun 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



0.07

Model-based estimation of aphid dispersal distances and its use to co-optimize sharka management strategies and the allocation of resistant cultivars

Picard C.^{1,2}, Pleydell D.^{1,3}, Rimbaud L.^{1,4}, Picheny V.⁵, Dallot S.¹, Jacquot E.¹, Soubeyrand S.⁶, **Thébaud G.**¹

ANSES, UCAS, Lyon, France

Models can estimate epidemiological parameters, assess the effectiveness of different management strategies and optimize them. Here we present the PESO (parameter estimation-simulationoptimization) modelling framework (Picard et al., 2017), and we apply it to the management of sharka (caused by *Plum pox virus*), the most damaging Prunus disease. In a first step (Pleydell et al., 2018), using a spatially-explicit simulation model integrating disease dispersal, surveillance and control, we demonstrated that disease dispersal distances can be estimated accurately when disease control is ongoing. Applying this model to 15 years of sharka surveys in 605 orchards, we obtained the first estimate of aphid flight distances at the landscape scale. About 50% of aphid flights terminate beyond 90 m, and 10% exceed 1 km. By their impact on our quantitative understanding of winged aphid dispersal, these results can inform the design of management strategies for plant viruses, which are mainly aphid-borne. Indeed, we integrated these parameter estimates in an expanded version of the model, simulating sharka management options based on orchard surveillance, tree removal, plantation bans, and the replacement of at most half of the removed susceptible orchards with resistant cultivars. With this tool, we studied the spatial allocations of resistant orchards that minimize the direct and indirect economic impact of PPV epidemics in the landscape. We tested several allocations of resistant orchards for either emerging or established epidemics, on landscapes varying by their level of patch aggregation. This study (Picard, 2018) showed that the most promising allocations of durably resistant cultivars were (i) the regular mixing of susceptible and resistant orchards (in the absence of disease management), and (ii) any allocation of resistant orchards (under effective disease management). In addition, co-optimizing disease management and the spatial allocation of resistant cultivars slightly improved our economic criterion.

¹ INRA, UMR BGPI, Campus de Baillarguet, 34398, Montpellier Cedex 5, France

³ INRA, UMR ASTRE, Montpellier, France

⁴ INRA, UR Pathologie Végétale, Avignon, France


⁵ INRA, UR MIAT, Castanet-Tolosan, France

⁶ INRA, UR BioSP, Avignon, France

17emes Rencontres de Virologie Végétale

27 au 31 janvier 2019 Aussois - France

RVV 2019



Aussois du 27 au 31 janvier 2019