



# Comparative development of embryos and placenta in mammals: how to choose the right biomedical model

Pascale Chavatte-Palmer, Véronique Duranthon

## ► To cite this version:

Pascale Chavatte-Palmer, Véronique Duranthon. Comparative development of embryos and placenta in mammals: how to choose the right biomedical model. COST Action BM 1308 “ Sharing Advances on Large Animal Models - SAALAM ”, Ludwig Maximilians University of Munich (LMU). DEU. European Cooperation in Science and Technology (COST), BEL., Sep 2017, Halle, Germany. hal-02734928


**HAL Id: hal-02734928**

**<https://hal.inrae.fr/hal-02734928>**

Submitted on 2 Jun 2020

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.




Leopoldina  
Nationale Akademie  
der Wissenschaften

**COST Action BM 1308 "Sharing Advances on Large Animal Models – SALAAM"**

## **FINAL CONFERENCE**

**ADVANCES ON LARGE ANIMAL MODELS: BRIDGING THE GAP BETWEEN  
BIOMEDICAL RESEARCH AND CLINICAL TRANSLATION**

**September 28-29, 2017, Halle (Saale), DE**



## **PROGRAMME AND ABSTRACTS**

**Comparative development of embryos and placenta in mammals: how to choose the right biomedical model**

Pascale Chavatte-Palmer and Véronique Duranthon

UMR BDR, INRA, ENVA, Université Paris Saclay, 78350, Jouy en Josas, France

Most research on embryo development and subsequent pregnancy is performed in rodent models. There is considerable evidence, however, that the mice is not always appropriate to mimic human development. For example, the major embryonic genome activation takes place at the 2 cell stage (about 30-32 hours post-conception) in mice whereas it takes place later and after several cell cycles in humans and domestic species. Major differences in embryo metabolism and the apposition of epigenetic marks have also been demonstrated. In terms of placental development, the inverted yolk sack placenta of the mice, that is present in the first half of pregnancy, is unique to mammals. In humans, the placenta remains hypoxic in early development until the opening of the vascular plugs of the spiral arteries allow the onset of placental perfusion at the end of the first trimester. Perturbations of this phenomenon induce the human specific pregnancy diseases, pre-eclampsia and vascular induced intra-uterine growth retardation. Finally, the timing of organ development varies between species. Thus, the choice of an appropriate model for the study of healthy and sick pregnancy in humans remains a conundrum, but the use of alternate animal models such as rabbits, pigs or ruminants may provide useful alternatives to rodents.