

Does early growth play a role in the sex determination of european seabass dicentrarchus labrax?

Marc Vandeputte, Khaled Horri, François Allal, Sebastien Ferrari, Marie-Odile Vidal, François Ruelle, Marie-Laure Bégout, Béatrice Chatain

▶ To cite this version:

Marc Vandeputte, Khaled Horri, François Allal, Sebastien Ferrari, Marie-Odile Vidal, et al.. Does early growth play a role in the sex determination of european seabass dicentrarchus labrax?. 12th International Symposium on Genetics in Aquaculture (ISGA), Jun 2015, Santiago de Compostela, Spain. pp.1. hal-02735070

HAL Id: hal-02735070 https://hal.inrae.fr/hal-02735070

Submitted on 2 Jun 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

DOES EARLY GROWTH PLAY A ROLE IN THE SEX DETERMINATION OF EUROPEAN SEABASS Dicentrarchus labrax?

Vandeputte M.^{1,28}, Horri K.¹, Allal F.¹, Ferrari S.³, Vidal M.-O.¹, Ruelle F.¹, Bégout M.-L.³, Chatain B.¹

¹ Ifremer UMR9190 MARBEC, Chemin de Maguelone, 34250 Palavas-les-Flots, France ² INRA, UMR1313 Génétique animale et Biologie intégrative, 78350 Jouy-en-Josas, France ³ Ifremer, Ressources Halieutiques de La Rochelle, Place Gaby Coll, 17137 L'Houmeau, France

European sea bass has a polygenic sex determinism with temperature influences, which remains hard to predict and manipulate. To investigate the interactions between early growth and sex determination, we studied West Mediterranean domesticated sea bass from a 50&x10♀ full factorial cross, reared as a single batch at 15°C to promote female sex determination. Temperature was increased to 25°C from 70 to 119 days post-hatching (dph) to increase growth. Fish were individually tagged at 95 dph (575 mg) with microtags and were individually weighed at 95, 115, 136, 157, 200, 256 and 325 dph, and then sexed. Parentage was recovered with 12 microsatellites; 1134 fish had appropriate pedigree, sex and growth data. Multivariate sire models were run to investigate the links between growth and sex (considered a threshold trait with an underlying sex tendency). Heritability of sex tendency was 0.39±0.12 on the underlying scale. The sex dimorphism of body weight in favour of females rose from +27% at 95 dph (BW $_{05}$) to +40-45% between 115 and 325 dph, and the sex dimorphism of DGC was maximal for DGC_{os-115}. If differential growth is seen as a consequence of phenotypic sex, growth should be corrected by a fixed sex effect. In this case, there was a positive, although not significant (r_{*}=0.18±0.12) genetic correlation of sex tendency with DGC_{05,18}, while it never exceeded 0.05 later than 136 dph. The genetic correlation of BW with sex tendency regularly decreased from 0.45±0.18 for BW₉₅ to -0.20±0.17 for BW₃₂₅. If differential growth is seen as a cause of phenotypic sex, no fixed sex effect should be modelled, which then led to a r, of sex tendency with DGC decreasing from 0.60±0.11 for DGC_{05.115} to 0.21 ± 0.15 for DGC₂₅₆₋₃₃₅₇, while the r₄ of BW with sex tendency also decreased from 0.72 ± 0.10 for BW₁₁₅ to 0.48±0.14 for BW₃₂₅. In all cases, the genetic and environmental correlation of growth and sex tendency was maximal at the earlier stages, especially before the first reported signs of sex differentiation. This leads us to think that post-larval growth is, at least partially, a cause and not a consequence of sex determination in sea bass.

Keywords: Sex determination, quantitative genetics, growth

[§] Corresponding author. Tel.: + 33 4 67 13 04 07; Fax: + 33 4 67 13 04 58 E-mail address: + marc.vandeputte@jouy.inra.fr