



HAL
open science

Methane emissions in sheep infected with parasitic nematodes and supplemented with *Acacia mearnsii*

Paulo de Mello Tavares Lima, Pierre Crouzoulon, Geneviève Zabré, Adama Kaboré, Vincent Niderkorn, Herve Hoste, Adibe Luiz Abdalla, Helder Louvandin

► To cite this version:

Paulo de Mello Tavares Lima, Pierre Crouzoulon, Geneviève Zabré, Adama Kaboré, Vincent Niderkorn, et al.. Methane emissions in sheep infected with parasitic nematodes and supplemented with *Acacia mearnsii*. 7. GGAA Greenhouse Gases & Animal Agriculture Conference (GGAA), 2019, Iguassu Falls. hal-02735207

HAL Id: hal-02735207

<https://hal.inrae.fr/hal-02735207v1>

Submitted on 2 Jun 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/335149914>

Methane emissions in sheep infected with parasitic nematodes and supplemented with *Acacia mearnsii*

Conference Paper · August 2019

CITATIONS

0

READS

67

8 authors, including:



Paulo de Mello Tavares Lima

University of São Paulo

39 PUBLICATIONS 125 CITATIONS

[SEE PROFILE](#)



Adama Kaboré

116 PUBLICATIONS 382 CITATIONS

[SEE PROFILE](#)



Vincent Niderkorn

French National Institute for Agricultural Research

68 PUBLICATIONS 906 CITATIONS

[SEE PROFILE](#)



Hervé Hoste

French National Institute for Agricultural Research

243 PUBLICATIONS 7,213 CITATIONS

[SEE PROFILE](#)

Some of the authors of this publication are also working on these related projects:



Productive aspects and methane emission in Santa Inês sheep supplemented with the legume *Macrotyloma axillare* [View project](#)



index of quality of government institutions [View project](#)

DOCUMENTOS

135

ISSN 1980-6841
August/ 2019

7th GGAA – Greenhouse Gas and Animal Agriculture Conference
August 4th to 8th, Iguassu Falls/Brazil

Proceedings



Methane emissions in sheep infected with parasitic nematodes and supplemented with *Acacia mearnsii*

Paulo de Mello Tavares Lima¹, Pierre Crouzoulon², Geneviève Zabré³, Adama Kabore⁴, Vincent Niderkorn⁵, Hervé Hoste⁶, Adibe Luiz Abdalla⁷ and Helder Louvandini⁸

¹Universidade de São Paulo, Centro de Energia Nuclear na Agricultura, Piracicaba, São Paulo, Brasil – PhD - Post-Doctoral Researcher

²Universidade de São Paulo, Centro de Energia Nuclear na Agricultura, Piracicaba, São Paulo, Brasil – MSc Student

³Laboratoire de Biologie et Santé Animales-DPA/INERA, Ouagadougou, Burkina Faso MSc - PhD Student

⁴Laboratoire de Biologie et Santé Animales-DPA/INERA, Ouagadougou, Burkina Faso PhD - Professor

⁵Université Clermont Auvergne, INRA, VetAgro Sup, UMR Herbivores, F-63122 Saint-Genès Champanelle, France – PhD – Scientific Researcher

⁶INRA, UMR, Chemin des Capelles, Toulouse, France – PhD – Senior Researcher

^{7,8}Universidade de São Paulo, Centro de Energia Nuclear na Agricultura, Piracicaba, São Paulo, Brasil - PhD - Associate Professor

Our objective was to evaluate the effects of *Acacia mearnsii* bark extract (PAB) on methane (CH₄) production in sheep infected with *Trichostrongylus colubriformis* and *Haemonchus contortus*. Twenty 10 month-old Santa Inês lambs were used in a 50-day trial. Four treatment groups were formed: two control groups uninfected, one without PAB (C-) (n=4) and one with PAB (C+) (n=4); and two infected groups, one without PAB (I-) (n=6) and another receiving PAB (I+) (n= 6). Animals were kept in individual pens for 40 days, and received ad libitum chopped tifton 85 hay (*Cynodon* spp.) and 210 g/animal/day of concentrate (PAB supplementation: 15 g/animal/day – in concentrate). Animals were reared indoors to avoid helminthic infections (faecal egg count (FEC) was monthly controlled). At day 0, infected groups were artificially infected with L3 larvae of *T. colubriformis* (7500) and *H. contortus* (5000). After 28 days post-infection, establishment of infections was confirmed by FEC: I-: 1150 + 530.1 and I+: 567 + 544.6 eggs/gram of feces. After 40 days of trial (during which other parameters were evaluated - data not shown), CH₄ emissions in chambers were measured and variance analysis (proc GLM, SAS®) showed no PAB effects on methanogenesis (p > 0.05), however, both infected groups had higher CH₄ emissions than the control groups: C-: 16.7 + 10.91; C+: 23.7 + 12.88; I-: 43.1 + 6.37; I+: 40.6 + 6.02 (g CH₄/kg dry matter intake) (p < 0.05). The methanogenic effect of parasitic infections should be explored in studies addressing sustainability of animal production.

Keywords: Black wattle, *Haemonchus contortus*, Small chamber, *Trichostrongylus colubriformis*, Tannins