

Deciphering the genetic basis of tolerance to Cucumber mosaic virus in pepper

Judith Hirsch, Marion Szadkowski, Gregory Girardot, Allan Kampfer, Maxime Louis, Sandra Martin, Sabrina Azehaf, Yseult Cardona, Alexandra Schoeny, Pauline Millot, et al.

▶ To cite this version:

Judith Hirsch, Marion Szadkowski, Gregory Girardot, Allan Kampfer, Maxime Louis, et al.. Deciphering the genetic basis of tolerance to Cucumber mosaic virus in pepper. 17. Rencontres de Virologie Végétale (RVV 2019), Jan 2019, Aussois, France. 127 p., 2019, 17èmes Rencontres de Virologie Végétale: RVV 2019. hal-02735594

HAL Id: hal-02735594 https://hal.inrae.fr/hal-02735594

Submitted on 2 Jun 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



P.41

Deciphering the genetic basis of tolerance to Cucumber mosaic virus in pepper


Judith Hirsch¹, Marion Szadkowski¹, Grégory Girardot¹, Alan Kampfer¹, Maxime Louis¹, Sandra Martin¹, Sabrina Azehaf¹, Yseult Cardona¹, Alexandra Schoeny¹, Pauline Millot¹, Catherine Wipf-Scheibel¹, Patrick Gognalons¹, Cécile Desbiez¹, Karine Nozeran¹, Eric Verdin¹, Alain Palloix², Benoît Moury¹

Plant defense against parasites can be divided into two distinct components: mechanisms that reduce parasite accumulation (resistance sensu stricto) and mechanisms that reduce the negative impact of infection on host fitness, health or yield without impacting parasite concentrations (tolerance). As it is expected to exert a weaker selection pressure on parasite populations, plant tolerance to pathogens appears as an interesting alternative to resistance s.s. for sustainable disease management. However, little is known about the genetic determinants controlling tolerance to parasites. We have chosen the interaction between pepper and Cucumber mosaic virus (CMV) as a model to study plant tolerance to viral pathogens. We performed a screen using a pepper doubled haploid (DH) mapping population to map CMV tolerance and resistance QTLs. Both virus titer and plant health were simultaneously evaluated for each DH line. Virus accumulation was quantified using serological methods (DAS-ELISA, Double Antibody Sandwich Enzyme Linked Immunosorbent Assay). The impact of infection on plant health was measured using different methods. These methods included calculating the AUDPC (Area Under the Disease Progress Curve) index, which combines time of symptom emergence and symptom intensity, measuring plant growth parameters, such as the reduction in fresh weight of infected plants compared to mockinoculated plants, and measuring leaf chlorophyll content. Our screening efforts have shown that plant health and virus load are not correlated and allowed the identification of lines displaying contrasted levels of tolerance and resistance to CMV. Mapping efforts are underway to detect QTLs controlling tolerance or resistance s.s., and to compare the genetic architecture of these two defense mechanisms.

¹ INRA, Unité de Pathologie Végétale (UR407), CS 60094, 84143 Montfavet Cedex, France

² INRA, Unité GAFL (UR1052), CS 60094, 84143 Montfavet Cedex, France

17emes Rencontres de Virologie Végétale 27 au 31 janvier 2019 Aussois - France RVV 2019



Aussois du 27 au 31 janvier 2019