

Mobilizing tropical plants as a sustainable alternative to the issue of anthelmintic resistance in small ruminants

MARIE-MAGDELEINE C¹., BERNIER R¹., FÉLICITÉ Y¹., PHILIBERT L.¹, ARECE GARCIA J.²

¹ URZ Recherches Zootechniques, INRA, 97170, Petit-Bourg (Guadeloupe), France

² EEPF "Indio Hatuey", Universidad de Matanzas, Central España Republicana, CP44280-Matanzas, Cuba

Contacts: carine.marie-magdeleine-chevry@inra.fr

Caribbean Science and Innovation Meeting



Keywords : Condensed tannins, Gastrointestinal nematode, *Haemonchus contortus*, Anthelmintic resistance.

Introduction

Condensed tannins (CT) are **complex polyphenolic secondary metabolites** from plants. **Protein-CT complexes in ruminants** could result in affection of **rumen fermentation** (Min et al., 2001), allow **availability of amino-acids** and induce a gastrointestinal **nematicidal action** (Marie-Magdeleine et al., 2010). A preliminary study was conducted in order to evaluate *in vitro* the effect of several CT types, against various **chemical-resistant strains** of the parasite *Haemonchus contortus*.

Materials and Methods

Study of **CT from 8 plant species from 6 botanical families** (CT 1 to CT 8).

• **Extraction**: acetone/water (3:7; v/v).

• **Chemical quantification and elucidation**: thiolysis, LC-MS : Procyanidin/Prodelphinidin ratio

• **Biological activity** : **Drug resistance** against thiabendazole (Thia), Moxidectin (Mox), levamisole (Lev), ivermectin (Ivm) ; and CT **Anthelmintic *in vitro* assays** against L3 stage migration and exsheathment of 5 strains of *H. contortus* (B, C, G, J and R) : 5 concentrations of CT and 3 repetitions, PBS control. Calculation of efficient concentration for 50% inhibition EC50.

Results

Figure 1. Drug resistance of *H. contortus* strains

% L3 migration			
Chemical drug	<i>H. contortus</i> strain		
Thia	57.6 ^a	B	18.0 ^{cd}
Mox	33.4 ^b	C	47.9 ^a
Lev	8.5 ^c	G	11.1 ^d
Ivm	8.1	J	33.6 ^b
		R	24.1 ^{bc}

Strains G, J, B and R : Resistant to MOX and THIA, susceptible to IVM and LEV

Strain C : Multi-resistant to IVM, LEV, MOX and THIA

Figure 2. Prodelphinidin content in condensed tannins





Figure 3. Effects of condensed tannins on the exsheathment kinetics of three strains of *H. contortus*



Conclusion

It seems that the **structural complexity of condensed tannins**, the presence of other active compounds in the plant extracts, but also **the nature of the parasite resistance**, could affect the **tannin-protein interactions**.

Acknowledgements The authors would like to thank UE PTEA and URZ lab for experimental support, and CRB CARARE for parasite material. This research was supported by the FEDER Region Guadeloupe CPER CRB project.

References Marie-Magdeleine, C., Boval, M., Philibert, L., Borde, A., Archimède, H., 2010. Livestock Science 131, 234-239.

October 19th-23rd 2019

