



HAL
open science

Addressing organic viticulture environmental burdens by better understanding causes of inter-annual impacts variations

Christel Renaud-Gentié, Valentin Dieu, Marie Thiollet-Scholtus, Aurélie Perrin, Séverine Julien, Sandra Beauchet, Frédérique Jourjon, Anne Merot

► To cite this version:

Christel Renaud-Gentié, Valentin Dieu, Marie Thiollet-Scholtus, Aurélie Perrin, Séverine Julien, et al.. Addressing organic viticulture environmental burdens by better understanding causes of inter-annual impacts variations. 11. International Conference on Life Cycle Assessment of Food 2018 (LCA Food), Kasetsart University [Siracha Campus] (KU). THA., Oct 2018, Bangkok, Thailand. hal-02735830

HAL Id: hal-02735830

<https://hal.inrae.fr/hal-02735830v1>

Submitted on 2 Jun 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

LCAF-2018-07-00307

Addressing organic viticulture environmental burdens by better understanding causes of inter-annual impacts variations

Christel Renaud-Gentié^{1*}, Valentin Dieu², Marie Thiollet-Scholtsus³, Aurélie Perrin¹, Séverine Julien¹, Sandra Beauchet^{1,4,5}, Frédérique Jourjon¹, Anne Mérot².

¹USC 1422 GRAPPE, Univ. Bretagne Loire, Ecole Supérieure d'Agricultures (ESA)-INRA, SFR 4207 QUASAV, 55 rue Rabelais, 49007 Angers, France. ²INRA, UMR 1230 System, Montpellier France. ³INRA, UMR 0055 Aster, Colmar, France. ⁴ADEME, SAF, 20, Avenue du Grésillé, 49000 Angers, France

⁵Now with IFP Energies nouvelles, nouvelles, 1 et 4 Avenue de Bois-Préau, 92852 Rueil-Malmaison, France

Abstract

Organic viticulture is developing rapidly due to consumers demand and state incentives in some EU countries. Nevertheless, it has shown higher inter-annual variability in its environmental impacts than conventional viticulture. Therefore, the organic winegrowers would benefit from a better understanding of this variability in order to better address their environmental impacts. A study was conducted in four contrasted pedoclimatic conditions and two contrasted years in terms of climate and of pest and disease pressure. LCAs of organic wine grape was calculated based on detailed inventories of data from the eight real vineyard situations.

Main contributor to impacts was diesel combustion. The inter-annual variation was different between the plots. The impacts that varied the most were the freshwater and soil eco-toxicities, marine eutrophication, freshwater eutrophication and metal depletion. The main agricultural operations contributing to impact variations were disease management due to disease pressure related to climatic conditions variations.

Keywords: sustainable agriculture; Cradle-to-farm gate LCA; variability; climate, diesel, copper.

*Corresponding author. Tel.: +33-2-41-23-55-55

E-mail address: c.renaud@groupe-esa.com

1 Introduction

Organic viticulture covers 332,905 ha in the world, of which 88% is in Europe (Agence-Bio, 2017), and in some countries, such as Spain and Italy, there has been two-digit growth due to an increasing demand for organic wines and the support of public subsidies. However, organic viticulture may be more subject to inter-annual variations of intervention frequency than conventional one (Beauchet et al., accepted with minor corrections), especially due to fungal pressure and organic fungicides washability by rain leading to important variability in environmental impacts. This sector will benefit from maintaining its very good image in consumers' minds (Jourjon and Symoneaux, 2014) by addressing its environmental impacts, through a better understanding of the inter-annual and site-related variations of impacts and their causes.

This paper deals with quantification and causes identification of the environmental performance of four different organic technical management routes (TMRs), their causes, and variability in order to propose improvement drivers. After presenting the

cases studied and methods, we identify and discuss the main hotspots and their variation causes.

2 Material and methods

Four real vineyard plots were studied for two years with contrasting climate or diseases pressure in three French wine regions: One in Alsace, one in Provence Alpes Côte d'Azur and two in the Loire Valley, (Table 1). In each region, the cases were chosen so as to be representative of one or two different types of organic TMRs by applying Renaud-Gentié *et al.* (2014) method.

Life Cycle Assessment (LCA) was conducted using Recipe and USETox™ as characterization methods. Soil erosion [T.ha⁻¹] for heavy metals and phosphorus emission calculations was calculated thanks to Rusle2, then emission models were the same as Rouault *et al.* (2016). Inventory included all operations performed in the vineyard during the year studied and the related inputs.



Table 1: Main characteristics of the contrasted plots (*in italic*) and vineyard operations of the TMRs.

TMR Name	Rou68		Cre84		Hil49		Sav49	
<i>Grape variety</i>	<i>Riesling</i>		<i>Grenache Noir</i>		<i>Chenin Blanc</i>		<i>Chenin Blanc</i>	
<i>PDO/Region</i>	<i>Alsace/Alsace</i>		<i>Ventoux/ Provence- Alpes -côte d'Azur</i>		<i>Saumur Loire Valley</i>		<i>Savennières/ Loire Valley</i>	
<i>Soil type</i>	<i>Sandy-clay- silty</i>		<i>Sandy Silty</i>		<i>-Silty</i>		<i>Sandy-Clay</i>	
<i>Climate</i>	<i>Continental</i>		<i>Mediterranean</i>		<i>Temperate- oceanic</i>		<i>Temperate- oceanic</i>	
<i>% of soil covered by grass</i>	100		40		30		80	
Main Inventory elements								
Year	2013	2015	2013	2015	2010	2012	2010	2012
Number pesticide spraying	6	6	9	8	9	12	10	11
Quantity of copper applied in the year (g.ha⁻¹)	2832	5660	2880	2160	2648	5811	3480	4683
Quantity of sulfur applied in the year (g.ha⁻¹)	24500	69320	84850	83100	40000	51200	36480	50680
Number mechanical operations (except soil and pesticide)	6	6	4	4	6	4	3	2
Nb soil management operations	9	9	7	6	11	9	6	5
Total hours of tractor	39	39	19,6	17,63	51	45,33	11,75	33,41
Total hours manual labour	77,2	76,7	258	290	249	236	296	340
Yield (kg.ha⁻¹)	11250	11250	3075	8377	3600	5250	2250	2250

On the basis of LCA results, for each of the four studied situations, the percentage of inter-annual variation was calculated for each impact category and for each type of operation, in order to identify the main contributors to the total impact variation. Due to limited room, a selection of results is presented here.

3 Results and discussion

Considering the LCA results of the four cases for the two years, the main causes of the environmental impacts were identified (Table 2). Diesel was found as one of the major contributors in all the impacts, followed by the emissions from plant protection (copper) or not directly related to operations (N, P, heavy metals from the soil reserve). Fertilisation being not applied on all the plots did not appear as a major contributor.

We observed a variation of impacts between the contrasted years from 0 to 235% according to the plot and to the impact category for a cumulated variation of all impacts of 182% up to 564% (Figure1). The two plots situated in the Loire Valley showed the highest variability.

The impacts that varied the most (Figure 2) were relating to ecotoxicity (terrestrial, freshwater and marine) as well as freshwater eutrophication and metal depletion.

Amongst the seven groups of vineyard operations and two types of emissions non-related to operations, the main contributors to inter-annual variation are summarized in the table 3.

The annual climate, influenced:

- fungal pressure and thus the number (implying each time a mechanical operation) and dose of fungicide treatments required
- weed growth and thus the number of mechanical soil management operations
- vine growth and thus the number of mechanical operations
- erosion and lixiviation and as a consequence nitrogen and phosphorous emissions from fertilisers or from soil reserve

Table 2: Main causes of the impacts in the eight LCA results, causes are classified according to their number of occurrence as first contributor (in brackets, the related vineyard operations).

Main contributors	Climate Change	Terrestrial Acidification	Freswater ecotoxicity	Terrestrial ecotoxicity	Water depletion	Fossil depletion
First	Diesel combustion (soil, canopy and diseases management)	Sulfur production (diseases management sulfur)	Emissions grape production (not related to a specific operation)	Emissions of plant protection (diseases management copper)	Diesel combustion (soil, canopy and diseases management)	Diesel combustion (soil, canopy and diseases management)
Second	Background emissions in the field (not related to any operation)	Diesel combustion (soil, canopy and diseases management)	Diesel combustion (soil, canopy and diseases management)		Tractor manufacturing (soil, canopy and diseases management)	Tractor manufacturing (soil, canopy and diseases management)
Third	Tractor manufacturing (soil, canopy and diseases management)	Emissions from Fertilisation	Emissions plant protection (diseases management)		Treatments (diseases management)	Fertiliser manufacturing




Figure 1 (left): Cumulated percentage of variation of all impact categories between two contrasted years for four organic vineyard TMRs; Figure 2 (right): Average percentage of variation per impact category for four organic vineyard TMRs.

The winegrower's technical choices regarding type of organic fertiliser, or different forms of copper (used as fungicide), and tractor speed also influenced the inter-annual variation.

The main pathways for environmental performance improvements are i) reducing fuel consumption by either limiting the number of operations or/and using more fuel-efficient machines or different energy sources; adjusting the tractor speed; ii) reducing the dose and emission of copper products by the use of tunnel sprayers, and iii) limiting nitrogen emissions

through the choice of fertiliser type and management, limiting the use of fertilisers using manganese coming from manganese extraction industry; iv) design the vineyard plot in order to limit erosion (rows perpendicular to slope, use of grass cover) .

4 Conclusions

This study provides an overview of the hotspots, causes and variability of the environmental impacts of organic viticulture in contrasting climates and sites, as well as improvement paths.

Table 3: Main contributors and drivers to inter-annual variation of impacts for four organic vineyard TMRs.

Impact categories	Operations or non-related to operations emissions	Drivers
All impact categories	Soil management operations duration (mainly plot SAV (150% variation) but also the others to a lesser extent (up to 50%))	Weed growth due to climate of the spring, and change in the tractor speed (strategy of the winegrower)
	Occasional operations (replacement of dead plants) (plot HIL)	Strategy of the vinegrower (impact to be amortised on more years)
Water depletion metal depletion	Dose of manganese sulfate foliar fertiliser (plot CRE),	Strategy of the vinegrower
	Duration of soil management operations (plot SAV)	Weed growth due to climate of the spring, and change in the tractor speed (strategy of the winegrower)
Soil ecotoxicity, soil acidification, marine ecotoxicity, freshwater and marine eutrophication	Number of pesticide application and the nature of copper used (all the plots)	Fungal pressure due to climate of the spring and summer
Photochemical oxidation, climate change soil acidification	Background emissions (N and P) (plots HIL and CRE)	Effect of climate on emissions
Marine Eutrophication photochemical oxidation soil acidification	Mechanical operation (plots HIL and SAV)	Canopy growth and fungal pressure, due to the climate of the spring

Acknowledgements: The authors thank F. Ajem, Z. Bibes, N. Oumarou-Koura, A. Rouault, and the winegrowers; Vibrato & Qualenvic projects for their financial support, the reviewers for their contribution to the improvement of the paper.

References

Agence-Bio. La bio dans le Monde. Paris, France: Agence française pour le développement et la promotion de l'agriculture biologique; 2017.

Beauchet, S.; Rouault, A.; Renouf, M.A.; Thiollet-Scholtus, M.; Jourjon, F.; Renaud-Gentié, C. Inter-annual variability in the environmental performance of viticultural technical management routes - a case study in the Middle Loire Valley (France). international Journal of Life Cycle Assessment. accepted with minor corrections.

Jourjon, F.; Symoneaux, R. AOC versus environnement : quelle perception par les consommateurs et quel levier pour la compétitivité des vins français ? , 37th World Congress of Vine and Wine. Mendoza, Argentina; 2014.

Renaud-Gentié, C.; Burgos, S.; Benoît, M. Choosing the most representative technical management routes within diverse management practices: Application to vineyards in the Loire Valley for environmental and quality assessment. Eur J Agron. 2014;56(0):19-36.

Rouault, A.; Beauchet, S.; Renaud-Gentie, C.; Jourjon, F. Life Cycle Assessment of viticultural technical management routes (TMRs): comparison between an organic and an integrated management route. OENO One. 2016 2016-07-27;50(2).

Conference Proceedings

11th International Conference on
Life Cycle Assessment of Food 2018 (LCA Food)

in conjunction with

6th LCA AgriFood Asia

and 7th International Conference on
Green and Sustainable Innovation (ICGSI)

*“Global food challenges
towards sustainable
consumption and production”*

16 – 20 October 2018
Bangkok, Thailand

Organized by



Programme and Book of Abstracts

The 11th International Conference on
Life Cycle Assessment Food 2018 (LCA Food 2018)

“Global food challenges towards sustainable consumption and production”

in conjunction with

the 6th LCA AgriFood Asia

and

the 7th International Conference on Green and Sustainable Innovation (ICGSI 2018)

Organized by

Centre of Excellence on enVironmental strategy for GREEN business (VGREEN) – Kasetsart University (KU)

The Joint Graduate School of Energy and Environment (JGSEE) – King Mongkut’s University
of Technology Thonburi (KMUTT)

National Science and Technology Development Agency (NSTDA)