



HAL
open science

Monitoring aphid population dynamics: towards a better understanding of virus epidemics in melon crops

Alexandra Schoeny, Patrick Gognalons, Gregory Girardot, Pauline Millot, Karine Nozeran, Catherine Wipf-Scheibel, Hervé Lecoq

► To cite this version:

Alexandra Schoeny, Patrick Gognalons, Gregory Girardot, Pauline Millot, Karine Nozeran, et al.. Monitoring aphid population dynamics: towards a better understanding of virus epidemics in melon crops. 14. International Plant Virus Epidemiology Symposium (IPVE 2019), Korean Society for Plant Pathology (KSPP)., May 2019, Séoul, South Korea. hal-02736004

HAL Id: hal-02736004

<https://hal.inrae.fr/hal-02736004v1>

Submitted on 2 Jun 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Monitoring aphid population dynamics: towards a better understanding of virus epidemics in melon crops

A. Schoeny, P. Gognalons, G. Girardot, P. Millot, K. Nozeran, C. Wipf-Scheibel, and H. Lecoq

Pathologie Végétale, INRA, Montfavet, France
(*alexandra.schoeny@inra.fr*)

In France, open field melon crops are regularly impacted by four aphid-borne viruses: cucurbit aphid-borne yellows virus (CABYV), cucumber mosaic virus (CMV), watermelon mosaic virus (WMV) and zucchini yellow mosaic virus (ZYMV). The efficiency of control methods is likely to be enhanced with an accurate knowledge of epidemic drivers in particular those linked with aphid vectors. Field experiments were conducted in Southeastern France between 2010 and 2018 to investigate the relationships between aphid population dynamics and virus epidemics. Winged aphids visiting melon crops were sampled daily using non-biased suction traps and aphid species were identified under a stereomicroscope. Viruses were monitored weekly by DAS-ELISA. Gompertz models were fitted to virus incidence data sets by nonlinear regression and AUDPCs (Area Under the Disease Progress Curve) were calculated. A statistical analysis was performed to explore the relationships existing between several “aphid” variables (total aphid abundances and specific abundances over different periods of time) and several “virus” variables (cumulative total of infected plants over different periods of time, newly infected plants per week, AUDPCs, Gompertz model parameter estimates). No significant relationship was highlighted between aphids and non-persistent viruses (CMV, WMV, ZYMV). Interestingly, a predictive relationship was established between *Aphis gossypii* population dynamics and CABYV epidemics suggesting that an early control of the population of *Aphis gossypii* could impact favourably the epidemic onset and progress of this persistent virus in melon crops.



14th IPVE
International Plant Virus Epidemiology
Symposium

14th IPVE

International Plant Virus Epidemiology
Symposium

Coex, Seoul, South Korea | May 13-17th 2019



Rural Development
Administration



National Institute of
Horticultural and Herbal Science



Korea Institute of Planning and
Evaluation for Technology
in Food, Agriculture and Forestry



The Korean Society of Plant Pathology