



HAL
open science

Genomics to estimate additive and dominance genetic variances in purebred and crossbred pig traits

Llibertat Tusell, Hélène Gilbert, Zulma Vitezica, Marie-José Mercat, Andres Legarra, Catherine Larzul

► To cite this version:

Llibertat Tusell, Hélène Gilbert, Zulma Vitezica, Marie-José Mercat, Andres Legarra, et al.. Genomics to estimate additive and dominance genetic variances in purebred and crossbred pig traits. 68. Annual Meeting of the European Association for Animal Production (EAAP), Aug 2017, Tallinn, Estonia. Wageningen Academic Publishers, Annual Meeting of the European Association for Animal Production, 23 (1ère Ed.), 513 p., 2017, Book of Abstracts of the 68th Annual Meeting of the European Federation of Animal Science. hal-02736114

HAL Id: hal-02736114

<https://hal.inrae.fr/hal-02736114v1>

Submitted on 2 Jun 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - ShareAlike 4.0 International License

Genomics to estimate additive and dominance genetic variances in purebred and crossbred pig traits*L. Tusell¹, H. Gilbert¹, Z.G. Vitezica¹, M.J. Mercat², A. Legarra¹ and C. Larzul¹**¹INRA, UMR1388, INPT ENSAT, INPT ENVT GenPhySE, 31326, Castanet-Tolosan, France, ²IFIP/BIOPORC, La Motte au Vicomte, 35651 Le Rheu, France; helene.gilbert@inra.fr*

This study aims at assessing the contribution of the additive and dominance genomic variances to the phenotypic expression of several purebred Piétrain and crossbred (Piétrain × Large White) pig performances. A total of 636 purebred and 720 crossbred male piglets were phenotyped for 22 traits that can be classified into the trait groups growth rate and feed efficiency, carcass composition, meat quality, behaviour, boar taint and puberty. Additive and dominance variances estimated in univariate genotypic models including additive and dominance genotypic effects and a genomic inbreeding covariate allowed us to retrieve the additive and dominance SNP variances for purebred and crossbred performances. These estimated variances were used, together with the allelic frequencies of the parental populations, to obtain additive and dominance variances in terms of genomic breeding values and dominance deviations. Estimates of additive genetic variances across traits were consistent with previous results without dominance indicating that additive and dominance genetic effects were non-confounded. Some traits showed a relevant amount of dominance genetic variance in both populations (i.e. growth rate 8%, feed conversion ratio 9-12%, backfat thickness 14-12%, lean meat 10-8%, carcass lesions 9%, in purebreds and crossbreds, respectively) or increased amounts in crossbreds (i.e. ham cut 8-13%, loin 7-16%, pH semimembranosus 13-18%, pH Longissimus dorsi 9-14%, dressing yield 5-15%, androstenone 5-13% and estradiol 6-11%). Results suggest that accounting for dominance in the models of these traits could lead to an increased GEBV accuracy and that using crossbred information can be beneficial to evaluate purebred candidates to selection for crossbred performance. Further research will compare additive and dominance marker effects between crossbred and purebred performances.