



HAL
open science

Cadmium and Deoxynivalenol in durum wheat grains: physiological and biological basis of the co-contamination

Florence Richard-Forget, Vessela Atanasova-Penichon, Sylvain Chéreau, Jean-yves Cornu, Christine Ducos, Valerie Nicaise, Laetitia Pinson-Gadais, Nadia Ponts, Marie-Noëlle Verdal-Bonnin, Quentin Gras

► To cite this version:

Florence Richard-Forget, Vessela Atanasova-Penichon, Sylvain Chéreau, Jean-yves Cornu, Christine Ducos, et al.. Cadmium and Deoxynivalenol in durum wheat grains: physiological and biological basis of the co-contamination. 14. European Fusarium Seminar, Universität für Bodenkultur Wien [Vienne, Autriche] (BOKU). AUT., Apr 2018, Tulln, Austria. 122 p. hal-02736250

HAL Id: hal-02736250

<https://hal.inrae.fr/hal-02736250>

Submitted on 2 Jun 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution| 4.0 International License



April 8 – 11, 2018
Minoritenkloster
Tulln, AUSTRIA

ORGANIZING COMMITTEE

Gerhard Adam
(BOKU, AT)

Rainer Schuhmacher
(BOKU, AT)

Hermann Bürstmayr
(BOKU, AT)

Joseph Strauss
(BOKU, AT)

Rudolf Krska
(BOKU, AT)

Gerlinde Wiesenberger
(BOKU, AT)

Dieter Moll
(BIOMIN, AT)

Alexandra Khassidov
(ÖGMBT, AT)

INTERNATIONAL SCIENTIFIC COMMITTEE

Sarah M. De Saeger
(BE)

Ruth Dill-Macky
(US)

Doris Marko
(AT)

Thomas Miedaner
(DE)

Antonio Moretti
(IT)

Michele Suman
(IT)

B. Gillian Turgeon
(US)

Table of Contents

Table of Contents	2
General Information	3
Welcome Address	4
Invited Speakers	5
Detailed Program	13
SUNDAY - April 8	13
MONDAY - April 9	13
TUESDAY - April 10	16
WEDNESDAY - April 11	19
Posters Table	21
Lectures	22
Welcome Note & Plenary Session	23
Session 1: Fusarium genomics and virulence mechanisms, population genetics and diversity	25
Session 2: Host resistance: mechanisms, genetics, genomics and breeding	32
Session 3: Fusarium secondary metabolites and metabolomics of Fusarium-host plant interactions	39
Session 4: Fusarium mycotoxins - Toxicology, Metabolism and Remediation	46
Session 5: Integrated Fusarium management (pre/postharvest, epidemiology and forecasting, fungicide resistance, utilization of contaminated batches)	54
Poster Presentations	61
Author Index	110
Participant Index	115
Imprint	123

PP-83 Changes in expression of stress-responsive genes in *Fusarium proliferatum* exposed to host plant extracts

Łukasz Stępień¹, Natalia Witaszak¹, Agnieszka Waśkiewicz², Justyna Lalak-Kańczugowska¹

¹ Department of Pathogen Genetics and Plant Resistance, Institute of Plant Genetics of the Polish Academy of Sciences, Poland

² Department of Chemistry, Poznań University of Life Sciences, Wojska Polskiego 75, 60-625 Poznań, Poland

Stress factors (biotic or abiotic) are critical players in functional characteristics of plant-pathogenic fungi. Our previous studies have shown that *Fusarium proliferatum*, a casual fungal pathogen, besides increased fumonisin biosynthesis, activates the expression and accumulation of numerous proteins to minimize the effect of potential stressing factors, such as host plant extracts. Some of them are related with stress response, e.g. the heat-shock proteins (HSPs) which often show protective and stimulating biological functions. Therefore, here we tried to understand the role of HSPs in response to stress conditions exerted by the presence of plant metabolites from host species. The main objective of present study was to examine the changes in the expression of two *Hsp* genes after the addition of aqueous extracts from four potential host species in *Fusarium proliferatum*.

Asparagus-derived *F. proliferatum* strains was cultured *in vitro* using a liquid medium, which at the 5th day of cultivation was supplemented with plant extracts made from asparagus, maize, garlic and pineapple tissues. Mycelia were collected every two days from liquid cultures before and after extract added and subjected to the total mRNA extraction. To validate the expression profiles of *Hsp70* and *Hsp88* genes, specific primers were designed based on previously obtained partial genomic sequences and quantitative RT-PCR technique was used and normalized against the constitutively expressed *tub2* gene.

The expression levels of *Hsp70* and *Hsp88* genes varied relating to host extract used. The *Hsp88* gene was expressed in all samples whereas the expression of the *Hsp70* gene was observed 24 h after the extract supplementation. The asparagus extract induced the highest increase of *Hsp70* and *Hsp88* transcript levels and garlic extract was the second most effective inducer. In the *Hsp88* expression profile we observed two expression peaks: after 24 h of extract supplementation and after 12 days of cultivation (7 days after extract application). We have demonstrated that dissimilar nutritive environments have significant impact on stress-related gene expression profiles in *F. proliferatum*.

This research was supported by the Polish National Science Centre Project 2015/17/B/NZ9/03577.

PP-84 Cadmium and Deoxynivalenol in durum wheat grains: physiological and biological basis of the co-contamination.

Florence Richard-Forget¹, Vessela Atanasova-Penichon¹, Sylvain Chereau¹, Jean-Yves Cornu², Christine Ducos¹, Valérie Nicaise², Laetitia Pinson-Gadais¹, Nadia Ponts¹, Marie-Noëlle Verdal-Bonnin¹, Quentin Gras¹

¹ UR1264 MycSA, INRA, France

² INRA, UMR 1391 ISPA, 71 Avenue Edouard Bourlaux, CS20032, 33 883 Villenave d'Ornon, France

Cadmium (Cd) and mycotoxins are among the most worrying contaminants that threaten the safety of food products derived from cereal kernels. Indeed, as recently supported by the last French total diet survey (Anses, 2011), deoxynivalenol (DON) and Cd human exposure through food mainly results from the consumption of cereal-derived products. Durum wheat is the most sensitive cereal culture for both DON and Cd accumulation in kernels, leading to a high frequency of co-contaminated harvests. This frequent co-occurrence (even though each toxic substance is in concentration within the EU regulatory limits) combined with the fact that Cd and DON are likely to be distributed in the same milling fractions raises the concern of consumer exposure to the cocktail Cd+DON.

To address the issue of DON+Cd co-contamination, the CaDON initiative (funded by the French National Agency for Research, 2015-2019) investigates the relations between Cd and DON occurrence in durum wheat, from crops co-contamination in the field to the milling end products, as well as the toxicity of Cd and DON mixtures upon ingestion. One of the objectives pursued by the CaDON project aims to elucidate the physiological bases of Cd+DON contamination of durum wheat kernels. The effect of soil Cd contamination on *Fusarium graminearum* infection and DON accumulation in kernels and conversely of *F. graminearum* infection on Cd accumulation are investigated through the implementation of *in vitro* and greenhouse experiments. As a first step, the way *F. graminearum* exposition to Cd affects the production of DON is analysed and the mechanisms underlying this modulation investigated with a strong focus on the relation between Cd/oxidative stress/DON biosynthesis. These insights will be the subject of the present communication together with the first data delivered by greenhouse experiments.

Anses, 2011. <https://www.anses.fr/fr/content/les-etudes-de-l'alimentation-totale-eat>