



HAL
open science

A first insight into the effect of Lotilaner on GABA-gated channels from the european tick *Ixodes ricinus*

Claude Charvet, Clément Auger, Claude Risper, Caroline Hervet, Elise Courtot, Steeve Thany, Olivier Plantard, Cédric Neveu

► **To cite this version:**

Claude Charvet, Clément Auger, Claude Risper, Caroline Hervet, Elise Courtot, et al.. A first insight into the effect of Lotilaner on GABA-gated channels from the european tick *Ixodes ricinus*. 27. Conference of the World Association for the Advancement of Veterinary Parasitology, Jul 2019, Madison, United States. 349 p. hal-02736790


HAL Id: hal-02736790

<https://hal.inrae.fr/hal-02736790v1>

Submitted on 2 Jun 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



WAAVP

27th Conference of the World Association for
the Advancement of Veterinary Parasitology

JULY 7 - 11, 2019 | MADISON, WI, USA

Dedicated to the legacy of Professor Arlie C. Todd

Sifting and Winnowing the Evidence in Veterinary Parasitology



f @WAAVP2019  @WAAVP_2019

Abstract Book

*Joint meeting with the 64th American Association of Veterinary Parasitologists
Annual Meeting & the 63rd Annual Livestock Insect Workers Conference*

www.WAAVP2019.com **#WAAVP2019**

humidity). Geometric means of combined live tick counts were calculated for attachment intervals of 1-2 hours, 3-4 hours, 5-6 hours, and 7-8 hours, and least squares means were compared by ANOVA with a two-sided significance level set at $\alpha=0.05$. Significantly fewer live *I. scapularis* were removed from treated than control dogs after 3-4 hours of attachment ($p=0.0012$); upon holding 24 hours, significantly fewer *I. scapularis* that had previously been attached to treated dogs for 1-2 hours were alive ($p=0.0049$). Significantly fewer live *A. americanum* were removed from treated than control dogs after 7-8 hours of attachment ($p=0.0254$); upon holding 24 hours, significantly fewer *A. americanum* that had previously been attached to treated dogs for 3-4 hours were alive ($p=0.0003$). These data indicate that acaricidal activity of sarolaner against *I. scapularis* and *A. americanum* begins after 1-2 hours and 3-4 hours of attachment, respectively, on sarolaner treated dogs.

OA13.04 A First Insight Into the Effect of Lotilaner on GABA-Gated Channels From the European Tick *Ixodes Ricinus*

Dr Claude Charvet¹, Clément Auger¹, Dr Claude Rispe², Caroline Hervet², Dr Elise Courtot¹, Pr Steeve Thany³, Dr Olivier Plantard², Dr Cédric Neveu¹

¹INRA, ISP, Université Tours, UMR1282, Nouzilly, France, ²INRA, Oniris, BIOEPAR, Nantes, France, ³Université d'Orléans, LBLGC, USC1328 INRA, Orléans, France

Ticks are strict blood-feeding arthropods (Acari), which represent a major health issue for wild or domesticated animals and humans, due to their potential to transmit disease agents. Control of ticks is increasingly difficult due to the development of drug-resistant parasites. Ligand-gated ion channels of the tick central nervous system are the primary targets of acaricides. Among those receptors, the γ -aminobutyric acid-gated chloride ion channels (GABACl) are the main synaptic inhibitory receptors. Lotilaner is a recently developed parasiticide from the isoxazoline chemical class that was shown to be a non-competitive antagonist of GABACl from the livestock tick *Rhipicephalus microplus*. In the

present study, we characterized the GABACl from the European tick species *Ixodes ricinus*.

We extracted RNAs from *Ixodes ricinus* nymphs. Taking advantage of the phylogenetic closeness of *I. ricinus* and *R. microplus* in the Arthropoda phylum, we identified the *I. ricinus* GABACl subunit homologue. The cDNA encoding the Iri-GABACl was cloned and the corresponding in vitro synthesized cRNAs were micro-injected into *Xenopus laevis* oocytes to investigate its pharmacological properties. Functional expression and two-electrode voltage clamp studies demonstrated that the GABACl subunit formed a homomeric receptor gated by GABA. Importantly, the insecticides like lotilaner, fipronil and picrotoxin efficiently blocked the GABA currents as previously observed for the *R. microplus* GABACl. Surprisingly, *I. ricinus* GABACl was not sensitive to the pesticide dieldrin, suggesting a potential naturally existing resistance mechanism involving alternative exons.

Here we report the functional characterization of the first GABACl of *I. ricinus* demonstrating that it is an important molecular target for lotilaner. Transcriptomic analysis of *I. ricinus* are in progress to identify new acaricidal targets.

OA13.05 Clinical Efficacy of Afoxolaner in Dogs Naturally Infected With *Sarcoptes Scabiei* and Concomitant Modifications of the Skin Microbiota

Raluca Mindru², Andrei Cristian Lupu², Dr. Constantin Roman², Dr. Charlotte Bernigaud⁵, Dr. Amaury Briand¹, Dr. Frédéric Beugnet³, Dr. Katja Fischer⁴, Professor Liviu Miron², **Professor Jacques Guillot¹**

¹Veterinary College of Alfort, Maisons-alfort, France, ²Ion Ionescu de la Brad, Faculty of Veterinary Medicine, Iasi, Romania, ³Boehringer Ingelheim Animal Health, Lyon, France, ⁴QIMR Berghofer, Medical Research Institute, Brisbane, Australia, ⁵Henri Mondor Hospital, France

Infection by *Sarcoptes scabiei* remains a common disease in dogs, especially in tropical countries where most of stray or non-controlled dogs show cutaneous