



Mono (2-ethylhexyl) phthalate induces transcriptomic and proteomic alterations in bovine oocytes.

Dorit Kalo, Anais Vitorino Carvalho, Catherine Archilla, Véronique Duranthon, Marco Moroldo, Yishai Levin, Meital Kupervaser, Yoav Smith, Zvi Roth

► To cite this version:

Dorit Kalo, Anais Vitorino Carvalho, Catherine Archilla, Véronique Duranthon, Marco Moroldo, et al.. Mono (2-ethylhexyl) phthalate induces transcriptomic and proteomic alterations in bovine oocytes.. 35. Annual Meeting A.E.T.E. 2019, Sep 2019, Murcia, Spain. Animal Reproduction, 6 (3), 2019, Proceedings of the 35th Meeting of AETE. hal-02736954

HAL Id: hal-02736954

<https://hal.inrae.fr/hal-02736954>

Submitted on 2 Jun 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



A169E Embryology, developmental biology and physiology of reproduction

Mono (2-ethylhexyl) phthalate induces transcriptomic and proteomic alterations in bovine oocytes

Dorit Kalo^{1,2}, Anaïs Vitorino Carvalho³, Catherine Archilla³, Veronique Duranthon³, Marco Moroldo⁴, Yishai Levin⁵, Meital Kupervaser⁵, Yoav Smith⁶, Zvi Roth^{1,2}

¹Department of Animal Science, Robert H. Smith Faculty of Agriculture, Food and Environment, The Hebrew University of Jerusalem, Israel; ²Center of Excellence in Agriculture and Environmental Health, The Hebrew University of Jerusalem, Israel; ³UMR BDR, INRA, ENVA, Université Paris Saclay, Jouy-en-Josas 78350, France; ⁴Animal Biological Resources for Integrative and Digital Genomics Facilities, Jouy-en-Josas 78350, France; ⁵The de Botton Institute for Protein Profiling, The Nancy and Stephen Grand Israel National Center for Personalized Medicine, Weizmann Institute of Science, Rehovot 76100, Israel; ⁶The Bioinformatics Unit, Faculty of Medicine, the Hebrew University, Jerusalem 91120, Israel.

Keywords: phthalates, oocyte developmental competence, transcriptome.

Phthalates are plasticizers, used in a variety of industrial plastics. Di(2-ethylhexyl) phthalate (DEHP), the most commonly used plasticizer, and its main metabolite, mono(2-ethylhexyl) phthalate (MEHP), are known reproductive toxicants. A residual concentration of MEHP (~20 nM) has been found in follicular fluid aspirated from IVF-treated women and DEHP-treated cows. Previously we have reported that exposure of oocyte during maturation to MEHP impairs nuclear maturation, reduces cleavage and blastocyst formation rates. However, the effect of 20 nM MEHP on the transcriptomic profile of oocytes and their derived blastocysts is not entirely clear. Bovine oocytes were in-vitro matured with or without 20 nM MEHP for 22 h. At the end of maturation, they were collected for transcriptomic (by microarray; n = 20 per sample; 4 replicates) and proteomic (n = 200 per group) analyses to examine a possible direct effect of MEHP on the oocyte transcriptomic and proteomic profiles. The remaining oocytes were in-vitro fertilized and embryonic development was recorded 42–44 h and 7 days postfertilization. Blastocysts were also collected for microarray analysis (n = 10 per sample; 4 replicates). Transcriptomic data were analyzed using Partek Genomics Suite software. Control probes were removed; signals were log2 transformed followed by interslide quantile normalization. Genes were considered differentially expressed if the *P*-value by one-way ANOVA was lower than 0.05 and absolute fold change was 1.5 between the control and MEHP-treated group. Proteomic raw data were imported into Expressionist® followed by Mascot software. Data were searched against the bovine sequences from UniProtKB. Proteins were considered differentially expressed at a fold change of ± 1.5 with at least 2 unique peptides. Oocyte transcriptome analysis revealed MEHP-induced alterations in the expression of 456 genes. The differentially expressed genes were associated with actin cytoskeleton (n = 47 genes; e.g., *ACTG1*), metabolic pathway (n = 43) including oxidative phosphorylation (n = 12; e.g., *ND5*), oocyte maturation (n = 9; e.g., *PIK3CA*), and embryonic development (n = 14; e.g., *SOX10*, *NOTCH*); 191 proteins were affected by MEHP in mature oocytes, associated with methylation and acetylation (n = 51), metabolic pathway (n = 33) including mitochondrial oxidative phosphorylation (n = 7; e.g., *ATP5E*), and cytoskeleton structure (n = 32; e.g., *ACTN1*, *EGFR*). In control vs. MEHP-derived blastocysts, 290 genes were differentially expressed, associated with transcription process, cytoskeleton regulation and metabolic pathway; 9 of these genes were impaired in both oocytes exposed to MEHP (i.e., direct effect) and blastocysts developed from those oocytes (i.e., carryover effect). The study explores, for the first time, the risk associated with exposing oocytes to relevant MEHP concentrations (i.e., those found in the follicular fluid) to the maternal transcripts. Although it was the oocytes that were exposed to MEHP, alterations carried over to the blastocyst stage, following embryonic genome activation, implying that these embryos are of low quality.