



HAL
open science

Improving the durability and efficiency of plant resistance deployment using eco-evolutionary modelling

Loup Rimbaud, Julien Papaix, Benoît Moury, Luke G. Barrett, Peter H. Thrall

► To cite this version:

Loup Rimbaud, Julien Papaix, Benoît Moury, Luke G. Barrett, Peter H. Thrall. Improving the durability and efficiency of plant resistance deployment using eco-evolutionary modelling. 17. Eucarpia Meeting on Genetics and Breeding of Capsicum and Eggplant, Sep 2019, Avignon, France. INRA, Centre de recherche Provence-Alpes-Côte d'Azur, 263 p., 2019, Innovations in Genetics and Breeding of Capsicum and Eggplant. Proceedings of the 17th EUCARPIA Meeting on Genetics and Breeding of Capsicum and Eggplant September 11-13, 2019 | Avignon - France. hal-02737088

HAL Id: hal-02737088

<https://hal.inrae.fr/hal-02737088>

Submitted on 2 Jun 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial - NoDerivatives 4.0 International License

Improving the durability and efficiency of plant resistance deployment using eco-evolutionary modelling

Rimbaud L¹, Papaix J², Moury B¹, Barrett LG³, Thrall PH³

¹INRA, UR407 Pathologie Végétale, Avignon, France. ²INRA, UR BioSP, Avignon, France. ³CSIRO Agriculture & Food, Canberra, Australia

BACKGROUND Genetically-controlled plant resistance can reduce the damage caused by pathogens. However, pathogens have the ability to evolve and overcome the resistance. This often occurs very quickly after resistance is deployed, resulting in significant crop losses and continuing needs to breed new resistant cultivars. To tackle this issue, several strategies have been proposed to constrain the evolutionary potential of pathogen populations and thus increase resistance durability. These strategies mainly rely on using different combinations of resistance genes (e.g. qualitative and/or quantitative resistance) in time, space, or both (e.g. via gene pyramiding, cultivar rotations, cultivar mixtures, field mosaics). However, experimental assessment of the efficiency (i.e. ability to reduce disease impact) and the durability (i.e. ability to limit pathogen evolution and delay resistance breakdown) of different deployment strategies presents a major challenge.

MATERIALS & METHODS Therefore, we developed a spatially-explicit stochastic model [1] to assess the epidemiological and evolutionary outcomes of the major deployment options described above, for both qualitative (major resistance genes) and quantitative resistance (e.g. QTLs affecting different pathogen life-history traits). In addition, we analyzed the impact of landscape organization (as defined by the proportion of fields cultivated with a resistant cultivar, and their spatial aggregation) and epidemiological or evolutionary parameters (e.g. mutation probability, cost of infectivity).

RESULTS Our main results on resistance to rusts indicate that evolutionary and epidemiological control are not necessarily correlated [2], and that no deployment strategies is universally optimal [3].

DISCUSSION & CONCLUSION The model was first parameterized for cereal resistance to rusts (caused by fungi of the genus *Puccinia*), and is destined to be applied to other pathosystems including viruses of vegetable crops, especially Potato virus Y on pepper (*Capsicum spp.*).

REFERENCES

- [1] Rimbaud L, Papaix J, et al. 2018. PLoS Computational Biology 14(4): e1006067. DOI: 10.1371/journal.pcbi.1006067
- [2] Papaix J, Rimbaud L, et al. 2018. Evolutionary Applications 11(5): 705-717. DOI: 10.1111/eva.12570
- [3] Rimbaud L, Papaix J, et al. 2018. Evolutionary Applications 11(10):1791-1810. DOI: 10.1111/eva.12681




Figure 1. Model overview. (A) Model architecture. Healthy hosts can be contaminated by propagules and may become infected. Following a latent period, infectious hosts produce new propagules which may mutate and disperse across the landscape. At the end of the infectious period, infected hosts become epidemiologically inactive. Qualitative resistance prevents transition to the latent infected state (L). Green boxes indicate healthy hosts which contribute to crop yield and host growth, in contrast to latent hosts (dark blue box) and diseased hosts (i.e. symptomatic, red boxes). Model parameters associated with epidemiological processes are indicated in grey. Distributions used to simulate stochasticity in model transitions are indicated in red; B: binomial, Γ : gamma, P: Poisson, M: multinomial. Host growth is deterministic. (B) Two-dimensional representation of the power-law dispersal kernel calibrated for rust pathogens. Top panel indicates the logarithm of the probability to disperse from the origin to any point of the landscape; bottom panel indicates the cumulative probability of dispersing over a given distance. (C-D) Example of simulation with two major resistance genes deployed as a mosaic: (C) dynamic of diseased hosts; (D) landscape. Blue vertical lines indicate the durability of the two resistant cultivars. These lines delineate the three periods used to compute epidemiological outputs from AUDPC: short-term (ST, green area), transitory period (TP, grey) and long-term (LT, red).



Innovations in Genetics and Breeding of Capsicum and Eggplant

Proceedings of the 17th EUCARPIA Meeting on Genetics
and Breeding of Capsicum and Eggplant,

September 11-13, 2019 | Avignon - France

Editors: Véronique Lefebvre & Marie-Christine Daunay

Editors

Véronique Lefebvre & Marie-Christine Daunay

Title

Innovations in Genetics and Breeding of Capsicum and Eggplant

Sub-title

Proceedings of the 17th EUCARPIA Meeting on Genetics and Breeding of Capsicum and Eggplant
September 11-13, 2019 | Avignon - France

Publisher

Institut National de la Recherche Agronomique (INRA)

Centre de recherche Provence-Alpes-Côte d'Azur

228 route de l'aérodrome

CS 40 509 - Domaine Saint Paul, Site Agroparc,

84914 Avignon Cedex 9 - France

Visual identity

© Armelle Favery

Artistic director

© Lyonel Liger assisted by Sabine Laugier

Layout design and editing

Salima Kherchache

Printed by

SUD LABO, 35 avenue Pierre Sépard, 84000 Avignon - France