



HAL
open science

Impact of HTST pasteurization of human milk on the kinetic of digestion of macronutrients after in vitro dynamic digestion

S. Sottemano, Giulia Maiocco, Carlo Stefano Nebbia, Marzia Giribaldi, Olivia Ménard, Didier Dupont, Amélie Deglaire, Laura Cavallarin, Alexandra Coscia, Enrico Bertino

► To cite this version:

S. Sottemano, Giulia Maiocco, Carlo Stefano Nebbia, Marzia Giribaldi, Olivia Ménard, et al.. Impact of HTST pasteurization of human milk on the kinetic of digestion of macronutrients after in vitro dynamic digestion. 40th Congress of the Union of Middle-Eastern and Mediterranean Pediatric Societies, Oct 2019, Cagliari, Sardinia, Italy. 2019. hal-02737463

HAL Id: hal-02737463

<https://hal.inrae.fr/hal-02737463>

Submitted on 2 Jun 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Impact of HTST pasteurization of human milk on the kinetic of digestion of macronutrients after in vitro dynamic digestion

Stefano Sottemano¹, Giulia Maiocco¹, Stefano Nebbia², Marzia Giribaldi², Olivia Ménard³, Didier Dupont³, Amélie Deglaire³, Laura Cavallarin², Alessandra Coscia¹, Enrico Bertino¹.

1. Neonatal Unit of the University, City of Health and Science of Turin, Turin, Italy. 2. Institute of the Science of Food Production. CNR. Grugliasco. Italy. 3. Science et Technologie du Lait et de l'Oeuf. INRA. Agrocampus Ouest. Rennes. France.

Introduction

Donor human milk (HM) represents the best alternative when mother's own milk is not available, a common occurrence in Neonatal Intensive Care Units. Heat treatment of donor HM is mandatory for safety reasons: Holder pasteurization (HoP, 62.5°C-30') is recommended by all HM bank guidelines. A recent study [1] has demonstrated that HoP affects the digestion profile and behavior of several HM components. High Temperature-Short Time pasteurization (HTST, 71°C-15'') is currently under evaluation as a promising alternative technology to limit the denaturation of some biological compounds of raw HM and, very recently [2], it was shown to better preserve the milk antiviral activity with respect to HoP

Objectives

The aim of the present work was to assess whether the different types of pasteurization (HoP, HTST) impact the digestive kinetics of human milk during in vitro dynamic digestion.

Methodology

The experimental design is summarized in Fig. 1. HoP was performed by using the Human Milk Bank equipment (Metalardredinox, Italy), and HTST by a patented proprietary small-scale device [3].

Raw, HoP and HTST human milk were subjected to gastrointestinal digestion using DIDGI[®] system [4]; samples were characterized for:

- **Particle size distribution (PSD)** on undigested and gastric samples (Mastersizer 2000 laser light scattering with two laser sources, Malvern Instruments);
- **Confocal microscopy**: The microstructure was observed using Nikon C1Si confocal laser scanning microscopy (CLSM) on inverted microscope TE2000-E (Nikon, Champigny-sur-Marne, France) as previously described by Bourlieu et al. (2015). FastGreen[®] and LipidTox[®] (Thermo-Fisher Scientific) were used for staining simultaneously proteins (Blue) and apolar lipids (Green), respectively.
- **Protein profile**: (reducing and non reducing NuPAGE[®] 4-12%, Life Technologies) and MS identification of differential bands (Q Exactive[™] Mass Spectrometer, Thermo Scientific);
- **Amino acid (AA) profile** on undigested and intestinal samples (cation exchange chromatography, Biochrom 30 automatic AA analyzer, Biochrom);
- **Triglyceride content** (thin layer chromatography and densitometry analysis by Image Quant TL (GE Healthcare).

Statistical analyses were performed using the PAST3 software package.

Results

- Both pasteurization methods led to heat-induced protein aggregates (blue dots), both in the soluble phase and at the interface of the HM fat globule membrane. The protein-fat interaction phenomena, which could be seen also in raw human milk, seems to be more relevant following HoP treatment, with respect to HTST (Fig. 2)
- During gastric digestion, both pasteurization methods modified PSD, as compared to RHM (Fig. 3);
- Caseins were rapidly hydrolyzed in the gastric phase. Lactoferrin was hydrolyzed faster in the pasteurized samples in comparison to Raw HM, in which lactoferrin was more resistant to gastro-intestinal digestion. In particular, the kinetics digestion of iGa was statistically similar between RHM and HTST milk (Fig. 4);
- Heat-treatments, consequently, affected the intestinal release of AA, and a **significantly higher bioaccessibility** ($p < 0.05$) of AA was found for HTST, as compared to HoP (Fig. 5);
- **No difference** were found about Lipolysis between HoP and HTST (Fig. 6).

Conclusion

This work provides the first important evidences on the **differential impact of HoP and HTST pasteurization techniques on bioaccessibility of aa AND IgA of DHM** for preterm newborns.




Fig. 1: Experimental Design




Fig. 2: Confocal microscopy, protein aggregation in blue




Fig. 3: Size distributions of Gastric Digests particles of Raw and Pasteurized HM




Fig. 4: Non reducing NuPAGE profiles of HM protein digests from Raw (R) and pasteurized (P: HoP; H: HTST)




Fig. 5: % Free AA over Total sum (A) and Essential AAs (B).




Fig. 6: % Free TAG over initial quantity during Gastric and Intestinal phases.

Reference

- [1] Deglaire A, De Oliveira SC, Jardin J, Briard-Bion V, ..., Dupont D. Impact of human milk pasteurization on the kinetics of peptide release during in vitro dynamic term newborn digestion. Electrophoresis. 2016;37(13):1839-50
- [2] Donalisio M, Rittà M, Francese R, Civra A, ..., Lembo D. High temperature-short time pasteurization has a lower impact on the antiviral properties of human milk than holder pasteurization. Front Pediatr. 2018;6
- [3] Giribaldi M, Coscia A, Peila C, Antoniazzi S, ..., Cavallarin L. Pasteurization of human milk by a benchtop high-temperature short-time device. Innovative Food Sci Emerg Technol. 2016;36:228-33
- [4] Ménard O, Cattenoz T, Guillemin H, Souchon I, ..., Picque D. Validation of a new in vitro dynamic system to simulate infant digestion. Food Chem. 2014;145:1039-45