



**HAL**  
open science

## Role of Paneth cells during infection of neonatal mice by *Cryptosporidium parvum*

Alessandra Nicolosi, Tiffany Pézier, William Guesdon, Thierry Pedron,  
Fabrice Laurent, Sonia Lacroix-Lamandé

### ► To cite this version:

Alessandra Nicolosi, Tiffany Pézier, William Guesdon, Thierry Pedron, Fabrice Laurent, et al.. Role of Paneth cells during infection of neonatal mice by *Cryptosporidium parvum*. 7. International Giardia and Cryptosporidium Conference, Jun 2019, Rouen, France. , 243 p., 2019, 7. International Giardia and Cryptosporidium Conference. hal-02737527

**HAL Id: hal-02737527**

**<https://hal.inrae.fr/hal-02737527>**

Submitted on 2 Jun 2020

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# VII<sup>th</sup> International *Giardia* and *Cryptosporidium* Conference



**Conference Abstracts on USB Key**  
**June 23-26, 2019**



**UFR Santé, University of Rouen, France**

**Role of Paneth cells during infection of neonatal mice by *Cryptosporidium parvum*.**

Alessandra Nicolosi<sup>1</sup>, Tiffany Pezier<sup>1</sup>, William Guesdon<sup>1</sup>, Thierry Pedron<sup>2</sup>, Fabrice Laurent<sup>1</sup>, Sonia Lacroix-Lamandé<sup>1</sup>.

1- Equipe Apicomplexes et Immunité Mucosale (AIM), UMR1282 Infectiologie et Santé Publique (ISP) INRA – Université, Centre Val de Loire, 37380 Nouzilly. 2- Equipe Pathogénie Microbienne Moléculaire, Institut Pasteur de Paris.

*Cryptosporidium parvum* is a zoonotic apicomplexan parasite responsible for a diarrheal disease named cryptosporidiosis. This protozoan parasite is found worldwide and is transmitted by contaminated water. The immature intestinal immune system in very young animals and children under 5 places them at high risk of developing severe cryptosporidiosis. Paneth cells (PC) are specialized intestinal epithelial cells located at the base of intestinal crypts producing antimicrobial peptides (AMPs) that develop and mature after birth. We and others have already described in vitro that antimicrobial peptides such as CRAMP and CCL20 can alter the viability of sporozoites of *C. parvum* (1). We therefore wondered whether PCs and the AMPs that they produce can participate in the protective innate immune response against the parasite. By using a mouse model of neonatal cryptosporidiosis, we investigated the role of Paneth cells in the innate immune response against *C. parvum*. We first compared the susceptibility to *C. parvum* of mice genetically modified to be depleted of PCs (Sox9flox/flox-vil-Cre mice) and observed an increased level of infection when PCs are absent, associated with a reduced expression of AMPs. We also determined the effect of *Cryptosporidium parvum* infection on PC development and activity. By immunofluorescence, we observed on intestinal sections that *C. parvum* infection decreases the number of granule-positive-PCs and lysozyme-positive-PCs in neonatal mice. Altogether, these first results clearly demonstrate that PCs are important contributors of the innate protective immune response in mice and that lysozyme, already described to be efficient in vitro on *C. parvum* sporozoite viability, may be involved in this effect.

(1) Guesdon W, Auray G, Pezier T, Bussière FI, Drouet F, Le Vern Y, Marquis M, Potiron L, Rabot S, Bruneau A, Werts C, Laurent F, Lacroix-Lamandé S. CCL20 Displays Antimicrobial Activity Against *Cryptosporidium parvum*, but Its Expression Is Reduced During Infection in the Intestine of Neonatal Mice. J Infect Dis. 2015 .